

VA Influenza Toolkit Manual

2005 - 2006

Office of Public Health & Environmental Hazards
Employee Education System
National Center for Patient Safety
Infectious Diseases Program Office
VA Infection Control Professionals
National Center for Health Promotion and Disease Prevention

Veterans Health Administration
U.S. Department of Veterans Affairs

www.publichealth.va.gov/flu
www.publichealth.va.gov/InfectionDontPassItOn

September 2005



Foreword

Jonathan B. Perlin, MD, PhD, MSHA, FACP
Under Secretary for Health



Dear Colleagues,

The VA health care system has made influenza vaccination a priority and the results are evident. We have achieved self-reported vaccination rates of as high as 88% among eligible veteran patients over age 64—all of you deserve great credit for your efforts in achieving this milestone!

We must remember, however, that influenza is a serious and preventable illness that kills 36,000 Americans and leads to more than 200,000 excess hospitalizations annually. We know that influenza vaccine prevents influenza and the serious medical consequences of influenza quite well, and that as a Nation, we could do much, much better in our acceptance of this valuable annual preventive health measure.

We *must* continue to emphasize influenza vaccination of patients. Vaccinating our veteran patients protects them, helps protect their families, and keeps them from transmitting influenza to other patients and staff in our health care facilities.

What about ourselves? Health care worker vaccination rates are low—only about 40% of all U.S. health care workers report receiving influenza vaccine in any year. Yet, research has repeatedly shown that health care worker vaccination helps protect not only the vaccinated individual from illness, but prevents the spread of influenza to family members, patients, and other staff in the workplace. Vaccinated health care workers lose fewer days from their work and home lives to illness than unvaccinated health care workers. Influenza vaccine is safe and effective, and with all of us striving to make influenza vaccination easy and accessible, we *will* achieve VA staff vaccination rates equivalent to those of our veteran patients.

As I write this message to accompany the 2005-2006 Influenza Toolkit there are several uncertainties facing us regarding the upcoming influenza season. We don't know for sure what the influenza vaccine supply will be, what the demand will be, how severe the season will be, or which strain of influenza that will be most prevalent. Because of these uncertainties, I ask that you follow the guidelines and timing that will be communicated by my office through advisories and electronic mail throughout the influenza season.

Thank you for all of your dedication and your efforts for this year's influenza vaccination campaign.

A handwritten signature in blue ink, reading "Jonathan B. Perlin". The signature is fluid and cursive, written in a professional style.

Jonathan B. Perlin, MD, PhD, MSHA, FACP
July 2005

Introduction

Kristin Nichol, MD, MPH, MBA
Minneapolis VA Medical Center

Influenza and pneumococcal diseases are major causes of morbidity and mortality in the United States. Each year they are responsible for tens of millions of illnesses, hundreds of thousands of hospitalizations, tens of thousands of deaths, and billions of dollars in health care costs. Together they are responsible for more deaths due to vaccine preventable diseases by 10 to 50 fold than all other vaccine preventable diseases combined! The elderly and others with chronic medical conditions are at increased risk for the serious complications of influenza and pneumococcal diseases that might result in hospitalization and death, and many of the veterans we serve fall into one of these high risk categories.



Influenza and pneumococcal vaccinations are safe and effective. In the VA we have made influenza and pneumococcal vaccination a priority and have included monitoring vaccination rates in our performance measurement portfolio. Despite our impressive accomplishments to date, there is still room for improvement, especially among high risk veterans under the age of 65.

This year there is also increased emphasis on the importance of health care workers receiving their influenza vaccinations every year. Health care workers have been implicated as sources and/or vectors for influenza transmission in the health care setting, sometimes with serious complications for the patients who have contracted influenza from their health care provider. Nationally, only about 40% of health care workers report receiving their influenza vaccinations. It is likely in the VA system that we have similar vaccination rates.

We have much that we can and should to do to sustain past successes and improve further in our efforts to prevent and control influenza and pneumococcal diseases.

Evidence-based reviews of the literature have identified successful strategies for improving vaccination rates for elderly patients. These include:

- Providers' recommendations for vaccination.
- Systems approaches to ensure that patients are assessed and that appropriate vaccinations are offered and administered. Among such approaches are reminders to patients and providers, standing orders for nurses and other health care professionals, and walk-in clinics.

Keys to successfully increasing health care worker influenza vaccination rates are:

- Education about the disease, the safety of vaccination, and the rationale for targeting health care workers (first and foremost to protect our patients)
- Convenient access to vaccination (e.g. mobile carts, etc)
- Removal of cost barriers
- Top management buy-in

Improving influenza and pneumococcal vaccination rates will enhance the health and well being of our veterans and also reduce their health care costs. This toolkit will help you do just that.

Important Note on Influenza Vaccination

No two flu seasons are alike. Epidemics of influenza typically occur during the winter months in temperate regions and have been responsible for an average of 36,000 deaths/year in the United States. Influenza viruses also can cause pandemics, during which rates of illness and death from influenza-related complications can increase worldwide. Influenza viruses cause disease whose rates of infection are highest among persons aged ≥ 65 years, children aged < 2 years, and persons of any age who have medical conditions that place them at increased risk for complications from influenza. Influenza vaccination is the primary method for preventing influenza and its severe complications.

What happened last year?

The 2004-2005 flu season began with the announcement that a major U.S. supplier was unable to deliver its promised supply. It was anticipated that the national shortage would have significant impact on the Veterans Health

Administration (VHA), so seven advisories were written with essential input from colleagues across VHA. These contained information on policies, supplies, distribution, and

priority groups to receive vaccine. VA was able to respond to the initial influenza vaccine shortage by communicating priorities to patients and staff, establishing which of the CDC priority groups were vaccinated first, organizing clinics, and continuing the work of vaccinating late into the season.

VA Flu Advisories

VA staff and providers can review the latest information on 2005-2006 flu vaccine found in flu advisories on email and on the web by visiting www.publichealth.va.gov/flu.

It is important that all health care workers be vaccinated against flu unless they have a contraindication to the flu vaccine. Vaccinating for flu is one of the best ways to protect our entire VA community.

What's happening this year?

The Pharmacy Benefits Management (PBM) Strategic Health Care Group has announced that an award for Influenza Virus Vaccine (Fluzone) has been made for the 2005-2006 season. Contract # V797P-9170, was made to sanofi pasteur, inc., formerly Aventis Pasteur, Inc. This information has already been communicated to pharmacy staff throughout VHA, but it is important to reiterate it again here. As far as we know today, there may be only one approved manufacturer of influenza vaccine for the U.S.

As a health care system with a large percentage of patients in high priority groups, we need to communicate, follow guidance, and use our vaccine supplies fully and appropriately.

For 2005-2006 we do expect to contend with a vaccine supply challenge. Lessons from last year help us prepare for this year.

Sanofi pasteur will be splitting shipments into two or more deliveries. All ordering facilities will receive a partial shipment by October 15, 2005, with the balance to be completed by December 31, 2005.

Flu Advisory #1 for 2005 – 2006 flu season was released in May 2005. To keep informed on new flu information and to see the latest VA Flu Advisory, visit www.publichealth.va.gov/flu.

Contents

Section 1	Overview of This Year's Manual.....	11
Section 2	Vaccine Information.....	13
	<u>Inactivated Influenza Vaccine</u>	
	Information about Inactivated Influenza Vaccine.....	13
	CDC Inactivated Influenza Vaccine Information Statement (VIS).....	15
	<u>Live, Attenuated Intranasal Influenza Vaccine (LAIV)</u>	
	Guidance for Use in VA Workers, 2005-2006.....	17
	CDC Live, Intranasal Influenza Vaccine Information Statement (VIS).....	19
	<u>Pneumococcal Vaccine</u>	
	Information about Pneumococcal Vaccine.....	21
	CDC Pneumococcal Polysaccharide (PPV23) Vaccine Information.....	23
	Statement (VIS)	
Section 3.	How to Improve Health Care Worker Vaccination Rates.....	25
Section 4.	Best Strategies for Increasing Adult Influenza Vaccination Rates.....	29
Section 5.	Tools for Flu Prevention.....	33
Section 6.	Frequently Asked Questions.....	39
	General Questions.....	39
	Novel or Pandemic Influenza.....	41
	Eligibility for Influenza Vaccination in VA.....	41
	HIV/AIDS and Influenza Vaccination.....	42
	Influenza Vaccine Storage.....	43
	Live, attenuated, intranasal influenza vaccine (LAIV).....	44
Section 7.	Appendices.....	47
	A. How to Administer Flu and Pneumococcal Vaccines.....	49
	B. CDC Prevention and Control of Influenza. Recommendations of the Advisory Committee on Immunization Practices (ACIP). MMWR. July 29, 2005 / 54....	51
	C. CDC Adult Immunization Schedule, 2004-2005 [2005-2006 to be released in fall 2005].....	97
	D. Prevention and Treatment of Influenza with Antiviral Drugs.....	103
	E. Resources, References, and Web sites.....	105
	F. Acknowledgements.....	111

Section 1

Overview of This Year's Manual

This year's *VA Influenza Toolkit Manual* contains basic background information and resources designed to help VA staff carry out a successful program of vaccination for veterans and staff in the VA medical system.

The 2005-2006 Influenza Toolkit

Printed versions of the entire toolkit (this manual and other materials) are being sent to eight groups of key VA staff in each facility:

- Influenza vaccination (“flu”) coordinators
- Occupational health clinicians
- Prevention coordinators
- Infection control professionals
- Public affairs officers
- Patient education contacts
- Patient safety officers
- Education contacts (Employee Education System)

Goals for the vaccination program

- Increasing our rate of influenza vaccination of health care workers – keeping those with patient contact healthy is an essential aspect of keeping our medical system running and taking good care of patients. On the average, only about 40% of health care workers in the US get vaccinated against the flu – VA can do better.
- Maintaining or even increasing our already excellent rate of influenza vaccination of veteran patients. Some of our medical centers are reporting as many as 9 out of 10 eligible patients get a flu shot. VA can keep up the great work and make sure as many eligible veterans as possible are protected against the flu.
- Providing vaccination against pneumococcal illness, when appropriate.
- Promoting nonvaccine methods of preventing infection, particularly through hand and respiratory hygiene.
- Asking all VHA staff and providers, even those who do not directly deliver vaccinations, to join this year's flu vaccinations efforts by keeping informed of vital flu information, discussing influenza with patients on the importance of getting vaccinated, hanging posters, and supporting “flu” coordinators and infection control professionals with flu prevention efforts during influenza season.

Section 2 of the manual provides fact sheets or guidance for both the inactivated and live forms of influenza vaccines as well as pneumococcal vaccine. It also provides reproducible copies of the most current Vaccine Information Statements (VISs) available from the Centers for Disease Control and Prevention (CDC). A VIS should be provided to each person receiving a given vaccine.

Section 3 was written by VHA's Occupational Health Program to support this year's emphasis on health care worker vaccination and gives practical information on improving rates of vaccination, what the barriers are, and suggestions for overcoming these barriers.

Section 4 is a compilation of material from published studies as well as feedback from VA staff on strategies and ideas for carrying out successful vaccination programs in VA.

Section 5 of the manual is an overview of the tools in the 2005 – 2006 Flu Toolkit (posters, stickers, and buttons) that were created separately from this manual and offers suggestions on how to use them.

Section 6 covers a variety of frequently asked questions that VA staff may have concerning influenza, vaccination, and other aspects of this year's flu vaccination program.

Section 7 contains the **Appendices**. These include an explanation of the steps involved in vaccinating individuals, reprints of the latest CDC guidance on influenza and adult immunization schedule, information on prevention and treatment of influenza with antiviral drugs, a long list of additional sources of information, and the list of coordinating offices and staff that assembled this material.

All of this material – and any changes that come about over the course of this flu season – will be posted on the Web at www.publichealth.va.gov/flu and www.publichealth.va.gov/InfectionDontPassItOn (and the VA Intranet sites <http://vaww.vhaco.va.gov/phshcg/Flu> and <http://vaww.vhaco.va.gov/phshcg/InfectionDontPassItOn>).

Section 2

Vaccine Information

Inactivated (Injectable) Influenza Vaccine 2005-2006

- **The 2005-06 trivalent influenza vaccine for the United States will contain:**
 - A/New Caledonia/20/99-like (H1N1)
 - A/California/7/2004-like (H3N2) or A/New York/55/2004
 - B/Shanghai/361/2002-like antigens or B/Jilin/20/2003 or B/Jiangsu/10/2003
- **What’s new this year?**
 - An increased emphasis on health care worker vaccination—*annually for all*
 - Persons with compromised respiratory functioning, compromised handling of respiratory secretions, or at-risk for aspiration are added to the high risk category
 - Vaccine programs should employ both forms of influenza vaccine—inactivated (injectable) and live attenuated (intranasal) influenza vaccine in order to extend the supply available for persons only eligible to receive the inactivated form.
 - The possibility that vaccine may need to be given to members of risk groups in order of their risk from influenza, a *tiered timing* approach, will be announced as the vaccine supply is better defined.

- **Who should be vaccinated?**

*The Advisory Committee on Immunization Practices (ACIP) and CDC recommends influenza vaccination for people at high risk for complications from influenza, or who are important for the care of infants, are health care workers or who are in an age group where many members are at high risk . People in the following groups should seek vaccination this season, although they may be asked to follow a *tiered timing* approach announced by CDC and the VA Under Secretary for Health thru advisories and emails.*

Persons at high risk from influenza

- adults aged 65 years and older;
- residents of nursing homes and long-term care facilities
- persons aged 2–64 years with underlying chronic medical conditions¹;
- children 6 months-18 years who are on long-term aspirin therapy
- all children aged 6–23 months
- all women who will be pregnant during the influenza season

Persons from an age group with members at high risk from influenza

- persons age 50-64

Persons who may transmit influenza, are important in the care of infants, or who work in health care settings

- health-care workers
- out-of-home caregivers and household contacts of children aged <6 months.

*Persons who are not in any of the groups listed above are also encouraged to be vaccinated with inactivated vaccine as supplies allow. These persons should follow *tiered timing* guidance or get live, inactivated (intranasal) influenza vaccine if eligible.*

¹ for a complete list of chronic medical conditions see “Target Groups for Vaccination/Persons at Increased Risk for Complications” in Prevention and Control of Influenza: Recommendations of the Advisory Committee on Immunization Practices (ACIP), MMWR, July 29, 2005. Vol. 54. Available at <http://www.cdc.gov/mmwr/preview/mmwrhtml/rr5408a1.htm>

INACTIVATED INFLUENZA VACCINE

WHAT YOU NEED TO KNOW

1 Why get vaccinated?

Influenza (“flu”) is a very contagious disease.

It is caused by the influenza virus, which spreads from infected persons to the nose or throat of others.

Other illnesses can have the same symptoms and are often mistaken for influenza. But only an illness caused by the influenza virus is really influenza.

Anyone can get influenza. For most people, it lasts only a few days. It can cause:

- fever
- sore throat
- chills
- fatigue
- cough
- headache
- muscle aches

Some people get much sicker. Influenza can lead to pneumonia and can be dangerous for people with heart or breathing conditions. It can cause high fever and seizures in children. Influenza kills about 36,000 people each year in the United States, mostly among the elderly.

Influenza vaccine can prevent influenza.

2 Inactivated Influenza vaccine

There are two types of influenza vaccine:

An **inactivated** (killed) vaccine, given as a shot, has been used in the United States for many years.

A **live**, weakened vaccine was licensed in 2003. It is sprayed into the nostrils. *This vaccine is described in a separate Vaccine Information Statement.*

Influenza viruses are constantly changing. Therefore, influenza vaccines are updated every year, and an annual vaccination is recommended.

For most people influenza vaccine prevents serious illness caused by the influenza virus. It will *not* prevent “influenza-like” illnesses caused by other viruses.

It takes about 2 weeks for protection to develop after the shot, and protection can last up to a year.

Inactivated influenza vaccine may be given at the same time as other vaccines, including pneumococcal vaccine.

Some inactivated influenza vaccine contains thimerosal, a preservative that contains mercury. Some people believe thimerosal may be related to developmental problems in children. In 2004 the Institute of Medicine published a report concluding that, based on scientific studies, there is no evidence of such a relationship. If you are concerned about thimerosal, ask your doctor about thimerosal-free influenza vaccine.

3 Who should get inactivated influenza vaccine?

Influenza vaccine can be given to people 6 months of age and older. It is recommended for **people who are at risk of serious influenza or its complications**, and for **people who can spread influenza to those at high risk** (including all household members):

People at high risk for complications from influenza:

- **All children** 6-23 months of age.
- **People 65 years of age and older.**
- Residents of **long-term care facilities** housing persons with chronic medical conditions.
- People who have **long-term health problems** with:
 - heart disease
 - kidney disease
 - lung disease
 - metabolic disease, such as diabetes
 - asthma
 - anemia, and other blood disorders
- People with certain conditions (such as **neuromuscular disorders**) that can cause breathing problems.
- People with a **weakened immune system** due to:
 - HIV/AIDS or other diseases affecting the immune system
 - long-term treatment with drugs such as steroids
 - cancer treatment with x-rays or drugs
- People 6 months to 18 years of age on **long-term aspirin treatment** (these people could develop Reye Syndrome if they got influenza).
- Women who will be **pregnant** during influenza season.

People who can spread influenza to those at high risk:

- **Household contacts and out-of-home caretakers** of infants from 0-23 months of age.
- Physicians, nurses, family members, or anyone else in **close contact with people at risk** of serious influenza.

Influenza vaccine is also recommended for adults 50-64 years of age and anyone else who wants to **reduce their chance of catching influenza.**

An annual flu shot should be *considered* for:

- People who provide **essential community services.**
- People living in **dormitories** or under other crowded conditions, to prevent outbreaks.
- People at high risk of influenza complications who **travel** to the Southern hemisphere between April and September, or to the tropics or in organized tourist groups at **any** time.

4

When should I get influenza vaccine?

The best time to get influenza vaccine is in **October** or **November**.

Influenza season usually peaks in February, but it can peak any time from November through May. So getting the vaccine in December, or even later, can be beneficial in most years.

Some people should get their flu shot in **October** or earlier:

- people **50 years of age and older**,
- younger people at **high risk** from influenza and its complications (including **children 6 through 23 months of age**),
- **household contacts** of people at high risk,
- **healthcare workers**, and
- **children younger than 9 years of age** getting influenza vaccine for the first time.

Most people need one flu shot each year. **Children younger than 9 years of age getting influenza vaccine for the first time** should get 2 doses, given at least one month apart.

5

Some people should talk with a doctor before getting influenza vaccine

Some people should not get inactivated influenza vaccine or should wait before getting it.

- Tell your doctor if you have any **severe** (life-threatening) allergies. Allergic reactions to influenza vaccine are rare.
 - Influenza vaccine virus is grown in eggs. People with a severe egg allergy should not get the vaccine.
 - A severe allergy to any vaccine component is also a reason to not get the vaccine.
 - If you have had a severe reaction after a previous dose of influenza vaccine, tell your doctor.
- Tell your doctor if you ever had Guillain-Barré Syndrome (a severe paralytic illness, also called GBS). You may be able to get the vaccine, but your doctor should help you make the decision.
- People who are moderately or severely ill should usually wait until they recover before getting flu vaccine. If you are ill, talk to your doctor or nurse about whether to reschedule the vaccination. People with a **mild illness** can usually get the vaccine.

6

What are the risks from inactivated influenza vaccine?

A vaccine, like any medicine, could possibly cause serious problems, such as severe allergic reactions. The risk of a vaccine causing serious harm, or death, is extremely small.

Serious problems from influenza vaccine are very rare. The viruses in inactivated influenza vaccine have been killed, so you cannot get influenza from the vaccine.

Mild problems:

- soreness, redness, or swelling where the shot was given
- fever
- aches

If these problems occur, they usually begin soon after the shot and last 1-2 days.

Severe problems:

- Life-threatening allergic reactions from vaccines are very rare. If they do occur, it is within a few minutes to a few hours after the shot.
- In 1976, a certain type of influenza (swine flu) vaccine was associated with Guillain-Barré Syndrome (GBS). Since then, flu vaccines have not been clearly linked to GBS. However, if there is a risk of GBS from current flu vaccines, it would be no more than 1 or 2 cases per million people vaccinated. This is much lower than the risk of severe influenza, which can be prevented by vaccination.

7

What if there is a severe reaction?

What should I look for?

- Any unusual condition, such as a high fever or behavior changes. Signs of a serious allergic reaction can include difficulty breathing, hoarseness or wheezing, hives, paleness, weakness, a fast heart beat or dizziness.

What should I do?

- **Call** a doctor, or get the person to a doctor right away.
- **Tell** your doctor what happened, the date and time it happened, and when the vaccination was given.
- **Ask** your doctor, nurse, or health department to report the reaction by filing a Vaccine Adverse Event Reporting System (VAERS) form.

Or you can file this report through the VAERS web site at www.vaers.hhs.gov, or by calling 1-800-822-7967.

VAERS does not provide medical advice.

8

The National Vaccine Injury Compensation Program

In the event that you or your child has a serious reaction to a vaccine, a federal program has been created to help pay for the care of those who have been harmed.

For details about the National Vaccine Injury Compensation Program, call 1-800-338-2382 or visit their website at www.hrsa.gov/osp/vicp

9

How can I learn more?

- Ask your immunization provider. They can give you the vaccine package insert or suggest other sources of information.
- Call your local or state health department.
- Contact the Centers for Disease Control and Prevention (CDC):
 - Call 1-800-232-4636 (1-800-CDC-INFO)
 - Visit CDC's website at www.cdc.gov/flu



DEPARTMENT OF HEALTH AND HUMAN SERVICES
CENTERS FOR DISEASE CONTROL AND PREVENTION
NATIONAL IMMUNIZATION PROGRAM

Vaccine Information

Live, Attenuated Intranasal Influenza Vaccine (LAIV) Guidance for Use in VA Workers 2005-2006

For VA Health Care Workers, Other VA Employees, Students, Trainees, and Volunteers

- A single LAIV is licensed in the U.S.—FluMist™ (MedImmune, Inc.). LAIV is a live, trivalent, intranasally-administered vaccine that induces broad mucosal and systemic immune response.
- **LAIV is an option for vaccinating healthy VA employees under the age of 50.** VA health care facilities may use it whether or not there is a shortage of inactivated (injectable) vaccine. But, especially in the event of a shortage of inactivated vaccine, use of LAIV conserves inactivated vaccine for those are not eligible to receive LAIV.
- Side effects that may occur after administration of LAIV to adults include runny nose, nasal congestion, headache, sore throat, and cough.
- **LAIV should NOT be given**
 - to VA health care workers and employees who are 50 or over
 - persons who are at high risk of influenza complications (including all persons ≥ 65 years old; pregnant women; the immunocompromised; those who have asthma, cystic fibrosis, or chronic obstructive pulmonary disease; those with chronic metabolic disease like diabetes; those with renal disease).
- **LAIV should NOT be administered by**
 - severely immunosuppressed persons because of the small risk of acquiring vaccine virus from the environment during administration.
- **LAIV MAY be administered by**
 - others considered at high risk of influenza complications (including persons ≥ 65 years old, pregnant women, those who have asthma, cystic fibrosis, or chronic obstructive pulmonary disease; those with chronic metabolic disease like diabetes; those with renal disease).
- **Restrictions from working after receiving LAIV:**
Shedding of vaccine virus occurs, mostly in the first 3 days after administration. One unpublished study reported transmission of one of the vaccine component viruses from one child to another in a daycare setting, but the transmission did not result in clinical illness in the recipient child. Person to person transmission of vaccine virus in adults has not been reported. After 2 seasons of safe use of LAIV, concern about transmission of vaccine virus has diminished.
 - ***Employees who work with the severely immunosuppressed in a protective environment should not get LAIV*** (e.g. patients who are in hospital in a protective environment after a hematopoietic stem cell transplant). If an employee who works with severely immunosuppressed patients in a protective environment receives LAIV, the employee should refrain from contact with severely immunosuppressed patients for 7 days.

Exclusion from the workplace after receiving LAIV is otherwise not necessary.

- **LAIV is a useful alternative to inactivated influenza vaccine for some VA workers.** Influenza vaccination protects VA workers and those around them from influenza illness. All VA workers (unless they have a contraindication) should receive an annual influenza vaccination.

References:

VA Influenza Directive 05-06 (pending)

Prevention and Control of Influenza: Recommendations of the Advisory Committee on Immunization Practices (ACIP), MMWR, July 29, 2005. Vol. 54. Available at <http://www.cdc.gov/mmwr/preview/mmwrhtml/rr5408a1.htm>

LIVE, INTRANASAL INFLUENZA VACCINE

WHAT YOU NEED TO KNOW

1 Why get vaccinated?

Influenza (“flu”) is a very contagious disease.

It is caused by the influenza virus, which spreads from infected persons to the nose or throat of others.

Other illnesses can have the same symptoms and are often mistaken for influenza. But only an illness caused by the influenza virus is really influenza.

Anyone can get influenza, but rates of infection are highest among children. For most people, it lasts only a few days. It can cause:

· fever · sore throat · chills · fatigue
· cough · headache · muscle aches

Some people get much sicker. Influenza can lead to pneumonia and can be dangerous for people with heart or breathing conditions. It can cause high fever and seizures in children. Influenza kills about 36,000 people each year in the United States.

Influenza vaccine can prevent influenza.

2 Live, attenuated influenza vaccine (nasal spray)

There are two types of influenza vaccine:

Live, attenuated influenza vaccine (LAIV) was licensed in 2003. LAIV contains live but attenuated (weakened) influenza virus. It is sprayed into the nostrils rather than injected into the muscle. It is recommended for healthy children and adults from 5 through 49 years of age, who are not pregnant.

Inactivated influenza vaccine, sometimes called the “flu shot,” has been used for many years and is given by injection. *This vaccine is described in a separate Vaccine Information Statement.*

Influenza viruses are constantly changing. Therefore, influenza vaccines are updated every year, and annual vaccination is recommended.

For most people influenza vaccine prevents serious illness caused by the influenza virus. It will *not* prevent “influenza-like” illnesses caused by other viruses.

It takes about 2 weeks for protection to develop after vaccination, and protection can last up to a year.

3 Who can get LAIV?

Live, intranasal influenza vaccine is approved for healthy children and adults from 5 through 49 years of age, including most health-care workers and household contacts of most people at high risk for influenza complications. However, LAIV should not be given to pregnant women or people with certain medical conditions.

4 Who should *not* get LAIV?

The following people should not get live intranasal influenza vaccine. They should check with their health-care provider about getting the inactivated vaccine.

- **Adults 50 years of age or older or children younger than 5.**
- People who have **long-term health problems** with:
 - heart disease
 - kidney disease
 - lung disease
 - metabolic disease, such as diabetes
 - asthma
 - anemia, and other blood disorders
- People with a **weakened immune system** due to:
 - HIV/AIDS or other diseases affecting the immune system
 - long-term treatment with drugs that weaken the immune system, such as steroids
 - cancer treatment with x-rays or drugs
- Children or adolescents on **long-term aspirin treatment** (these people could develop Reye syndrome if they get influenza).
- **Pregnant women.**
- Anyone with a history of **Guillain-Barré syndrome** (a severe paralytic illness, also called GBS).

Inactivated influenza vaccine (the flu shot) is the preferred vaccine for people (including health-care workers, and family members) coming in **close contact with anyone who has a severely weakened immune system** (that is, anyone who requires care in a protected environment).

Some people should talk with a doctor before getting *either* influenza vaccine:

- Anyone who has ever had a serious allergic reaction to **eggs** or to a **previous dose** of influenza vaccine.
- People who are moderately or severely ill should usually wait until they recover before getting flu vaccine. If you are ill, talk to your doctor or nurse about whether to reschedule the vaccination. People with a **mild illness** can usually get the vaccine.

5 When should I get influenza vaccine?

The best time to get influenza vaccine is in **October** or **November**. Influenza season usually peaks in February, but it can peak any time from November through May. So getting the vaccine in December, or even later, can be beneficial in most years.

Most people need one dose of influenza vaccine each year. **Children younger than 9 years of age getting influenza vaccine for the first time** should get 2 doses for LAIV, these doses should be given 6-10 weeks apart.

LAIV may be given at the same time as other vaccines. This includes other live vaccines, such as MMR or chickenpox. But if two live vaccines are not given on the same day, they should be given at least 4 weeks apart.

6 What are the risks from LAIV?

A vaccine, like any medicine, could possibly cause serious problems, such as severe allergic reactions. However, the risk of a vaccine causing serious harm, or death, is extremely small.

Live influenza vaccine viruses rarely spread from person to person. Even if they do, they are not likely to cause illness.

LAIV is made from weakened virus and does not cause influenza. The vaccine *can* cause mild symptoms in people who get it (see below).

Mild problems:

Some children and adolescents 5-17 years of age have reported mild reactions, including:

- runny nose, nasal congestion or cough
- headache and muscle aches
- fever
- abdominal pain or occasional vomiting or diarrhea

Some adults 18-49 years of age have reported:

- runny nose or nasal congestion
- sore throat
- cough, chills, tiredness/weakness
- headache

These symptoms did not last long and went away on their own. Although they can occur after vaccination, they may not have been caused by the vaccine.

Severe problems:

- Life-threatening allergic reactions from vaccines are very rare. If they do occur, it is within a few minutes to a few hours after the vaccination.

- If rare reactions occur with any new product, they may not be identified until thousands, or millions, of people have used it. Over two million doses of LAIV have been distributed since it was licensed, and no serious problems have been identified. Like all vaccines, LAIV will continue to be monitored for unusual or severe problems.

7 What if there is a severe reaction?

What should I look for?

- Any unusual condition, such as a high fever or behavior changes. Signs of a serious allergic reaction can include difficulty breathing, hoarseness or wheezing, hives, paleness, weakness, a fast heart beat or dizziness.

What should I do?

- **Call** a doctor, or get the person to a doctor right away.
- **Tell** your doctor what happened, the date and time it happened, and when the vaccination was given.
- **Ask** your doctor, nurse, or health department to report the reaction by filing a Vaccine Adverse Event Reporting System (VAERS) form.

Or you can file this report through the VAERS website at www.vaers.hhs.gov, or by calling 1-800-822-7967.

VAERS does not provide medical advice.

8 The National Vaccine Injury Compensation Program

In the event that you or your child has a serious reaction to a vaccine, a federal program has been created to help pay for the care of those who have been harmed.

For details about the National Vaccine Injury Compensation Program, call **1-800-338-2382** or visit their website at www.hrsa.gov/osp/vicp

9 How can I learn more?

- Ask your immunization provider. They can give you the vaccine package insert or suggest other sources of information.
- Call your local or state health department.
- Contact the Centers for Disease Control and Prevention (CDC):
 - Call **1-800-232-4636 (1-800-CDC-INFO)**
 - Visit CDC's website at www.cdc.gov/flu



**DEPARTMENT OF HEALTH AND HUMAN SERVICES
CENTERS FOR DISEASE CONTROL AND PREVENTION
NATIONAL IMMUNIZATION PROGRAM** 20

Vaccine Information

Information about Pneumococcal Vaccine for 2005-2006

Pneumococcal vaccine (Pneumococcal Polysaccharide Vaccine, PPV 23) is used to decrease the risk of serious pneumococcal disease and its complications.

It may be administered to adults *at any time during the year* who meet any of the criteria or have conditions listed below:

- Age 65 or older
- HIV infection or AIDS
- Absent or malfunctioning spleen
- Sickle cell disease
- Organ or bone marrow transplant candidate or recipient
- Immunosuppressive treatment with X-ray, cancer drugs, or long-term steroids
- Persons living in special environments or social situations (such as Native Americans or residents of long term care facilities)
- Nephrotic syndrome or renal failure
- Candidate for or recipient of a cochlear implant
- Lung disease
- Cancer, including leukemia, lymphoma, and multiple myeloma
- Diabetes mellitus
- Spinal cord injury or disease
- Cerebrospinal fluid leaks
- Alcoholism or cirrhosis of the liver

How often should pneumococcal vaccine be given?

- *Once* to adults age 65 or older if they have not received an earlier dose
- Adults who received a dose before age 65 should receive a 2nd dose if 5 or more years have passed since the first dose.
- Adults at the highest risk of pneumococcal infections should receive a 2nd dose 5 or more years after the first dose, regardless of what age the 1st dose was given. Adults at the highest risk are those with: HIV infection or AIDS, absent or malfunctioning spleen, sickle cell disease, organ or bone marrow recipients, nephrotic syndrome or renal failure, immunosuppressive treatment with X-rays, cancer drugs, or long-term steroids.
- Only 2 doses at most are given. Repeating doses every 5, 6, or 10 years is not recommended.

What about repeating a dose if a patient is uncertain of having received it before?

If the patient's vaccination status is unknown, those in the recommended group should be administered pneumococcal vaccine. Studies have shown that adults who were revaccinated 4 years or more after an initial vaccination did not have an increased incidence of side effects. Consider supplying patients who have trouble remembering their vaccination history with a personal immunization card. Examples of cards are available at <https://www.immunize.org/adultizcards/index.htm>

PNEUMOCOCCAL POLYSACCHARIDE VACCINE

WHAT YOU NEED TO KNOW

1 Why get vaccinated?

Pneumococcal disease is a serious disease that causes much sickness and death. In fact, pneumococcal disease kills more people in the United States each year than all other vaccine-preventable diseases combined. Anyone can get pneumococcal disease. However, some people are at greater risk from the disease. These include people 65 and older, the very young, and people with special health problems such as alcoholism, heart or lung disease, kidney failure, diabetes, HIV infection, or certain types of cancer.

Pneumococcal disease can lead to serious infections of the lungs (pneumonia), the blood (bacteremia), and the covering of the brain (meningitis). About 1 out of every 20 people who get pneumococcal pneumonia dies from it, as do about 2 people out of 10 who get bacteremia and 3 people out of 10 who get meningitis. People with the special health problems mentioned above are even more likely to die from the disease.

Drugs such as penicillin were once effective in treating these infections; but the disease has become more resistant to these drugs, making treatment of pneumococcal infections more difficult. This makes prevention of the disease through vaccination even more important.

2 Pneumococcal polysaccharide vaccine (PPV)

The pneumococcal polysaccharide vaccine (PPV) protects against 23 types of pneumococcal bacteria. Most healthy adults who get the vaccine develop protection to most or all of these types within 2 to 3 weeks of getting the shot. Very old people, children under 2 years of age, and people with some long-term illnesses might not respond as well or at all.

3 Who should get PPV?

- All adults 65 years of age or older.
- Anyone over 2 years of age who has a long-term health problem such as:
 - heart disease
 - lung disease
 - sickle cell disease
 - diabetes
 - alcoholism
 - cirrhosis
 - leaks of cerebrospinal fluid
- Anyone over 2 years of age who has a disease or condition that lowers the body's resistance to infection, such as:
 - Hodgkin's disease
 - lymphoma, leukemia
 - kidney failure
 - multiple myeloma
 - nephrotic syndrome
 - HIV infection or AIDS
 - damaged spleen, or no spleen
 - organ transplant
- Anyone over 2 years of age who is taking any drug or treatment that lowers the body's resistance to infection, such as:
 - long-term steroids
 - certain cancer drugs
 - radiation therapy
- Alaskan Natives and certain Native American populations.

4

How many doses of PPV are needed?

Usually one dose of PPV is all that is needed.

However, under some circumstances a second dose may be given.

- A second dose is recommended for those people aged 65 and older who got their first dose when they were under 65, if 5 or more years have passed since that dose.
- A second dose is also recommended for people who:
 - have a damaged spleen or no spleen
 - have sickle-cell disease
 - have HIV infection or AIDS
 - have cancer, leukemia, lymphoma, multiple myeloma
 - have kidney failure
 - have nephrotic syndrome
 - have had an organ or bone marrow transplant
 - are taking medication that lowers immunity (such as chemotherapy or long-term steroids)

Children 10 years old and younger may get this second dose 3 years after the first dose. Those older than 10 should get it 5 years after the first dose.

5

Other facts about getting the vaccine

- Otherwise healthy children who often get ear infections, sinus infections, or other upper respiratory diseases do not need to get PPV because of these conditions.
- PPV may be less effective in some people, especially those with lower resistance to infection. But these people should still be vaccinated, because they are more likely to get seriously ill from pneumococcal disease.
- **Pregnancy:** The safety of PPV for pregnant women has not yet been studied. There is no evidence that the vaccine is harmful to either the mother or the fetus, but pregnant women should consult with their doctor before being vaccinated. Women who are at high risk of pneumococcal disease should be vaccinated before becoming pregnant, if possible.

6

What are the risks from PPV?

PPV is a very safe vaccine.

About half of those who get the vaccine have very mild side effects, such as redness or pain where the shot is given.

Less than 1% develop a fever, muscle aches, or more severe local reactions.

Severe allergic reactions have been reported very rarely.

As with any medicine, there is a very small risk that serious problems, even death, could occur after getting a vaccine.

Getting the disease is much more likely to cause serious problems than getting the vaccine.

7

What if there is a serious reaction?

What should I look for?

- Severe allergic reaction (hives, difficulty breathing, shock).

What should I do?

- **Call** a doctor, or get the person to a doctor right away.
- **Tell** your doctor what happened, the date and time it happened, and when the vaccination was given.
- **Ask** your doctor, nurse, or health department to report the reaction by filing a Vaccine Adverse Event Reporting System (VAERS) form.

Or you can file this report through the VAERS web site at www.vaers.org, or by calling 1-800-822-7967.

VAERS does not provide medical advice.

8

How can I learn more?

- Ask your doctor or nurse. They can give you the vaccine package insert or suggest other sources of information.
- Call your local or state health department.
- Contact the Centers for Disease Control and Prevention (CDC):
 - Call **1-800-232-4636 (1-800-CDC-INFO)** or
 - Visit the National Immunization Program website at www.cdc.gov/nip



**DEPARTMENT OF HEALTH AND HUMAN SERVICES
CENTERS FOR DISEASE CONTROL AND PREVENTION
NATIONAL IMMUNIZATION PROGRAM**

Section 3

Improving Health Care Worker Vaccination Rates

CDC recommends that all health care workers receive an annual influenza vaccination to prevent transmission to patients. Influenza immunization rates among health care workers remain low, with only 36%-40% of health care workers reporting influenza vaccination each year. Influenza vaccination is an important patient safety issue because unvaccinated staff can spread influenza to patients, coworkers, and family members, leading to influenza-related illness and death. When health care workers become ill with influenza, absenteeism and disruption of care may result. When health care workers transmit influenza to patients, some of them may experience serious, even life-threatening complications or secondary pneumonias. When promoting vaccination among health care workers, emphasize the reasons to get the influenza vaccine:

- Prevents death
- Prevents severe illness
- Protects patients
- Protects families
- Protects you and your coworkers
- Decreases use of sick leave

Obstacles – Individual Beliefs

Some health care workers have misperceptions and misunderstandings about influenza vaccine. The scientific literature suggests several reasons for low immunization rates among health care workers including: concern about side effects or vaccine safety, perception of low personal risk of illness, inconvenience, ignorance of CDC recommendations, and dislike of needles. Therefore, there should be continuous and ongoing vaccine education updates emphasizing the seriousness of influenza and addressing employee misconceptions about influenza and the vaccine. Common misconceptions must be addressed (the flushot does NOT give you influenza). Examples of notices that might be posted include:

You know that the influenza shot works, so why don't more people get vaccinated?

Some people are concerned about side effects. They think that the influenza shot will make them sick. However, mild soreness of the arm at the injection site is the most common side effect. The shot itself will NOT give you influenza. Influenza vaccination is the best protection against the influenza. Protect VA patients, yourself, your co-workers and your family. Get vaccinated. Check with your supervisor for information on how to get your influenza shot.

Did you get your influenza shot last year?

If you didn't, you may have harmed the health of some of our patients, your coworkers and family members. You can spread influenza to patients, putting them at risk for influenza and its complications. Studies show that vaccination of health care workers is associated with decreased mortality among nursing home patients. Protect yourself and your patients--get a shot. Ask your supervisor about information on how to get a flu shot.

“I don't need to get a flu shot.” Is this you?

Influenza can cause serious illness and death even in young, healthy people. It's not just a disease that affects the elderly. If you get influenza, you can spread it to your patients, putting them at risk for severe illness and complications from the flu. Protect yourself, your coworkers, and your patients—get a flu shot. Ask your supervisor for information on how to get vaccinated.

Successful Campaigns for Vaccinating Health Care Workers

- ➔ Sending a letter, postcard or email to employees prior to the start of the vaccine season reminding them of the importance of vaccination, where and when they will be able to get the flu vaccine.
- ➔ Writing something in the employee newsletter or posting information on staff bulletin boards and providing fact sheets with pay stubs to dispel misconceptions and increase acceptance.
- ➔ Increasing the number of sites where the vaccine is given. Use mobile carts to transport to different clinic areas, service meetings, grand rounds or near cafeteria entrances. This approach can minimize inconvenience as well as means to advertise the vaccine availability. Carts should be stocked with vaccine, safety syringes, vaccine information statements, sharps disposal containers, alcohol hand rub, alcohol wipes, adhesive bandages, documentation forms, and injectable epinephrine with orders for administration in the event of an acute hypersensitivity reaction.
- ➔ Making appointments with services to attend service meetings. A schedule should be posted or email sent to those in the service announcing that the vaccine is available at the staff meeting.
- ➔ If your occupational health unit has a website, adding information to the website regarding flu shot locations and times.
- ➔ In late November identifying employees not yet vaccinated and reminding them by email or a phone call that the flu vaccine is available.
- ➔ Working closely with Pharmacy to get your supply of vaccine for employees.
- ➔ Modifying education materials from CDC and elsewhere to address risks to employees if they are not vaccinated.
- ➔ Encouraging the facility director, service chiefs, and other managers to lead the way by getting their vaccine and encouraging their staff to get vaccinated.
- ➔ Giving out buttons or stickers to all staff who are vaccinated showing that they have been vaccinated. It is an additional advertising strategy for both employees and patients to be vaccinated against influenza.
- ➔ Sponsoring a kick off event.
- ➔ Adding an influenza reminder to occupational health's telephone recording. When employees call they can automatically be reminded about the availability of the vaccine. If the recording capacity exists, add specific information regarding dates, times and locations for flu shots as well as any other pertinent information. These reminders should begin September 15 and conclude after the flu season has peaked, which usually occurs in February or March.
- ➔ Extending hours that the vaccine is available to staff to include all shifts and days of the week. Plans must be made to have additional staff available during the extended hours of the clinic or available during off hours.
- ➔ If there is a vaccine shortage, using additional strategies if necessary to ensure those who are identified as needing the vaccine are targeted. If the shortage resolves, there also should be a mechanism in place to remind those not vaccinated that it is not too late to get the vaccine.

Key elements of a successful staff flu vaccination campaign include:

- informing employees about the availability of the vaccine and the plan (awareness),
- educating employees about its importance (marketing),
- making the vaccine convenient,
- notifying employees regarding the scheduling of administration and
- keeping track of who has been vaccinated.

Additional Measures to Take to Prevent the Spread of Influenza

Remind employees that although the influenza vaccination may be the best way to protect against the influenza, there are other measure that they should also take to protect themselves, their families, and patients.

- Stay at home when they are sick, especially if running a fever
- Keep tissues at their desk so that they can cough or sneeze into it – NOT into their hands
- Keep a trash can at their desk to dispose of used tissues
- Frequently wipe down their keyboard, mouse and phone with antibacterial wipes
- Wash their hands or wipe with hand sanitizer frequently, especially after using copy machines, fax machines, someone else's computer or phone, or after sneezing or other contact with their own secretions
- Avoid contact with people who are sick, except of course the patients they are here to help
- Wash their hands before eating food
- Wash their hands frequently with water and soap or alcohol-based rubs

Remember health care workers may also have health problems and conditions which put them at increased risk of complications from influenza. These include:

- Chronic cardiac or pulmonary disorders severe enough to require regular medical follow-up care.
- Being 65 and older.
- Chronic health conditions such as diabetes mellitus and other metabolic diseases, cancer, immunodeficiency, renal disease, anemia and hemoglobinopathy.
- Any conditions that can compromise respiratory function or the handling of respiratory secretions or that can increase the risk of aspiration.
- Pregnant women.

Immunization is the primary method to prevent influenza, limit transmission, and prevent complications from the flu. Trivalent inactivated influenza virus vaccine may be administered to all categories of health care workers unless there is a contraindication for the vaccine. Live attenuated influenza vaccine (FluMist™) may also be administered to health care workers. It is a good option for those health care workers who are in good health, are not pregnant, those who have a dislike of needles, and meet the criteria for LAIV.

Section 4

Best Strategies for Increasing Adult Influenza Vaccination Rates

These strategies have been shown to be effective for increasing adult influenza immunization rates. They are most effective when combined in multi-component efforts to increase immunizations. In the event of a vaccine shortage, it is important that patients at highest risk for complications from influenza and the health care workers who provide direct service to them be vaccinated. Patient and health care worker strategies are described separately, and more details can be obtained from Appendix E.

Getting Patients Vaccinated

1. Organizational
 - Make influenza vaccination an organizational priority
 - Provide written policy stressing importance and effectiveness of patient influenza immunization with clear direction from VHA leadership (i.e., Directive or Flu Advisory)
 - Consider customizing information for local distribution (e.g., bulletins, announcements, email messages)
 - Solicit local leadership buy-in and involvement
 - Use photos of hospital director or other dignitaries getting their flu shot in newsletters, VA TV/monitor displays
 - Use performance feedback
 - Set goals/benchmarks
 - Monitor and inform providers about number and % of high-risk patients vaccinated
 - Encourage friendly competition among providers or clinics
 - Provide incentives to providers with high patient vaccination rates

2. Systems
 - Use computerized clinical record reminders
 - Remove administrative barriers
 - Use standing orders for inpatients (acute as well as long-term care, and psychiatric settings), outpatients and Home Care patients
 - Use patient reminders (postcards/letters) and recall systems

3. Convenience
 - Expand access/outreach
 - Extended clinic hours/days
 - Drop-In vaccination days, ‘Drive-by’ vaccination
 - Vaccination in settings previously not used routinely for this purpose (hospital lobbies, Vet Centers, Domiciliaries)
 - Target special populations in clinics where they are likely to be seen (SCI, HIV/ID clinics, Homeless programs)

4. Communication
 - Use multiple message formats
 - Provider email, “on hold” messages for patient callers, newsletters, posters, buttons, stickers, pens, cafeteria tray liners, table tents
 - Phone calls and/or mailed reminders to outpatients
 - Include reminders with pharmacy refills, appointment letters
 - Home visits can include influenza immunization

- Encourage patients to help keep track of their vaccinations
- Education for patients
 - Provide fact sheets to all patients sitting in clinic waiting areas
 - Direct and simple
 - Appropriate languages and reading level
 - Info can also be broadcast through medical media to inpatients or presented on VA TV/monitors throughout the medical center
 - Enlist providers from multiple disciplines and trainees to assist with patient education efforts (PharmD, nursing students, interns and residents)
- Education for employees
 - Provide training on importance and effectiveness of influenza vaccination (Grand Rounds, staff meetings)
 - Provide information and tools to facilitate successful patient influenza immunization campaigns

Getting Health Care Workers and Staff Vaccinated

1. Organizational
 - Make influenza vaccination an organizational priority
 - Provide written policy stressing importance of vaccination for health care workers (HCW) with clear direction from leadership (i.e., Directive or Flu Advisory)
 - Customize information for local distribution with local leadership buy-in and involvement
 - Use photos of hospital director or other dignitaries getting their flu shot (newsletters, posters, TV/Monitor displays)
 - Enlist peer vaccination champions to encourage HCW immunizations
 - Performance feedback
 - Set goals/benchmarks, encourage friendly competition among staff in different clinical settings, provide incentives to staff who receive vaccine through worksite or private source
 - Gift incentives may include buttons, canteen vouchers, movie passes or raffle tickets for specific items
2. Systems
 - Provide standing orders
 - Develop ways to monitor employee vaccination rates and provide feedback to specific clinics or settings
 - As an alternative to flu shots, consider accessing FluMist™ (nasal or LAIV) for staff under age 50 who do not routinely treat severely immunocompromised patients and have no contradiction.
3. Convenience
 - Expand access/outreach
 - Extended Occupational Health hours
 - Drop-In vaccination days, ‘Drive-by’ vaccination
 - Rolling carts—bringing vaccine directly to the employees’ work setting
 - Set up vaccination clinics in high-traffic areas (canteen, main hallways, etc.)
 - Include options for off-tour and offsite employees
4. Communication
 - Use multiple message formats
 - Provider email, newsletters, posters, buttons, pens, cafeteria table tents
 - Pay check stubs, Hey VA email messages

- Post schedules ahead of time for mobile carts and flu clinics
- Education
 - Provide training on importance and effectiveness of influenza vaccination (Grand Rounds, staff meetings)
 - Add to standard curricula of annual employee training session
 - Emphasize high risk to patients when staff not vaccinated
 - Emphasize low risk of side effects from vaccine, debunk myths

Section 5

Tools for Influenza Prevention

The 2005 - 2006 FLU TOOLKITS include a manual, buttons, stickers and new posters. Eight key contact groups at each VA medical facilities were sent the toolkit. These groups included:

- Influenza vaccination (“flu”) coordinators
- Occupational health clinicians
- Prevention coordinators
- Infection control professionals
- Public affairs officers
- Patient education contacts
- Patient safety officers
- Education contacts (Employee Education System)

WHAT IS THE VA INFLUENZA TOOLKIT MANUAL?

This manual is a resource for VA providers to use during flu season. It contains information on:

- **Influenza Illness**
- **Inactivated Influenza Vaccine**
- **Live, Attenuated Intranasal Influenza Vaccine (LAIV, FluMist™)**
- **Pneumococcal Vaccine**
- **How to Improve Health Care Worker Vaccination Rates**
- **Best Strategies for Increasing Adult Influenza Vaccination Rates**
- **Frequently Asked Questions**
- **And much more....**

Additional copies can be downloaded from www.publichealth.va.gov/InfectionDontPassItOn.

HOW CAN I USE THE BUTTONS AND STICKERS DURING FLU SEASON?

Buttons Two different buttons were designed. They are to be worn by VA health care workers. These are designed to encourage conversation between staff and patients on influenza vaccinations.





Stickers A sticker was designed to be distributed to health care workers and patients who have received their flu shots. Every time someone in VA receives a flu shot, she/he should get a sticker to wear.

ARE THERE MORE POSTERS?

Posters New posters were created to prevent infection acquired in the health care setting, especially influenza, but also other infections. The Flu Toolkit includes 27 different 8.5” x 11” posters designed for flu prevention, hand & respiratory hygiene, and bathroom etiquette.



WHO SHOULD SEE THE POSTERS ?

The code in the lower right of each poster includes the target audience with the words “ALL,” “CLINICAL.” The posters marked:

- “ALL” can be used in areas frequented by patients, visitors, families, VA employees, and staff including health care clinicians and providers of all types.
- “CLINICAL” are directed specifically to providers and should be hung where clinicians will see them.

WHERE DO I HANG THE POSTERS ?

Use the posters in places that will get these messages to the VA community. Place them in multiple sites throughout hospitals, clinics, domiciliaries, Vet Centers, etc. Posters can be hung or placed at reception desks, waiting areas, exam rooms, rest rooms, meeting/conference rooms, cafeterias, established kiosks, and bulletin boards. Rotate them often (weekly or monthly). The “Restroom” and “Wash” posters may be used in restrooms, hung near urinals, in stalls, on mirrors or sinks or soap dispensers. Posters may also be hung on restroom doors, especially so they can be seen upon exiting the restroom.

HOW CAN I PRINT POSTERS DIRECTLY FROM THE WEB ?

The poster number is located on the bottom right corner of the poster. Find the poster you want at the Web site below and right click on it to either print it or save (download) it to print later:

<http://www.publichealth.va.gov/InfectionDontPassItOn/>

New Posters “INFECTION: DON’T PASS IT ON” Campaign

Poster Title	Poster Description	Audience	Format
INFLUENZA			
Flu 1	Staying Healthy	All	Portrait
Flu 2	At 65 I’m healthy	All	Portrait
Flu 3	If you’re 65	All	Portrait
Flu 4	Get the Flu Shot	All	Portrait
Flu 5	Ask for a Flu Shot	All	Portrait
Flu 6	Keep Veterans Healthy	All	Portrait
Flu 7	Do Your Part Keep Veterans Healthy	All	Portrait
Flu 8	Are you ready?	Clinical	Portrait
Flu 9	When you care for a patient	Clinical	Portrait
Flu 10	Infection Control Management	Clinical	Portrait
HAND WASHING HYGIENE using soap & water			
Wash 1	Where to Wash	All	Portrait
Wash 2	Help a Veteran	All	Portrait
Wash 3	What are the Top 10?	All	Portrait
Wash 4	Hand Washing Checklist	All	Portrait
Wash 5	Wash Your Hands	All	Portrait
Wash 6	Wash Your Hands	All	Portrait
Wash 7	Handwashing - Good!	All	Portrait
Wash 8	Wash Your Hands	All	Portrait
Wash 9	Patients & Visitors: Wash Your Hands	All	Portrait
Wash 10	Patients & Visitors: Wash Your Hands	All	Portrait
RESTROOM			
Restroom 1	Bathroom Rules	All	Portrait
Restroom 2	After Using the Bathroom	All	Landscape
Restroom 3	Before you leave	All	Portrait
Restroom 4	What’s the Rush?	All	Landscape
OTHERS			
Hands 37	Patients & Visitors: It’s OK to ask	All	Portrait
Prevent 14	Keep Veterans Healthy	All	Portrait
Prevent 15	Do Your Part Keep Veterans Healthy	All	Portrait

WHAT DO THE POSTERS LOOK LIKE ?

DO YOUR PART TO KEEP VETERANS HEALTHY

COVER your coughs and sneezes


CLEAN your hands



Protect yourself and others

infection: Don't Pass It On www.publichealth.va.gov/InfectionDontPassItOn Department of Veterans Affairs

What Are the Top 10 Ways to Spread Germs?



CLEAN YOUR HANDS

Wash with soap & water

infection: Don't Pass It On www.publichealth.va.gov/InfectionDontPassItOn Department of Veterans Affairs

Hand Washing Checklist

- Make lather with soap and water
- Rub palms, back of hands, and between fingers
- Rub for 15 seconds
- Rinse and dry well



infection: Don't Pass It On www.publichealth.va.gov/InfectionDontPassItOn Department of Veterans Affairs

Handwashing - Good!

1. Make lather with soap and water
2. Rub palms, back of hands, and between fingers
3. Rub for 15 seconds
4. Rinse and dry well

Germs - Bad!



infection: Don't Pass It On www.publichealth.va.gov/InfectionDontPassItOn Department of Veterans Affairs

WASH YOUR HANDS

- Make lather with soap and water
- Rub palms, back of hands, and between fingers
- Rub for 15 seconds
- Rinse and dry well



infection: Don't Pass It On www.publichealth.va.gov/InfectionDontPassItOn Department of Veterans Affairs

Patients & Visitors:

WASH YOUR HANDS

BEFORE:
Eating or preparing food
Caring for someone sick

AFTER:
Going to the bathroom
Caring for someone sick
Blowing your nose
Coughing or sneezing
Touching garbage
Handling raw meat
Touching an animal




infection: Don't Pass It On www.publichealth.va.gov/InfectionDontPassItOn Department of Veterans Affairs

WASH	YOUR	HANDS
	1 Make lather with soap and water	
2 Rub palms, back of hands, and between fingers		3 Rub for 15 seconds
	4 Rinse and dry well	

infection: Don't Pass It On www.publichealth.va.gov/InfectionDontPassItOn Department of Veterans Affairs

PATIENTS & VISITORS:
Wash Your Hands

BEFORE:
 Eating or preparing food • Caring for someone sick



AFTER:
 Going to the bathroom • Caring for someone sick
 Blowing your nose • Coughing or sneezing
 Touching garbage • Handling raw meat • Touching an animal

Infection: Don't Pass It On
 www.publichealth.va.gov/infectionDontPassItOn
 Department of Veterans Affairs
 Health 10 .04

BATHROOM RULES:

ALWAYS FLUSH

ALWAYS WASH YOUR HANDS

Infection: Don't Pass It On
 www.publichealth.va.gov/infectionDontPassItOn
 Department of Veterans Affairs
 Health 10 .04

1. Make lather with soap and water



2. Rub palms, back of hands, and between fingers



WASH YOUR HANDS

3. Rub for 15 seconds




4. Rinse and dry well

Infection: Don't Pass It On
 www.publichealth.va.gov/infectionDontPassItOn
 Department of Veterans Affairs
 Health 10 .04

**What's the RUSH?
 Remember to FLUSH!**



AND, WASH YOUR HANDS!

Infection: Don't Pass It On
 www.publichealth.va.gov/infectionDontPassItOn
 Department of Veterans Affairs
 Health 10 .04

Patients & Visitors:



It's okay to ask health care providers if they have cleaned their hands.

Infection: Don't Pass It On
 www.publichealth.va.gov/infectionDontPassItOn
 Department of Veterans Affairs
 Health 10 .04

Help A Veteran

WASH YOUR HANDS

It only takes seconds



Infection: Don't Pass It On
 www.publichealth.va.gov/infectionDontPassItOn
 Department of Veterans Affairs
 Health 10 .04

Before you leave...

1. Flush



2. Wash Hands



stop the spread of germs

Infection: Don't Pass It On
 www.publichealth.va.gov/infectionDontPassItOn
 Department of Veterans Affairs
 Health 10 .04

KEEP VETERANS HEALTHY

Clean Your Hands

Cover Your Coughs & Sneezes

Get Your Flu Shot

Infection: Don't Pass It On www.publichealth.va.gov/InfectionDontPassItOn Department of Veterans Affairs

Keep Veterans Healthy

DEPARTMENT OF VETERANS AFFAIRS
UNITED STATES OF AMERICA

Clean your hands
Cover your coughs and sneezes

Infection: Don't Pass It On www.publichealth.va.gov/InfectionDontPassItOn Department of Veterans Affairs

DO YOUR PART TO KEEP VETERANS HEALTHY

COVER your coughs and sneezes

CLEAN your hands

Ask about your Flu Shot

Infection: Don't Pass It On www.publichealth.va.gov/InfectionDontPassItOn Department of Veterans Affairs

Are you ready?

Influenza season is here

Get your influenza vaccination
An annual flu vaccine can help protect you, and prevent you from giving flu to your patients, your colleagues and your family.

Vaccinate your patients
The flu vaccine is the best way to prevent influenza hospitalizations and deaths. Influenza causes an average of 36,000 deaths and 114,000 hospitalizations per year in the U.S.

Protect your community
Promote the flu vaccine throughout the influenza season. October and November are the best months to vaccinate, but the vaccine can be given as early as September and can be given in December and beyond.

Infection: Don't Pass It On www.publichealth.va.gov/InfectionDontPassItOn Department of Veterans Affairs CDC

ASK FOR A FLU SHOT

Flu is STILL spreading

Flu season can last into springtime

FLU SHOTS ARE AVAILABLE HERE

Protect yourself and your loved ones from flu & other germs by:

1. Keeping your hands clean
2. Covering your sneezes & coughs
3. Getting a flu shot

Infection: Don't Pass It On www.publichealth.va.gov/InfectionDontPassItOn Department of Veterans Affairs

"If you're 65 or older, you should be among the first to get a flu shot every year."

The flu kills 36,000 people each year. Most are 65 and over.

Stay healthy and strong.
Get a flu shot.

Infection: Don't Pass It On www.publichealth.va.gov/InfectionDontPassItOn Department of Veterans Affairs CDC

Get the Flu Shot!

The reasons are all around you.

Infection: Don't Pass It On www.publichealth.va.gov/InfectionDontPassItOn Department of Veterans Affairs

Section 6

Frequently Asked Questions (FAQs) on Influenza Vaccination

Topic Areas

1. General Questions
2. Novel or Pandemic Influenza
3. Eligibility for Influenza Vaccination in VA
4. HIV/AIDS and Influenza Vaccination
5. Influenza Vaccine Storage
6. Live, attenuated, intranasal influenza vaccine (LAIV)

1. General questions

How is influenza illness defined?

Influenza is a febrile respiratory illness caused by influenza virus that can be prevented by vaccination. The table below differentiates influenza from a ‘cold’.

<i>Signs and Symptoms</i>	<i>Flu</i>	<i>Cold</i>
Onset	sudden	gradual
Fever	high (over 100°F); lasting 3 to 4 days	less common; usually low-grade
Cough	dry; can become severe	hacking or congested
Headache	common	rare
Muscle aches and pains	usual; often severe	slight
Tiredness and weakness	can last 2 to 3 weeks	very mild
Extreme exhaustion	early and prominent	rare or never
Chest discomfort	common	mild to moderate
Stuffy nose	sometimes	common
Sneezing	sometimes	usual
Sore throat	sometimes	common

What should everyone know about the flu season?

- The first cases of influenza in the U.S. are usually identified in October.
- Widespread influenza activity appears 6-10 weeks after the first case.
- Influenza kills about 36,000 and hospitalizes more than 200,000 persons in the U.S. each year
- The influenza vaccine changes each year to match the currently circulating type of influenza. The 2005-06 influenza vaccine contains the following types:
 - A/California (H3N2)
 - A/New Caledonia (H1N1)
 - B/Shanghai
- One needs an influenza vaccine *each year* to get the latest protection.
- Influenza vaccination usually begins in September and October in the VA health care system, and continues through March or April.
- Influenza vaccine takes 1-2 weeks to generate protective immunity.

What are the recommendations for vaccination of health care workers against influenza?

All health care workers with patient contact should receive annual influenza vaccination unless they have a contraindication to the vaccine. Vaccination is recommended for health care workers for at least 3 reasons:

- health care workers can give influenza to their patients, co-workers, family members, and others;
- health care workers are at risk of getting influenza from patients with influenza; and
- preventing influenza by vaccinating keeps health care workers healthy and available to come to work to take care of patients.

How will we know whom to vaccinate when? For example, early in the 2004-2005 season, employees not involved in direct care were not supposed to be vaccinated by VA even if the individual employee was over 65 and/or had medical risks from consequences of flu. Will this be the correct approach this year?

For the 2005-2006 season, it is uncertain what the total influenza vaccine supply will be. The CDC and the VA Undersecretary for Health will issue regular advisories and updates to define the timing of priority groups for vaccination. Watch for VA Influenza Vaccine Advisories sent out broadly by email and posted on <http://www.publichealth.va.gov/flu/>

What else (besides vaccine) can one do to protect oneself and others from influenza illness?

- Cover your nose and mouth with a tissue when you cough or sneeze, and dispose of the tissue afterward.
- If you don't have a tissue, cough or sneeze into your sleeve.
- Clean your hands after you cough or sneeze with soap and warm water or an alcohol-based hand cleaner.
 - This VA website includes posters, information, and links about hand and respiratory hygiene: <http://www.publichealth.va.gov/infectiondontpassiton/>
- If you get the flu, avoid exposing others. Stay home from work or school until your fever is gone, and you feel ready to resume normal activities.
- Get pneumococcal vaccine if you're age 65 or older or have a chronic health condition.

What about antiviral medications?

The antiviral medications amantadine (Symmetrel®) and rimantadine (Flumadine®) are on the VA National Formulary and approved for prevention and treatment of influenza A. An alternative for treatment is oseltamivir (Tamiflu®). A supply of oseltamivir is maintained in a VA stockpile for outbreaks of a very serious nature.

Antivirals are most often used to help contain influenza outbreaks in setting such as nursing homes or to protect a high-risk person who is in direct contact with someone who has influenza.

There are some risks in taking antivirals. Some people have serious side effects from them.

To be effective, antivirals should be taken within 24-48 hours of being exposed to influenza or onset of symptoms.

2. Novel pandemic or avian influenza

What is the difference between regular influenza that is around every year and novel pandemic or avian flu?

Influenza virus circulates in humans every year, usually in winter. Several times each century, a strain that is new to humans originates from the re-assortment of a human and animal (sometimes bird or avian) strains. These new or novel strains cause pandemics that can be very serious, because humans have little pre-existing immunity to them and vaccines and antiviral medications take time to develop, supply, and distribute. The 1918-19 pandemic caused as many as 500,000 deaths in the U.S. and 50 million globally. Public health experts around the world and within VA are concerned about and are taking steps to prepare for a pandemic of novel influenza.

3. Eligibility for influenza vaccination in VA

Many wives and children of 100% service connected veterans who have CHAMPVA (Civilian Health and Medical Program of the Department of Veterans Affairs) visit VA medical facilities for flu shots. In some cases, vaccination has been provided to these groups if they meet criteria for flu vaccine. Is this the correct approach?

VA medicines, supplies, etc, cannot legally be given to non-veterans--even if giving influenza vaccine to a non-veteran caregiver of a veteran might prevent the transmission of the flu to a veteran.

Our facility has permitted veterans not currently enrolled for VA health care to go through screening for income or service-connected criteria for enrollment in order to get influenza vaccination. If they meet VA enrollment criteria, then they are given vaccine if they are within one of the priority groups for vaccine. Is this the correct procedure for veterans who might not be cared for by VA?

As long as a supply of vaccine is available, vaccine should be provided to veterans who are enrolled in VA care and who are currently in one of the priority groups as advised by the Under Secretary for Health's Annual Influenza Directive or by a current Influenza Vaccine Advisory. These documents will be posted and kept current at <http://www.publichealth.va.gov/flu/>

Requests for vaccine have been received from our state Soldiers Homes and Nursing Homes that house VA patients. It is difficult to control the supply in these situations, especially in nursing homes where only a few are VA patients. Also, is VA mandated to provide vaccine for employees in Soldier's Homes and Nursing Homes?

State Soldiers Homes' supplies are provided via individual contracts. Unless a VA facility has an agreement to vaccinate residents in a state Soldiers Home or Nursing Home, those veterans would need to visit a VA facility for their vaccination if they are enrolled for VA health care and belong to one of a priority group for vaccination. Only appropriate (i.e. direct patient care) employees within VA facilities can be vaccinated (unless an existing agreement has been negotiated for the State Home or Nursing Home employees). Even if a place houses and cares for veterans, it is not a *de facto* responsibility of VA to go on site and deliver care unless there are existing agreements to provide care.

Our facility does not vaccinate direct patient care volunteers based on the assumption that they can stay at home if necessary. Is this interpretation correct?

The decision with regard to volunteers is an individual VA facility decision. Some VA medical facilities consider volunteers to be necessary for direct patient care functions and include them in the vaccination plans as they would paid employees. Individual VA facilities must evaluate their volunteers' functions and decide on vaccine eligibility.

Should we offer vaccine to medical residents and other trainees who provide direct patient care at the VA during the flu season through our Occupational Health Department?

The decision with regard to residents and other trainees is an individual VA facility decision and should take into account the contractual agreement with the academic affiliate and the availability of the vaccine. Facilities may want to make the same decisions about providing flu vaccine for rotating or temporary trainees (e.g. house staff/medical residents) as they would for volunteers: if they provide needed, direct, hands-on patient care and the VA facility has vaccine available, these individuals should be vaccinated.

Are homeless veterans who attend stand-downs eligible for influenza vaccine?

Flu vaccine given by VA is for veterans who are enrolled for VA health care and who meet existing priority group and tiered vaccination timing plans (if any). If there is any question about a homeless veteran falling within a priority group, VA staff should use their best medical judgment. A very large percentage of homeless veterans are likely to have qualifying medical conditions that meet CDC criteria.

4. HIV/AIDS and influenza vaccination

Should people with HIV/AIDS receive influenza vaccine?

People with chronic underlying medical conditions, including HIV/AIDS, should be vaccinated with inactivated influenza vaccine. People with HIV/AIDS are considered at increased risk from serious influenza-related complications. Persons with advanced HIV disease may have a poor response to immunization. Therefore, chemoprophylaxis (use of antiviral medications for prevention) should also be considered for these patients if they are likely to be exposed to people with influenza.

Are there people with HIV/AIDS who should NOT receive flu shots?

Contraindications to the use of the flu shot in persons with HIV/AIDS are the same as those for uninfected persons: a history of severe allergy (i.e., anaphylactic allergic reaction) to hens' eggs, or a history of onset of Guillain-Barre syndrome during the 6 weeks after vaccination.

Can people with HIV/AIDS receive the live attenuated flu vaccine (LAIV, sold commercially as FluMist™)?

No. Persons with HIV/AIDS are not recommended to receive the live influenza vaccine. LAIV is approved for use only among healthy persons between the ages of 5 and 49 years.

When should people with HIV/AIDS be prescribed antiviral medications for chemoprophylaxis (prevention)?

Persons at high risk of serious influenza-related complications should be given antiviral medications if they are likely to have been exposed to other people with influenza. For example, when a family or household member is diagnosed with influenza, the exposed person with HIV/AIDS should be given chemoprophylaxis for 7 days. Vaccinated and unvaccinated HIV-infected persons who are residents of institutions experiencing an influenza outbreak should be given chemoprophylaxis for the duration of the outbreak or until discharge. People with advanced HIV disease who are not expected to mount an adequate antibody response to influenza vaccination should consider chemoprophylaxis with antiviral medications for the duration of influenza activity in the community, if antiviral medications are available in adequate supply.

There are no published data on interactions between anti-influenza agents such as amantidine and rimantidine and drugs used in the management of HIV infected persons. Patients should be observed for adverse drug reactions to anti-influenza chemoprophylaxis agents, especially when neurologic conditions or renal insufficiency are present.

Should health care workers who have contact with HIV/AIDS patients be vaccinated?

All health care workers with patient contact should receive annual influenza vaccination unless they have a contraindication to the vaccine. Vaccination is recommended for health care workers for at least 3 reasons:

- health care workers can give influenza to their patients, co-workers, family members, and others;
- health care workers are at risk of getting influenza from patients with influenza;
- preventing influenza by vaccinating keeps health care workers healthy and available to come to work to take care of patients

5. Influenza vaccine storage

CDC recommends that influenza vaccine be refrigerated during shipment and immediately upon being received and to prevent from freezing. Would pre-filling syringes and leaving them out of the refrigerator for use during high volume vaccination efforts affect the potency of the vaccine?

A contact at CDC knew of no data on vaccine stability once the vaccine is in the syringe, but the following cautions make sense:

- Don't prefill a large number of syringes because of
 - Increased risk for administration errors
 - Chance of wasting vaccine
 - Risk of inappropriate storage conditions
 - Potential for bacterial overgrowth in vaccines that do not contain a preservative
 - Reduced vaccine potency
- Prefill only a few syringes at a time
 - Prefill the smallest logical number according to your patient flow
 - Try to fill no more than 10 prefilled syringes at a time (1 vial) per person vaccinating
 - Discard any prefilled syringes remaining at the end of the clinic session.
 - Mark the container of prefilled syringes with the date and time of filling
 - Mark each syringe with the date, time and medication.
- In setting up a mass vaccination clinic

- Administer only one type of vaccine per station (keep influenza and pneumococcal vaccines separate)
- Transport the vaccine to the clinic in the manufacturer-supplied packaging at the recommended temperatures
- Keep vaccine vials and prefilled syringes in a cooler (but not in direct contact with ice)

6. Live, attenuated, intranasal influenza vaccine (LAIV or Flumist™)

LAIV is so new. Is it a safe vaccine?

The development of the live attenuated influenza vaccine has been going on since the 1960s. Prior to licensure, the safety of LAIV was studied in 20 clinical trials. More than 6,000 clinical trial participants were in the approved age range of 5-49 years. In healthy children there were no significant differences between vaccine and placebo recipients. Serious adverse reactions have been identified in less than 1% of LAIV recipients, either children or adults, since licensure.

There have been no instances of Guillain-Barré Syndrome reported among LAIV recipients.

Is LAIV really an option for health care workers?

Yes, LAIV is an option for healthy health care workers up through age 49, especially when there is a shortage of inactivated influenza vaccine. Choosing LAIV, currently available as FluMist™, means you are helping to conserve when there is limited inactivated influenza vaccine for high-risk persons who do not have the option of live attenuated influenza vaccine.

Is FluMist™ the only live attenuated influenza vaccine available in the United States?

Yes. FluMist™ is the only live attenuated influenza vaccine licensed for use in the United States. FluMist™ was approved by the FDA in June 2003 and is given intranasally (in the nose).

What does “attenuated” mean?

Attenuated means that the influenza viruses used to make LAIV have been weakened so they cannot grow well in human tissue. Six parts of the genetic material used to make the vaccine have been modified. In order for the vaccine to become as strong as the wild influenza virus, all six of these components would have to undergo genetic changes. Throughout clinical trials, the stability of the virus was studied and it was found that all six of the genetic components of LAIV remained stable maintaining its attenuated, cold-adapted characteristics.

I read that LAIV is “cold adapted.” What does that mean?

Cold adapted means the vaccine survives and replicates only at temperatures less than 25° Celsius. This temperature range allows the vaccine to grow well in the nose and throat. However, once the virus reaches the lower respiratory tract, the warm temperature destroys the virus. This means the vaccine virus, unlike influenza virus, cannot replicate and cause disease.

Is shedding the virus a problem for health care workers?

The FluMist™ package insert states that a person can shed the virus for up to three weeks because that is what the studies in humans showed, but shedding alone should not be equated with person-to-person transmission. In fact, studies have found that it is very rare. In a study conducted in a Finnish daycare center that was designed to maximize the chance of detecting vaccine virus transmission, one child shed the virus for 21 days. Other children in this study shed the virus a mean of 7.6 days. Estimated transmission rates were extremely low (0.6%-2.4%). There was actually only one documented case of LAIV transmission. An additional small study of 40 adults conducted since licensure found that only 50% of the adults were shedding the vaccine influenza virus on day three after vaccination, one adult shed the virus on day seven. That means that half the adults had stopped shedding the virus by day three. These post licensure studies prompted the Advisory Committee on Immunization Practices (ACIP, an independent committee that advises CDC) to reduce the recommended number of days a HCW should avoid contact with patients requiring protective isolation - from three weeks to seven days.

What if the shed virus is transmitted? Isn't that dangerous?

The virus is shed in lower titers than typically occur with shedding of wild-type influenza viruses. So even if transmission were to occur, there are not enough viral particles to make a person ill. The virus also retains its attenuated characteristics, thus cannot replicate in the lower respiratory tract. The bottom line? Shedding and rare subsequent transmission cannot be presumed to cause disease.

Is LAIV as effective as the regular influenza vaccine?

Yes. Studies conducted before the licensure of LAIV show that it is very protective against influenza disease. CDC discusses the efficacy as follows: “In one large study among children aged 15-85 months, the nasal-spray flu vaccine (LAIV) reduced the chance of influenza illness by 92% compared with placebo. In a study among adults, the participants were not specifically tested for influenza. However, the study found 19% fewer severe febrile respiratory tract illnesses, 24% fewer respiratory tract illnesses with fever, 23-27% fewer days of illness, 13-28% fewer lost work days, 15-41% fewer health care provider visits, and 43-47% less use of antibiotics compared with placebo.” Efficacy of inactivated influenza vaccine is about 70-90% in adults <65 years, and 77-91% effective in children 1-15 years of age. Thus, LAIV is at least as effective as inactivated vaccine.

Who can get LAIV?

People 5 through 49 years of age without any chronic health conditions are eligible for LAIV. This includes household contacts and out-of-home caregivers of infants less than 6 months of age, health care workers, and contacts of persons who have chronic health conditions.

Who cannot get LAIV?

Because no studies were completed at licensure to determine the efficacy or safety in certain persons, LAIV is not recommended for the following:

- People with medical conditions that place them at high risk for complications from influenza, including those with chronic heart or lung disease, such as asthma or reactive airways disease;

people with medical conditions such as diabetes or kidney failure; or people with illnesses that weaken the immune system or who take medications that can weaken the immune system.

- Children or adolescents receiving aspirin therapy.
- People with a history of Guillain-Barré Syndrome, a rare disorder of the nervous system.
- Pregnant women.
- People with a history of allergy to any of the components of LAIV or to eggs.

Can contacts of the persons listed above be vaccinated with LAIV?

Yes. There is only one instance in which healthy people should consider the inactivated flu injection over LAIV – when they are caring for persons requiring protective isolation, e.g., post bone marrow transplant.

Can contacts of persons with weakened immune systems be vaccinated with LAIV?

Yes, LAIV can be used in contacts of persons who have HIV/AIDS, who are on chemotherapy, or who have diseases that weaken their immune system. Remember that the vaccine uses a weakened virus that doesn't contain a viral dose big enough to cause disease. Also, once the virus gets to the lower respiratory tract the warmer temperatures kill it

Should health care workers who have a contraindication to LAIV administer it?

Environmental contamination with LAIV during administration is probably unavoidable. However, because it is an attenuated virus (weakened), that is designed not to replicate at the warm temperatures of the lower respiratory tract, the ACIP does not believe that administration of LAIV by a person with one of the contraindications to it (like asthma, chronic obstructive pulmonary disease, etc) is at risk from infection or illness from the vaccine virus.

What side effects are associated with LAIV?

In children, the most common side effects can include runny nose, headache, vomiting, muscle aches, and fever. In clinical trials upper respiratory symptoms, such as runny nose and nasal congestion, fever, or other systemic symptoms, were reported in 10%-40% of both vaccine and placebo recipients. Adults may experience a runny nose, headache, sore throat, and cough. Unlike children, fever is not a common side effect in adults receiving LAIV.

Can LAIV be given to patients when they are ill?

Persons with mild illnesses, e.g., diarrhea or mild upper respiratory tract infection, with or without fever, may receive the nasal spray flu vaccine (LAIV). Vaccination may need to be delayed for persons who have an illness that has caused nasal congestion because of the interference of vaccine delivery.

Some general questions and questions on LAIV in these FAQs were adapted from the Minnesota Department of Health's website on influenza, available at

<http://www.health.state.mn.us/divs/idepc/diseases/flu/index.html> Used with permission.

Section 7 Appendices

- A. How to Administer Flu and Pneumococcal Vaccines**
- B. CDC Prevention and Control of Influenza. Recommendations of the Advisory Committee on Immunization Practices (ACIP). MMWR. July 29, 2005 / 54**
- C. CDC Adult Immunization Schedule, 2004-2005 [2005-2006 to be released in fall 2005]**
- D. Prevention and Treatment of Influenza with Antiviral Drugs**
- E. Resources, References, and Web sites**
- F. Acknowledgements**

Appendix A

How to Administer Flu and Pneumococcal Vaccines

Note that the inactivated influenza vaccine MUST be administered intramuscularly with a 1”-2” 22-25 gauge needle. Shorter needles should not be used. The pneumococcal polysaccharide vaccine can be administered either IM (1”-2” 22-25 gauge needle) or SC (5/8”-3/4” 23-25 gauge needle).

The following is a brief guide to inactivated influenza and pneumococcal vaccine administration.

1. **Provide the patient with a Vaccine Information Statement (VIS).** The patient should be given a copy of the vaccine’s VIS prior to vaccine administration. This must be a copy that the patient can read and take home. Copies of influenza and pneumococcal VISs are included in this toolkit.
2. **Give the vaccine properly.**
 - **Examine and prepare the vaccine.** Always double check the vial label to make sure that you have the vaccine you want to administer. Shake the vial vigorously and visually inspect it for particulate matter. If you cannot shake the vaccine into a relatively even suspension, do not use it. After wiping the rubber stopper with an alcohol swab, load the syringe by injecting air into the vial, the same volume of air as the dose of vaccine to be drawn.
 - **Site and route of administration:** Inactivated influenza vaccines are administered intramuscularly (IM). The pneumococcal polysaccharide vaccine (PPV) can be administered either IM or subcutaneously (SC). IM injections should be injected directly into the deltoid muscle, below the shoulder on the upper arm. SC injections are given into the fatty tissue on the back of the upper arm.
 - **Proper needle gauge and length:** The proper needle length for IM injections is a 1”-2” 22-25 gauge needle. SC injections use a 5/8”-3/4” 23-25 gauge needle.
3. **Safely dispose of the needle and syringe.** Do not recap the needle after use. Discard the uncapped used needle into a sharps container. If a non safety product is being used, keep your eyes on the needle continuously until it is inside the container. If you are using a safety product, activate the safety mechanism before discarding in the sharps container.
4. **Prepare and watch for an allergic reaction (anaphylaxis).** Acute anaphylactic reactions are very rare, occurring after approximately 1 out of every 500,000 doses of vaccine. When they occur, however, you must take immediate action. No vaccine should ever be administered unless epinephrine, diphenhydramine, adult airways, and blood pressure cuffs are on hand. Staff should be familiar with an anaphylaxis protocol and with cardiopulmonary resuscitation (CPR). After you have administered a vaccine to a patient, instruct the patient to report any itching, redness (with or without hives), difficulty breathing, or abdominal pain within several minutes of injection. Having the patient wait 20 minutes in a post-injection area is suggested but is not officially recommended.

Content adapted from “Adults Only Vaccination: A Step-By-Step Guide,” Immunization Action Coalition (IAC), 2004.

For a detailed explanation and demonstration of immunization techniques, we recommend the 35-minute video “Immunization Techniques: Safe, Effective, Caring,” that can be ordered through the Immunization Action Coalition (IAC) at <http://www.immunize.org> and clicking the link for Video: IZ Techniques.

Instruction sheets on vaccine administration are also available from the IAC:

1. “How to administer IM and SC injections to adults,” available at:
<http://www.immunize.org/catg.d/p2020A.pdf>
2. “Administering vaccines to adults: Dose, route, site, needle size, and preparation,” available at: <http://www.immunize.org/catg.d/p3084.pdf>
3. Instructions for administration of FluMist™
<http://www.flumist.com/professional/admin/admin.asp>

Appendix B
CDC Prevention and Control of Influenza.
Recommendations of the Advisory Committee on Immunization
Practices (ACIP)

For published updates during the flu season visit: <http://www.cdc.gov/mmwr/>



MMWRTM

Morbidity and Mortality Weekly Report

Recommendations and Reports

July 29, 2005 / Vol. 54 / No. RR-8

Prevention and Control of Influenza

Recommendations of the Advisory Committee on Immunization Practices (ACIP)

The *MMWR* series of publications is published by the Coordinating Center for Health Information and Service, Centers for Disease Control and Prevention (CDC), U.S. Department of Health and Human Services, Atlanta, GA 30333.

SUGGESTED CITATION

Centers for Disease Control and Prevention. Prevention and control of influenza: Recommendations of the Advisory Committee on Immunization Practices (ACIP). *MMWR* 2005;54(No. RR-8):[inclusive page numbers].

Centers for Disease Control and Prevention

Julie L. Gerberding, MD, MPH
Director

Dixie E. Snider, MD, MPH
Chief Science Officer

Tanja Popovic, MD, PhD
(Acting) Associate Director for Science

Coordinating Center for Health Information and Service

Blake Caldwell, MD, MPH, and Edward J. Sondik, PhD
(Acting) Directors

National Center for Health Marketing*

Steven L. Solomon, MD
(Acting) Director

Division of Scientific Communications*

Maria S. Parker
(Acting) Director

Mary Lou Lindegren, MD
(Acting) Editor, MMWR Series

Suzanne M. Hewitt, MPA
Managing Editor, MMWR Series

Teresa F. Rutledge
*(Acting) Lead Technical Writer-Editor
Project Editor*

Beverly J. Holland
Lead Visual Information Specialist

Lynda G. Cupell
Malbea A. LaPete
Visual Information Specialists

Quang M. Doan, MBA
Erica R. Shaver
Information Technology Specialists

* Proposed.

CONTENTS

Introduction	1
Primary Changes and Updates in the Recommendations ..	2
Influenza and Its Burden	2
Options for Controlling Influenza	3
Recommendations for Using Inactivated and Live,	
Attenuated Influenza Vaccines	9
Target Groups for Vaccination	9
Additional Information Regarding Vaccination	
of Specific Populations	11
Comparison of LAIV with Inactivated Influenza Vaccine ..	12
Inactivated Influenza Vaccine Recommendations	13
Live, Attenuated Influenza Vaccine Recommendations	16
Recommended Vaccines for Different Age Groups	19
Timing of Annual Influenza Vaccination	19
Strategies for Implementing Vaccination	
Recommendations in Health-Care Settings	21
Influenza Vaccine Supply	22
Future Directions for Influenza Vaccine	
Recommendations	22
Recommendations for Using Antiviral Agents	
for Influenza	22
Role of Laboratory Diagnosis	23
Indications for Use	23
Dosage	25
Route	27
Pharmacokinetics	28
Side Effects and Adverse Reactions	28
Use During Pregnancy	29
Drug Interactions	29
Antiviral Drug-Resistant Strains of Influenza	30
Sources of Information Regarding Influenza	
and Its Surveillance	30
Additional Information Regarding Influenza Infection	
Control Among Specific Populations	30
References	31

Prevention and Control of Influenza

Recommendations of the Advisory Committee on Immunization Practices (ACIP)

Prepared by

Scott A. Harper, MD¹

Keiji Fukuda, MD¹

Timothy M. Uyeki, MD¹

Nancy J. Cox, PhD¹

Carolyn B. Bridges, MD²

¹*Division of Viral and Rickettsial Diseases
National Center for Infectious Diseases*

²*Epidemiology and Surveillance Division
National Immunization Program*

Summary

This report updates the 2004 recommendations by the Advisory Committee on Immunization Practices (ACIP) regarding the use of influenza vaccine and antiviral agents (CDC. Prevention and control of influenza: recommendations of the Advisory Committee on Immunization Practices [ACIP]. MMWR 2004;53[No. RR-6]:1–40). The 2005 recommendations include new or updated information regarding 1) vaccination of persons with conditions leading to compromise of the respiratory system; 2) vaccination of health-care workers; 3) clarification of the role of live, attenuated influenza vaccine (LAIV) in vaccine shortage situations; 4) the 2005–06 trivalent vaccine virus strains: A/California/7/2004 (H3N2)-like, A/New Caledonia/20/99 (H1N1)-like, and B/Shanghai/361/2002-like antigens (for the A/California/7/2004 [H3N2]-like antigen, manufacturers may use the antigenically equivalent A/New York/55/2004 virus, and for the B/Shanghai/361/2002-like antigen, manufacturers may use the antigenically equivalent B/Jilin/20/2003 virus or B/Jiangsu/10/2003 virus); and 5) the assessment of vaccine supply, timing of influenza vaccination, and prioritization of inactivated vaccine in shortage situations. A link to this report and other information can be accessed at <http://www.cdc.gov/flu>.

Introduction

Epidemics of influenza typically occur during the winter months in temperate regions and have been responsible for an average of approximately 36,000 deaths/year in the United States during 1990–1999 (1). Influenza viruses also can cause pandemics, during which rates of illness and death from influenza-related complications can increase worldwide. Influenza viruses cause disease among all age groups (2–4). Rates of infection are highest among children, but rates of serious illness and death are highest among persons aged ≥ 65 years, children aged < 2 years, and persons of any age who have medical conditions that place them at increased risk for complications from influenza (2,5–7).

The material in this report originated in the National Center for Infectious Diseases, Anne Schuchat, MD, Acting Director, and the Division of Viral and Rickettsial Diseases, James LeDuc, PhD, Director; and the National Immunization Program, Stephen Cochi, MD, Acting Director, and Epidemiology and Surveillance Division, Gina Mootrey, DO, MPH, Acting Director.

Corresponding preparer: Nancy Cox, PhD, National Center for Infectious Diseases, 1600 Clifton Road, NE, MS G-16, Atlanta, GA 30333. Telephone: 404-639-3591; Fax: 404-639-2334; E-mail: NJC1@cdc.gov.

Influenza vaccination is the primary method for preventing influenza and its severe complications. In this report from the Advisory Committee on Immunization Practices (ACIP), the primary target groups recommended for annual vaccination are 1) persons at increased risk for influenza-related complications (i.e., those aged ≥ 65 years, children aged 6–23 months, pregnant women, and persons of any age with certain chronic medical conditions); 2) persons aged 50–64 years because this group has an elevated prevalence of certain chronic medical conditions; and 3) persons who live with or care for persons at high risk (e.g., health-care workers and household contacts who have frequent contact with persons at high risk and who can transmit influenza to those persons at high risk). Vaccination is associated with reductions in influenza-related respiratory illness and physician visits among all age groups, hospitalization and death among persons at high risk, otitis media among children, and work absenteeism among adults (8–18). Although influenza vaccination levels increased substantially during the 1990s, further improvements in vaccine coverage levels are needed, chiefly among persons aged < 65 years who are at increased risk for influenza-related complications among all racial and ethnic groups, among blacks and Hispanics aged ≥ 65 years, among children aged 6–23 months,

and among health-care workers. ACIP recommends using strategies to improve vaccination levels, including using reminder/recall systems and standing orders programs (19–21). Although influenza vaccination remains the cornerstone for the control and treatment of influenza, information on antiviral medications is also presented because these agents are an adjunct to vaccine.

Primary Changes and Updates in the Recommendations

The 2005 recommendations include five principal changes or updates:

- ACIP recommends that persons with any condition (e.g., cognitive dysfunction, spinal cord injuries, seizure disorders, or other neuromuscular disorders) that can compromise respiratory function or the handling of respiratory secretions or that can increase the risk for aspiration be vaccinated against influenza (see Target Groups for Vaccination).
- ACIP emphasizes that all health-care workers should be vaccinated against influenza annually, and that facilities that employ health-care workers be strongly encouraged to provide vaccine to workers by using approaches that maximize immunization rates.
- Use of both available vaccines (inactivated and LAIV) is encouraged for eligible persons every influenza season, especially persons in recommended target groups. During periods when inactivated vaccine is in short supply, use of LAIV is especially encouraged when feasible for eligible persons (including health-care workers) because use of LAIV by these persons might considerably increase availability of inactivated vaccine for persons in groups at high risk.
- The 2004–05 trivalent vaccine virus strains are A/California/7/2004 (H3N2)-like, A/New Caledonia/20/99 (H1N1)-like, and B/Shanghai/361/2002-like antigens. For the A/California/7/2004 (H3N2)-like antigen, manufacturers may use the antigenically equivalent A/New York/55/2004 virus, and for the B/Shanghai/361/2002-like antigen, manufacturers may use the antigenically equivalent B/Jilin/20/2003 virus or B/Jiangsu/10/2003 virus (see Influenza Vaccine Composition).
- CDC and other agencies will assess the vaccine supply throughout the manufacturing period and will make recommendations preceding the 2005–06 influenza season regarding the need for tiered timing of vaccination of different risk groups. In addition, CDC will publish ACIP recommendations regarding inactivated vaccine subprioritization (tiering) on a later date in *MMWR*.

Influenza and Its Burden

Biology of Influenza

Influenza A and B are the two types of influenza viruses that cause epidemic human disease (22). Influenza A viruses are further categorized into subtypes on the basis of two surface antigens: hemagglutinin and neuraminidase. Influenza B viruses are not categorized into subtypes. Since 1977, influenza A (H1N1) viruses, influenza A (H3N2) viruses, and influenza B viruses have been in global circulation. In 2001, influenza A (H1N2) viruses that probably emerged after genetic reassortment between human A (H3N2) and A (H1N1) viruses began circulating widely. Both influenza A and B viruses are further separated into groups on the basis of antigenic characteristics. New influenza virus variants result from frequent antigenic change (i.e., antigenic drift) resulting from point mutations that occur during viral replication. Influenza B viruses undergo antigenic drift less rapidly than influenza A viruses.

Immunity to the surface antigens, particularly the hemagglutinin, reduces the likelihood of infection and severity of disease if infection occurs (23). Antibody against one influenza virus type or subtype confers limited or no protection against another type or subtype of influenza. Furthermore, antibody to one antigenic variant of influenza virus might not completely protect against a new antigenic variant of the same type or subtype (24). Frequent development of antigenic variants through antigenic drift is the virologic basis for seasonal epidemics and the reason for the usual incorporation of one or more new strains in each year's influenza vaccine.

Clinical Signs and Symptoms of Influenza

Influenza viruses are spread from person to person primarily through the coughing and sneezing of infected persons (22). The typical incubation period for influenza is 1–4 days, with an average of 2 days (25). Adults can be infectious from the day before symptoms begin through approximately 5 days after illness onset. Children can be infectious for ≥ 10 days, and young children can shed virus for several days before their illness onset. Severely immunocompromised persons can shed virus for weeks or months (26–29).

Uncomplicated influenza illness is characterized by the abrupt onset of constitutional and respiratory signs and symptoms (e.g., fever, myalgia, headache, malaise, nonproductive cough, sore throat, and rhinitis) (30). Among children, otitis media, nausea, and vomiting are also commonly reported with influenza illness (31–33). Respiratory illness caused by influenza is difficult to distinguish from illness caused by other respiratory pathogens on the basis of symptoms alone (see Role of Laboratory Diagnosis). Reported sensitivities and speci-

ficiencies of clinical definitions for influenza-like illness (ILI) in studies primarily among adults that include fever and cough have ranged from 63% to 78% and 55% to 71%, respectively, compared with viral culture (34,35). Sensitivity and predictive value of clinical definitions can vary, depending on the degree of co-circulation of other respiratory pathogens and the level of influenza activity (36). A study among older nonhospitalized patients determined that symptoms of fever, cough, and acute onset had a positive predictive value of 30% for influenza (37), whereas a study of hospitalized older patients with chronic cardiopulmonary disease determined that a combination of fever, cough, and illness of <7 days was 78% sensitive and 73% specific for influenza (38). However, a study among vaccinated older persons with chronic lung disease reported that cough was not predictive of influenza infection, although having a fever or feverishness was 68% sensitive and 54% specific for influenza infection (39).

Influenza illness typically resolves after 3–7 days for the majority of persons, although cough and malaise can persist for >2 weeks. Among certain persons, influenza can exacerbate underlying medical conditions (e.g., pulmonary or cardiac disease), lead to secondary bacterial pneumonia or primary influenza viral pneumonia, or occur as part of a coinfection with other viral or bacterial pathogens (40). Young children with influenza infection can have initial symptoms mimicking bacterial sepsis with high fevers (41,42), and ≤20% of children hospitalized with influenza can have febrile seizures (32,43). Influenza infection has also been associated with encephalopathy, transverse myelitis, Reye syndrome, myositis, myocarditis, and pericarditis (32,40,44,45).

Hospitalizations and Deaths from Influenza

The risks for complications, hospitalizations, and deaths from influenza are higher among persons aged ≥65 years, young children, and persons of any age with certain underlying health conditions (see Persons at Increased Risk for Complications) than among healthy older children and younger adults (1,6,8,46–52). Estimated rates of influenza-associated hospitalizations have varied substantially by age group in studies conducted during different influenza epidemics (Table 1).

Among children aged 0–4 years, hospitalization rates have ranged from approximately 500/100,000 children for those with high-risk medical conditions to 100/100,000 children for those without high-risk medical conditions (53–56). Within the 0–4 year age group, hospitalization rates are highest among children aged 0–1 years and are comparable to rates reported among persons aged ≥65 years (55,56) (Table 1).

During influenza epidemics from 1979–80 through 2000–01, the estimated overall number of influenza-associated hospitalizations in the United States ranged from

approximately 54,000 to 430,000/epidemic. An average of approximately 226,000 influenza-related excess hospitalizations occurred per year, with 63% of all hospitalizations occurring among persons aged ≥65 years (57). Since the 1968 influenza A (H3N2) virus pandemic, the greatest numbers of influenza-associated hospitalizations have occurred during epidemics caused by type A (H3N2) viruses (58).

Influenza-related deaths can result from pneumonia and from exacerbations of cardiopulmonary conditions and other chronic diseases. Deaths of older adults account for ≥90% of deaths attributed to pneumonia and influenza (1,52). In one study of influenza epidemics, approximately 19,000 influenza-associated pulmonary and circulatory deaths per influenza season occurred during 1976–1990, compared with approximately 36,000 deaths during 1990–1999 (1). Estimated rates of influenza-associated pulmonary and circulatory deaths/100,000 persons were 0.4–0.6 among persons aged 0–49 years, 7.5 among persons aged 50–64 years, and 98.3 among persons aged ≥65 years. In the United States, the number of influenza-associated deaths might be increasing in part because the number of older persons is increasing (59). In addition, influenza seasons in which influenza A (H3N2) viruses predominate are associated with higher mortality (60); influenza A (H3N2) viruses predominated in 90% of influenza seasons during 1990–1999, compared with 57% of seasons during 1976–1990 (1).

Deaths from influenza are uncommon among both children with and without high-risk conditions, but do occur (61,62). A study that modeled influenza-related deaths estimated that an average of 92 deaths (0.4 deaths per 100,000) occurred among children aged <5 years annually during the 1990's, compared with 32,651 deaths (98.3 per 100,000) among adults aged ≥65 years (1). Reports of 153 laboratory-confirmed influenza-related pediatric deaths from 40 states during the 2003–04 influenza season indicated that 61 (40%) were aged <2 years and, of 92 children aged 2–17 years, 64 (70%) did not have an underlying medical condition traditionally considered to place a person at risk for influenza-related complications (CDC, National Center for Infectious Diseases, unpublished data, 2005). Further information is needed regarding the risk for severe influenza-complications and optimal strategies for minimizing severe disease and death among children.

Options for Controlling Influenza

In the United States, the primary option for reducing the effect of influenza is immunoprophylaxis with vaccine. Inactivated (i.e., killed virus) influenza vaccine and live, attenuated influenza vaccine are available for use in the United States (see Recommendations for Using Inactivated and Live, At-

TABLE 1. Estimated rates of influenza-associated hospitalization, by age group and risk group for selected studies* — United States

Study years	Population	Age group	Hospitalizations/ 100,000 persons with high-risk conditions	Hospitalizations/ 100,000 persons without high-risk conditions
1973–1993† § ¶	Tennessee Medicaid	0–11 mos	1,900	496–1,038**
		1–2 yrs	800	186
		3–4 yrs	320	86
		5–14 yrs	92	41
1992–1997†† §§	Two health maintenance organizations	0–23 mos		144–187
		2–4 yrs		0–25
		5–17 yrs		8–12
1968–1969, 1970–1971, 1972–1973¶¶ ***	Health maintenance organization	15–44 yrs	56–110	23–25
		45–64 yrs	392–635	13–23
		≥65 yrs	399–518	—
1969–1995*** †††	National Hospital Discharge Data	<65 yrs	—	20–42§§§ ¶¶¶
1969–1995*** †††		≥65 yrs	—	125–228¶¶¶
1979–2001**** ††††	National Hospital Discharge Data	All ages	—	88.4§§§§

* Rates were estimated in years and populations with low vaccination rates. Hospitalization rates can be expected to decrease as vaccination rates increase. Vaccination can be expected to reduce influenza-related hospitalizations by 30%–70% among older persons and likely by even higher percentages among younger age groups when vaccine and circulating influenza virus strains are antigenically similar.

† **Source:** Neuzil KM, Mellen BG, Wright PF, Mitchel EF, Griffin MR. Effect of influenza on hospitalizations, outpatient visits, and courses of antibiotics in children. *N Engl J Med* 2000;342:225–31.

§ Outcomes were for acute cardiac or pulmonary conditions.

¶ **Source:** Neuzil KM, Wright PF, Mitchel EF, Griffin MR. Burden of influenza illness in children with asthma and other chronic medical conditions. *J Pediatr* 2000;137:856–64.

** The low estimate is for infants aged 6–11 months, and the high estimate is for infants aged 0–5 months.

†† **Source:** Izurieta HA, Thompson WW, Kramarz P, et al. Influenza and the rates of hospitalization for respiratory disease among infants and young children. *N Engl J Med* 2000;342:232–9.

§§ Outcomes were for acute pulmonary conditions. Influenza-attributable hospitalization rates for children at high risk were not included in this study.

¶¶ **Source:** Barker WH, Mullooly JP. Impact of epidemic type A influenza in a defined adult population. *Am J Epidemiol* 1980;112:798–811.

*** Outcomes were limited to hospitalizations in which either pneumonia or influenza was listed as the first condition on discharge records (Simonsen) or included anywhere in the list of discharge diagnoses (Barker).

††† **Source:** Simonsen L, Fukuda K, Schonberger LB, Cox NJ. Impact of influenza epidemics on hospitalizations. *J Infect Dis* 2000;181:831–7.

§§§ Persons at high risk and not at high risk for influenza-related complications are combined.

¶¶¶ The low estimate is the average during influenza A (H1N1) or influenza B-predominate seasons, and the high estimate is the average during influenza A (H3N2)-predominate seasons.

**** Outcomes were for rate of primary respiratory and circulatory hospitalizations.

†††† **Source:** Thompson WW, Shay DK, Weintraub E, et al. Influenza-associated hospitalizations in the United States. *JAMA* 2004;292:1333–40.

§§§§ Rate for all ages of persons, both with and without high-risk conditions.

tenuated Influenza Vaccine). Vaccinating persons at high risk for complications and their contacts each year before seasonal increases in influenza virus circulation is the most effective means of reducing the effect of influenza. Vaccination coverage can be increased by administering vaccine to persons during hospitalizations or routine health-care visits before the influenza season, making special visits to physicians' offices or clinics unnecessary. When vaccine and epidemic strains are well-matched, achieving increased vaccination rates among persons living in closed settings (e.g., nursing homes and other chronic-care facilities) and among staff can reduce the risk for outbreaks by inducing herd immunity (13). Vaccination of health-care workers and other persons in close contact with persons at increased risk for severe influenza illness can also

reduce transmission of influenza and subsequent influenza-related complications. Antiviral drugs used for chemoprophylaxis or treatment of influenza are a key adjunct to vaccine (see Recommendations for Using Antiviral Agents for Influenza). However, antiviral medications are not a substitute for vaccination.

Influenza Vaccine Composition

Both the inactivated and live, attenuated vaccines prepared for the 2005–06 season will include A/California/7/2004 (H3N2)-like, A/New Caledonia/20/99 (H1N1)-like, and B/Shanghai/361/2002-like antigens. For the A/California/7/2004 (H3N2)-like antigen, manufacturers may use the antigenically equivalent A/New York/55/2004 (H3N2) virus, and

for the B/Shanghai/361/2002-like antigen, manufacturers may use the antigenically equivalent B/Jilin/20/2003 virus or B/Jiangsu/10/2003 virus. These viruses will be used because of their growth properties and because they are representative of influenza viruses likely to circulate in the United States during the 2005–06 influenza season. Because circulating influenza A (H1N2) viruses are a reassortant of influenza A (H1N1) and (H3N2) viruses, antibody directed against influenza A (H1N1) and influenza (H3N2) vaccine strains provides protection against circulating influenza A (H1N2) viruses. Influenza viruses for both the inactivated and live attenuated influenza vaccines are initially grown in embryonated hens eggs. Thus, both vaccines might contain limited amounts of residual egg protein.

For the inactivated vaccine, the vaccine viruses are made noninfectious (i.e., inactivated or killed) (63). Subvirion and purified surface antigen preparations of the inactivated vaccine are available. Manufacturing processes differ by manufacturer. Manufacturers might use different compounds to inactivate influenza viruses and add antibiotics to prevent bacterial contamination. Package inserts should be consulted for additional information.

Thimerosal

Thimerosal, a mercury-containing compound, has been used as a preservative in vaccines (64) since the 1930s and is used in multi-dose vials of inactivated influenza vaccine to reduce the likelihood of bacterial contamination. Although no scientific evidence indicates that thimerosal in vaccines leads to serious adverse events in vaccine recipients, in 1999, the U.S. Public Health Service and other organizations recommended that efforts be made to eliminate or reduce the thimerosal content in vaccines to decrease total mercury exposure, chiefly among infants (64–66). Since mid-2001, vaccines routinely recommended for infants in the United States have been manufactured either without or with only trace amounts of thimerosal to provide a substantial reduction in the total mercury exposure from vaccines for children (67). Vaccines containing trace amounts of thimerosal have <1 mcg mercury/dose.

Influenza Vaccines and Thimerosal. LAIV does not contain thimerosal. Thimerosal preservative-containing inactivated influenza vaccines, distributed in multi-dose containers in the United States, contain 25 mcg of mercury/0.5-mL dose (64,65). Inactivated influenza virus vaccines distributed in the United States will also be available in 2005 in a *thimerosal-free* formulation in both 0.25 mL and 0.5-mL single-dose syringes and a preservative-free formulation (which contains trace amounts of thimerosal) in 0.25-mL-dose syringes.

Influenza vaccine is part of the routine childhood immunization schedule. Sanofi Pasteur, Inc. (formerly Aventis Pas-

teur, Inc.) produces FluZone[®], which is an inactivated influenza vaccine approved by the Food and Drug Administration (FDA) for persons aged ≥ 6 months. FluZone that is available in multi-dose vials contains thimerosal as a preservative. Thimerosal-free FluZone packaged as 0.25-mL unit dose syringes is available for use among persons aged 6–35 months. Thimerosal-free FluZone packaged as 0.5 mL unit dose syringes is available for use among persons aged ≥ 3 years. Fluvirin[®], produced by Chiron, is an inactivated influenza vaccine available in a preservative-free formulation, is packaged as 0.5-mL single-dose syringes, and is licensed for use in persons aged ≥ 4 years. The preservative-free Fluvirin vaccine contains trace amounts of thimerosal. The total amount of inactivated influenza vaccine available without thimerosal as a preservative will be increased as manufacturing capabilities are expanded.

The risks for severe illness from influenza infection are elevated among both young children and pregnant women, and both groups benefit from vaccination by preventing illness and death from influenza. In contrast, no scientifically conclusive evidence exists of harm from exposure to thimerosal preservative-containing vaccine, whereas evidence is accumulating of lack of any harm resulting from exposure to such vaccines (64,68). Therefore, the benefits of influenza vaccination outweigh the theoretical risk, if any, for thimerosal exposure through vaccination. Nonetheless, certain persons remain concerned regarding exposure to thimerosal. The U.S. vaccine supply for infants and pregnant women is in a period of transition during which the availability of thimerosal-reduced or thimerosal-free vaccine intended for these groups is being expanded by manufacturers as a feasible means of reducing an infant's total exposure to mercury because other environmental sources of exposure are more difficult or impossible to eliminate. Reductions in thimerosal in other vaccines have been achieved already and have resulted in substantially lowered cumulative exposure to thimerosal from vaccination among infants and children. For all of these reasons, persons recommended to receive inactivated influenza vaccine may receive either vaccine preparation, depending on availability.

Efficacy and Effectiveness of Inactivated Influenza Vaccine

The effectiveness of inactivated influenza vaccine depends primarily on the age and immunocompetence of the vaccine recipient and the degree of similarity between the viruses in the vaccine and those in circulation. The majority of vaccinated children and young adults develop high postvaccination hemagglutination inhibition antibody titers (69–71). These antibody titers are protective against illness caused by

strains that are antigenically similar to those strains of the same type or subtype included in the vaccine (70–73).

Adults Aged <65 Years. When the vaccine and circulating viruses are antigenically similar, influenza vaccine prevents influenza illness among approximately 70%–90% of healthy adults aged <65 years (9,12,74,75). Vaccination of healthy adults also has resulted in decreased work absenteeism and decreased use of health-care resources, including use of antibiotics, when the vaccine and circulating viruses are well-matched (9–12,75,76). In a case-control study of adults aged 50–64 years with laboratory-confirmed influenza during the 2003–04 season when the vaccine and circulating viruses were not well matched, vaccine effectiveness was estimated to be 52% among healthy persons and 38% among those with one or more high-risk conditions (77).

Children. Children aged 6 months can develop protective levels of anti-influenza antibody against specific influenza virus strains after influenza vaccination (69,70,78–81), although the antibody response among children at high risk for influenza-related complications might be lower than among healthy children (82,83). In a randomized study among children aged 1–15 years, inactivated influenza vaccine was 77%–91% effective against influenza respiratory illness and was 44%–49%, 74%–76%, and 70%–81% effective against influenza seroconversion among children aged 1–5, 6–10, and 11–15 years, respectively (71). One study (84) reported a vaccine efficacy of 56% against influenza illness among healthy children aged 3–9 years, and another study (85) determined vaccine efficacy of 22%–54% and 60%–78% among children with asthma aged 2–6 years and 7–14 years, respectively. A 2-year randomized study of children aged 6–24 months determined that ≥89% of children seroconverted to all three vaccine strains during both years (86). During year 1, among 411 children, vaccine efficacy was 66% (95% confidence interval [CI] = 34%–82%) against culture-confirmed influenza (attack rates: 5.5% and 15.9% among vaccine and placebo groups, respectively). During year 2, among 375 children, vaccine efficacy was –7% (95% CI = –247%–67%; attack rates: 3.6% and 3.3% among vaccine and placebo groups, respectively; the second year exhibited lower attack rates overall and was considered a mild season). However, no overall reduction in otitis media was reported. Other studies report that trivalent inactivated influenza vaccine decreases the incidence of influenza-associated otitis media among young children by approximately 30% (16,17). A retrospective study among approximately 5,000 children aged 6–23 months conducted during a year with a suboptimal vaccine match indicated vaccine effectiveness of 49% against medically attended, clinically diagnosed pneumonia or influenza (*International Classification of Diseases, Ninth Revision* [ICD-9] codes 480–

487) among children who had received 2 doses of influenza vaccine. No effectiveness was demonstrated among children who had received only 1 dose of influenza vaccine, illustrating the importance of administering 2 doses of vaccine to previously unvaccinated children aged <9 years (87).

Adults Aged ≥65 Years. Older persons and persons with certain chronic diseases might develop lower postvaccination antibody titers than healthy young adults and thus can remain susceptible to influenza infection and influenza-related upper respiratory tract illness (88–90). A randomized trial among noninstitutionalized persons aged ≥60 years reported a vaccine efficacy of 58% against influenza respiratory illness, but indicated that efficacy might be lower among those aged ≥70 years (91). The vaccine can also be effective in preventing secondary complications and reducing the risk for influenza-related hospitalization and death among adults aged ≥65 years with and without high-risk medical conditions (e.g., heart disease and diabetes) (13–15,18,92). Among elderly persons not living in nursing homes or similar chronic-care facilities, influenza vaccine is 30%–70% effective in preventing hospitalization for pneumonia and influenza (15,93). Among older persons who do reside in nursing homes, influenza vaccine is most effective in preventing severe illness, secondary complications, and deaths. Among this population, the vaccine can be 50%–60% effective in preventing influenza-related hospitalization or pneumonia and 80% effective in preventing influenza-related death, although the effectiveness in preventing influenza illness often ranges from 30% to 40% (94–96).

Efficacy and Effectiveness of LAIV

Healthy Children. A randomized, double-blind, placebo-controlled trial among 1,602 healthy children initially aged 15–71 months assessed the efficacy of trivalent LAIV against culture-confirmed influenza during two seasons (97,98). This trial included subsets of 238 healthy children (163 vaccinees and 75 placebo recipients) aged 60–71 months who received 2 doses and 74 children (54 vaccinees and 20 placebo recipients) aged 60–71 months who received a single dose during season one, and a subset of 544 children (375 vaccinees and 169 placebo recipients) aged 60–84 months during season two. Children who continued in the study remained in the same study group. In season one, when vaccine and circulating virus strains were well-matched, efficacy was 93% for all participants, regardless of age, among persons receiving 2 doses of LAIV. Efficacy was 87% in the 60–71-month subset for those who received 2 doses, and was 91% in the subset for those who received 1 or 2 doses. In season two, when the A (H3N2) component was not well-matched between vaccine and circulating virus strains, efficacy was 86% overall and 87% among those aged 60–84 months. The vaccine was 92% effi-

caxious in preventing culture-confirmed influenza during the two-season study. Other results included a 27% reduction in febrile otitis media and a 28% reduction in otitis media with concomitant antibiotic use. Receipt of LAIV also resulted in a 30% lower incidence of febrile otitis media and 21% fewer febrile illnesses. Another study assessing LAIV effectiveness in children aged 18 months–18 years indicated effectiveness against medically attended acute respiratory illness (MAARI) of 18%. However, applying a validation sample of surveillance cultures with MAARI demonstrated efficacy of 92% against influenza A (H1N1) and 66% against an influenza B drift variant (99).

Healthy Adults. A randomized, double-blind, placebo-controlled trial among 4,561 healthy working adults aged 18–64 years assessed multiple endpoints, including reductions in illness, absenteeism, health-care visits, and medication use during peak and total influenza outbreak periods (100). The study was conducted during the 1997–98 influenza season, when the vaccine and circulating A (H3N2) strains were not well-matched. The study did not include testing of viruses by a laboratory. During peak outbreak periods, no difference was identified between LAIV and placebo recipients experiencing any febrile episodes. However, vaccination was associated with reductions in severe febrile illnesses of 19% and febrile upper respiratory tract illnesses of 24%. Vaccination also was associated with fewer days of illness, fewer days of work lost, fewer days with health-care–provider visits, and reduced use of prescription antibiotics and over-the-counter medications.

Among the subset of 3,637 healthy adults aged 18–49 years, LAIV recipients ($n = 2,411$) had 26% fewer febrile upper-respiratory illness episodes; 27% fewer lost work days as a result of febrile upper respiratory illness; and 18%–37% fewer days of health-care provider visits caused by febrile illness, compared with placebo recipients ($n = 1,226$). Days of antibiotic use were reduced by 41%–45% in this age subset.

Another randomized, double-blind, placebo-controlled challenge study among 92 healthy adults (LAIV, $n = 29$; placebo, $n = 31$; inactivated influenza vaccine, $n = 32$) aged 18–41 years assessed the efficacy of both LAIV and inactivated vaccine (101). The overall efficacy of LAIV and inactivated influenza vaccine in preventing laboratory-documented influenza from all three influenza strains combined was 85% and 71%, respectively, on the basis of experimental challenge by viruses to which study participants were susceptible before vaccination. The difference in efficacy between the two vaccines was not statistically significant.

Cost-Effectiveness of Influenza Vaccine

Influenza vaccination can reduce both health-care costs and productivity losses associated with influenza illness. Economic

studies of influenza vaccination of persons aged ≥ 65 years conducted in the United States have reported overall societal cost savings and substantial reductions in hospitalization and death (15,93,102). Studies of adults aged < 65 years have reported that vaccination can reduce both direct medical costs and indirect costs from work absenteeism (8,10–12,75,103). Reductions of 13%–44% in health-care–provider visits, 18%–45% in lost workdays, 18%–28% in days working with reduced effectiveness, and 25% in antibiotic use for influenza-associated illnesses have been reported (10,12,104,105). One cost-effectiveness analysis estimated a cost of approximately \$60–\$4,000/illness averted among healthy persons aged 18–64 years, depending on the cost of vaccination, the influenza attack rate, and vaccine effectiveness against ILI (75). Another cost-benefit economic study estimated an average annual savings of \$13.66/person vaccinated (106). In the second study, 78% of all costs prevented were costs from lost work productivity, whereas the first study did not include productivity losses from influenza illness. Economic studies specifically evaluating the cost-effectiveness of vaccinating persons aged 50–64 years are not available, and the number of studies that examine the economics of routinely vaccinating children with inactivated or live, attenuated vaccine are limited (8,107–110). However, in a study of inactivated vaccine that included all age groups, cost utility (i.e., cost per year of healthy life gained) improved with increasing age and among those with chronic medical conditions (8). Among persons aged ≥ 65 years, vaccination resulted in a net savings per quality-adjusted life year (QALY) gained and resulted in costs of \$23–\$256/QALY among younger age groups. Additional studies of the relative cost-effectiveness and cost utility of influenza vaccination among children and among adults aged < 65 years are needed and should be designed to account for year-to-year variations in influenza attack rates, illness severity, and vaccine efficacy when evaluating the long-term costs and benefits of annual vaccination.

Vaccination Coverage Levels

One of the national health objectives for 2010 is to achieve vaccination coverage for 90% of persons aged ≥ 65 years (objective no. 14-29a) (111). Among persons aged ≥ 65 years, influenza vaccination levels increased from 33% in 1989 (112) to 66% in 1999 (113), surpassing the *Healthy People 2000* objective of 60% (114). Vaccine coverage in this group reached the highest levels recorded (68%) during the 1999–00 influenza season, using the percentage of adults reporting influenza vaccination during the previous 12 months who participated in the National Health Interview Survey (NHIS) during the first and second quarters of each calendar year as a proxy measure of influenza vaccine coverage for the previous

influenza season (113). Possible reasons for the increase in influenza vaccination levels among persons aged ≥ 65 years through the 1999–00 influenza season include 1) greater acceptance of preventive medical services by practitioners; 2) increased delivery and administration of vaccine by health-care providers and sources other than physicians; 3) new information regarding influenza vaccine effectiveness, cost-effectiveness, and safety; and 4) initiation of Medicare reimbursement for influenza vaccination in 1993 (8,14,15,94,95,115,116). Vaccine coverage increased more rapidly through the mid-1990s than during subsequent seasons (average annual percentage increase of 4% from 1988–89 to 1996–97 versus 1% from 1996–97 to 1999–00) and has remained relatively stable since 2000.

Estimated national influenza vaccine coverage in 2003 among persons aged ≥ 65 years and 50–64 years was 66% and 37%, respectively, based on 2003 NHIS data (Table 2). The estimated vaccination coverage among adults with high-risk conditions aged 18–49 years and 50–64 years was 24% and 46%, respectively, substantially lower than the *Healthy People*

2000 and 2010 objective of 60% (111,114). Continued annual monitoring is needed to determine the effects of vaccine supply delays and shortages, changes in influenza vaccination recommendations and target groups for vaccination, reimbursement rates for vaccine and vaccine administration, and other factors related to vaccination coverage among adults and children. New strategies to improve coverage will be needed to achieve the *Healthy People 2010* objective (21).

Reducing racial and ethnic health disparities, including disparities in vaccination coverage, is an overarching national goal (111). Although estimated influenza vaccination coverage for the 1999–00 season reached the highest levels recorded among older black, Hispanic, and white populations, vaccination levels among blacks and Hispanics continue to lag behind those among whites (113,117). Estimated vaccination coverage levels based on 2003 NHIS data among persons aged ≥ 65 years were 69% among non-Hispanic whites, 48% among non-Hispanic blacks, and 45% among Hispanics (CDC, National Immunization Program, unpublished data, 2005). Additional strate-

TABLE 2. Influenza vaccination coverage rates among adult target* population groups — National Health Interview Survey (NHIS), United States, 2003

Population group	Crude sample size	Weighted sample size	Influenza vaccination rate	
			%	(95% CI†)
All aged 50–64 yrs	6,666	46,000,500	36.8	(35.4–38.2)
All aged ≥ 65 yrs	5,662	33,677,900	65.5	(64.1–66.9)
Persons with high-risk conditions§				
Aged 18–64 yrs	4,347	29,746,400	34.2	(32.5–35.9)
Aged 18–49 yrs	2,341	16,324,700	24.2	(22.1–26.4)
Aged 50–64 yrs	2,006	13,421,800	46.3	(43.7–49.0)
Persons without high-risk conditions§				
Aged 18–49 yrs	15,654	113,504,600	15.8	(15.1–16.5)
Aged 50–64 yrs	4,637	32,425,100	32.7	(31.2–34.3)
Pregnant women¶	315	2,339,600	12.8	(9.0–17.9)
Health-care workers**	2,146	14,604,000	40.1	(37.5–42.7)
Household contacts of persons at high risk, including children aged <2 yrs††				
Aged 18–49 yrs	2,501	20,404,000	14.9	(13.4–16.6)
Aged 50–64 yrs	489	4,113,400	38.4	(33.6–43.5)

* As recommended by the Advisory Committee on Immunization Practices.

† Confidence interval.

§ Persons categorized as being at high risk for influenza-related complications self-reported one or more of the following: 1) ever being told by a physician they had diabetes, emphysema, coronary heart disease, angina, heart attack, or other heart condition; 2) having a diagnosis of cancer during the previous 12 months (excluding nonmelanoma skin cancer) or ever being told by a physician they have lymphoma, leukemia, or blood cancer during the previous 12 months; 3) being told by a physician they have chronic bronchitis or weak or failing kidneys; or 4) reporting an asthma episode or attack during the preceding 12 months.

¶ Aged 18–44 years, pregnant at the time of the survey and without high-risk conditions.

** Adults were classified as health-care workers if they were currently employed in a health-care occupation or in a health-care–industry setting, on the basis of standard occupation and industry categories recoded in groups by CDC’s National Center for Health Statistics.

†† Interviewed adult in each household containing at least one of the following: a child aged <2 years, an adult aged ≥ 65 years, or any person aged 2–17 years at high risk (see previous footnote §). To obtain information on household composition and high-risk status of household members, the sampled adult, child, and person files from NHIS were merged. Interviewed adults who were health-care workers or who had high-risk conditions were excluded. Information could not be assessed regarding high-risk status of other adults aged 18–64 years in the household, thus, certain adults 18–64 years who live with an adult aged 18–64 years at high risk were not included in the analysis.

gies are needed to achieve the *Healthy People 2010* objectives among all racial and ethnic groups.

In 1997 and 1998, vaccination coverage estimates among nursing home residents were 64%–82% and 83%, respectively (118,119). The *Healthy People 2010* goal is to achieve influenza vaccination of 90% among nursing home residents, an increase from the *Healthy People 2000* goal of 80% (111,114).

Reported vaccination levels are low among children at increased risk for influenza complications. One study conducted among patients in health maintenance organizations reported influenza vaccination percentages ranging from 9% to 10% among children with asthma (120). A 25% vaccination level was reported among children with severe to moderate asthma who attended an allergy and immunology clinic (121). However, a study conducted in a pediatric clinic demonstrated an increase in the vaccination percentage of children with asthma or reactive airways disease from 5% to 32% after implementing a reminder/recall system (122). One study reported 79% vaccination coverage among children attending a cystic fibrosis treatment center (123). Data from BRFSS collected in February 2005 indicated 48% vaccination coverage for 1 or more doses among children aged 6–23 months and 35% coverage among children aged 2–17 years who had one or more high-risk medical conditions during the 2004–05 season (124). Increasing vaccination coverage among persons who have high-risk conditions and are aged <65 years, including children at high risk, is the highest priority for expanding influenza vaccine use.

Annual vaccination is recommended for health-care workers. Nonetheless, NHIS reported vaccination coverage of only 40% among health-care workers in the 2003 survey (CDC, National Immunization Program, unpublished data, 2005). Vaccination of health-care workers has been associated with reduced work absenteeism (9) and fewer deaths among nursing home patients (125,126) and is a high priority for reducing the impact of influenza in health-care settings and for expanding influenza vaccine use (127,128).

Limited information is available regarding use of influenza vaccine among pregnant women. Among women aged 18–44 years without diabetes responding to the 2001 BRFSS, those who were pregnant were less likely to report influenza vaccination during the previous 12 months (13.7%) than those not pregnant (16.8%) (122,129). Only 13% of pregnant women reported vaccination according to 2003 NHIS data, excluding pregnant women who reported diabetes, heart disease, lung disease, and other selected high-risk conditions (CDC, National Immunization Program, unpublished data, 2004) (Table 2). These data indicate low compliance with the ACIP recommendations for pregnant women. In a study of influenza vaccine acceptance by pregnant women, 71% who

were offered the vaccine chose to be vaccinated (130). However, a 1999 survey of obstetricians and gynecologists determined that only 39% administered influenza vaccine to obstetric patients, although 86% agreed that pregnant women's risk for influenza-related morbidity and mortality increases during the last two trimesters (131).

Recent data indicate that self-report of influenza vaccination among adults, compared with extraction from the medical record, is both sensitive and specific (132). Patient self-reports should be accepted as evidence of influenza vaccination in clinical practice (132). However, information on the validity of parents' reports of pediatric influenza vaccination is not yet available.

Recommendations for Using Inactivated and Live, Attenuated Influenza Vaccines

Both the inactivated influenza vaccine and LAIV can be used to reduce the risk for influenza. LAIV is approved for use among healthy persons aged 5–49 years. Inactivated influenza vaccine is approved for persons aged ≥6 months, including those with high-risk conditions (see following sections on inactivated influenza vaccine and live, attenuated influenza vaccine).

Target Groups for Vaccination

Persons at Increased Risk for Complications

Vaccination with inactivated influenza vaccine is recommended for the following persons who are at increased risk for complications from influenza:

- persons aged ≥65 years;
- residents of nursing homes and other chronic-care facilities that house persons of any age who have chronic medical conditions;
- adults and children who have chronic disorders of the pulmonary or cardiovascular systems, including asthma (hypertension is not considered a high-risk condition);
- adults and children who have required regular medical follow-up or hospitalization during the preceding year because of chronic metabolic diseases (including diabetes mellitus), renal dysfunction, hemoglobinopathies, or immunosuppression (including immunosuppression caused by medications or by human immunodeficiency virus [HIV]);
- adults and children who have any condition (e.g., cognitive dysfunction, spinal cord injuries, seizure disorders, or other neuromuscular disorders) that can compromise

respiratory function or the handling of respiratory secretions or that can increase the risk for aspiration;

- children and adolescents (aged 6 months–18 years) who are receiving long-term aspirin therapy and, therefore, might be at risk for experiencing Reye syndrome after influenza infection;
- women who will be pregnant during the influenza season; and
- children aged 6–23 months.

In 2004, approximately 88 million persons in the United States were included in one or more of these target groups, including 36 million persons aged ≥ 65 years, 1.6 million long-term-care facility residents, 6 million children aged 6–23 months, 42 million persons aged 2–64 years with one or more conditions associated with an increased risk for influenza-related complications, and 4 million pregnant women (CDC, National Immunization Program, unpublished data, 2005).

Persons Aged 50–64 Years

Vaccination is recommended for persons aged 50–64 years because this group has an increased prevalence of persons with high-risk conditions. In 2002, approximately 43.6 million persons in the United States were aged 50–64 years, of whom 13.5 million (34%) had one or more high-risk medical conditions (133). Influenza vaccine has been recommended for this entire age group to increase the low vaccination rates among persons in this age group with high-risk conditions (see preceding section). Age-based strategies are more successful in increasing vaccine coverage than patient-selection strategies based on medical conditions. Persons aged 50–64 years without high-risk conditions also receive benefit from vaccination in the form of decreased rates of influenza illness, decreased work absenteeism, and decreased need for medical visits and medication, including antibiotics (9–12). Furthermore, 50 years is an age when other preventive services begin and when routine assessment of vaccination and other preventive services has been recommended (134,135).

Persons Who Can Transmit Influenza to Those at High Risk

Persons who are clinically or subclinically infected can transmit influenza virus to persons at high risk for complications from influenza. Decreasing transmission of influenza from caregivers and household contacts to persons at high risk might reduce influenza-related deaths among persons at high risk. Evidence from two studies indicates that vaccination of health-care workers is associated with decreased deaths among nursing home patients (125,126), and hospital-based influenza outbreaks frequently occur where unvaccinated health-care workers are employed. Administration of LAIV has been dem-

onstrated to reduce MAARI in contacts of vaccine recipients (136), and to reduce ILI-related economic and medical consequences (such as work days lost and number of health-care provider visits). In addition to health-care workers, additional groups that can transmit influenza to high-risk persons and that should be vaccinated include

- employees of assisted living and other residences for persons in groups at high risk;
- persons who provide home care to persons in groups at high risk; and
- household contacts (including children) of persons in groups at high risk.

In addition, because children aged 0–23 months are at increased risk for influenza-related hospitalization (54–56), vaccination is recommended for their household contacts and out-of-home caregivers, particularly for contacts of children aged 0–5 months, because influenza vaccines have not been approved by FDA for use among children aged <6 months (see Healthy Young Children).

Healthy persons aged 5–49 years in these groups who are not contacts of severely immunosuppressed persons (see Live, Attenuated Influenza Vaccine Recommendations) can receive either LAIV or inactivated influenza vaccine. All other persons in this group should receive inactivated influenza vaccine.

Health-Care Workers

All health-care workers should be vaccinated against influenza annually (128). Facilities that employ health-care workers are strongly encouraged to provide vaccine to workers by using approaches that maximize vaccination rates. This will protect health-care workers, their patients, and communities, and will improve prevention of influenza-associated disease, patient safety, and will reduce disease burden. Influenza vaccination rates among health-care workers should be regularly measured and reported. Although vaccination rates for health-care workers are typically <40%, with moderate effort, organized campaigns can attain higher rates of vaccination among this population (127,137). Currently, seven states have legislation requiring annual influenza vaccination of health-care workers or the signing of an informed declination (128), and 15 states have regulations regarding vaccination of health-care workers in long-term-care facilities (138). Physicians, nurses, and other workers in both hospital and outpatient-care settings, including medical emergency-response workers (e.g., paramedics and emergency medical technicians), should be vaccinated, as should employees of nursing home and chronic-care facilities who have contact with patients or residents.

Additional Information Regarding Vaccination of Specific Populations

Pregnant Women

Influenza-associated excess deaths among pregnant women were documented during the pandemics of 1918–19 and 1957–58 (49,139–141). Case reports and limited studies also indicate that pregnancy can increase the risk for serious medical complications of influenza (142–146). An increased risk might result from 1) increases in heart rate, stroke volume, and oxygen consumption; 2) decreases in lung capacity; and 3) changes in immunologic function during pregnancy. A study of the effect of influenza during 17 interpandemic influenza seasons demonstrated that the relative risk for hospitalization for selected cardiorespiratory conditions among pregnant women enrolled in Medicaid increased from 1.4 during weeks 14–20 of gestation to 4.7 during weeks 37–42, in comparison with women who were 1–6 months postpartum (147). Women in their third trimester of pregnancy were hospitalized at a rate (i.e., 250/100,000 pregnant women) comparable with that of nonpregnant women who had high-risk medical conditions. Researchers estimate that an average of 1–2 hospitalizations can be prevented for every 1,000 pregnant women vaccinated (147).

Because of the increased risk for influenza-related complications, women who will be pregnant during the influenza season should be vaccinated. Vaccination can occur in any trimester. One study of influenza vaccination of approximately 2,000 pregnant women demonstrated no adverse fetal effects associated with influenza vaccine (148).

Healthy Young Children

Studies indicate that rates of hospitalization are higher among young children than older children when influenza viruses are in circulation (53,55,56,149,150). The increased rates of hospitalization are comparable with rates for other groups considered at high risk for influenza-related complications. However, the interpretation of these findings has been confounded by co-circulation of respiratory syncytial viruses, which are a cause of serious respiratory viral illness among children and which frequently circulate during the same time as influenza viruses (151–153). One study assessed rates of influenza-associated hospitalizations among the entire U.S. population during 1979–2001 and calculated an average rate of approximately 108 hospitalizations per 100,000 person-years in children aged <5 years (46). Two recent studies have attempted to separate the effects of respiratory syncytial viruses and influenza viruses on rates of hospitalization among children who do not have high-risk conditions (54,55). Both studies reported that otherwise healthy children aged <2 years,

and possibly children aged 2–4 years, are at increased risk for influenza-related hospitalization compared with older healthy children (Table 1). Among the Tennessee Medicaid population during 1973–1993, healthy children aged 6 months–<3 years had rates of influenza-associated hospitalization comparable with or higher than rates among children aged 3–14 years with high-risk conditions (54,56). Another Tennessee study reported a hospitalization rate per year of 3–4/1,000 healthy children aged <2 years for laboratory-confirmed influenza (33).

Because children aged 6–23 months are at substantially increased risk for influenza-related hospitalizations, ACIP recommends vaccination of all children in this age group (154). ACIP continues to recommend influenza vaccination of persons aged ≥6 months who have high-risk medical conditions.

The current inactivated influenza vaccine is not approved by FDA for use among children aged <6 months, the pediatric group at greatest risk for influenza-related complications (54). Vaccinating their household contacts and out-of-home caregivers might decrease the probability of influenza infection among these children.

Beginning in March 2003, the group of children eligible for influenza vaccine coverage under the Vaccines for Children (VFC) program was expanded to include all VFC-eligible children aged 6–23 months and VFC-eligible children aged 2–18 years who are household contacts of children aged 0–23 months (155).

Persons Infected with HIV

Limited information is available regarding the frequency and severity of influenza illness or the benefits of influenza vaccination among persons with HIV infection (156,157). However, a retrospective study of young and middle-aged women enrolled in Tennessee's Medicaid program determined that the attributable risk for cardiopulmonary hospitalizations among women with HIV infection was higher during influenza seasons than during the peri-influenza periods. The risk for hospitalization was higher for HIV-infected women than for women with other well-recognized high-risk conditions, including chronic heart and lung diseases (158). Another study estimated that the risk for influenza-related death was 9.4–14.6/10,000 persons with acquired immunodeficiency syndrome (AIDS) compared with 0.09–0.10/10,000 among all persons aged 25–54 years and 6.4–7.0/10,000 among persons aged ≥65 years (159). Other reports indicate that influenza symptoms might be prolonged and the risk for complications from influenza increased for certain HIV-infected persons (160–162).

Inactivated influenza vaccination has been demonstrated to produce substantial antibody titers against influenza among

vaccinated HIV-infected persons who have minimal AIDS-related symptoms and high CD4+ T-lymphocyte cell counts (163–166). A limited, randomized, placebo-controlled trial determined that inactivated influenza vaccine was highly effective in preventing symptomatic, laboratory-confirmed influenza infection among HIV-infected persons with a mean of 400 CD4+ T-lymphocyte cells/mm³; a limited number of persons with CD4+ T-lymphocyte cell counts of <200 were included in that study (167). A nonrandomized study among HIV-infected persons determined that influenza vaccination was most effective among persons with >100 CD4+ cells and among those with <30,000 viral copies of HIV type-1/mL (162). Among persons who have advanced HIV disease and low CD4+ T-lymphocyte cell counts, inactivated influenza vaccine might not induce protective antibody titers (165,166); a second dose of vaccine does not improve the immune response in these persons (166,167).

One study determined that HIV RNA (ribonucleic acid) levels increased transiently in one HIV-infected person after influenza infection (168). Studies have demonstrated a transient (i.e., 2–4 week) increase in replication of HIV-1 in the plasma or peripheral blood mononuclear cells of HIV-infected persons after vaccine administration (165,169). Other studies using similar laboratory techniques have not documented a substantial increase in the replication of HIV (170–173). Deterioration of CD4+ T-lymphocyte cell counts or progression of HIV disease have not been demonstrated among HIV-infected persons after influenza vaccination compared with unvaccinated persons (166,174). Limited information is available concerning the effect of antiretroviral therapy on increases in HIV RNA levels after either natural influenza infection or influenza vaccination (156,175). Because influenza can result in serious illness and because vaccination with inactivated influenza vaccine can result in the production of protective antibody titers, vaccination will benefit HIV-infected persons, including HIV-infected pregnant women.

Breastfeeding Mothers

Influenza vaccine is safe for mothers who are breastfeeding and their infants. Breastfeeding does not adversely affect the immune response and is not a contraindication for vaccination.

Travelers

The risk for exposure to influenza during travel depends on the time of year and destination. In the tropics, influenza can occur throughout the year. In the temperate regions of the Southern Hemisphere, the majority of influenza activity occurs during April–September. In temperate climate zones of the Northern and Southern Hemispheres, travelers also can be exposed to influenza during the summer, especially when

traveling as part of large organized tourist groups (e.g., on cruise ships) that include persons from areas of the world where influenza viruses are circulating (176,177). Persons at high risk for complications of influenza who were not vaccinated with influenza vaccine during the preceding fall or winter should consider receiving influenza vaccine before travel if they plan to

- travel to the tropics,
- travel with organized tourist groups at any time of year, or
- travel to the Southern Hemisphere during April–September.

No information is available regarding the benefits of revaccinating persons before summer travel who were already vaccinated in the preceding fall. Persons at high risk who receive the previous season's vaccine before travel should be revaccinated with the current vaccine the following fall or winter. Persons aged ≥50 years and persons at high risk should consult with their physicians before embarking on travel during the summer to discuss the symptoms and risks for influenza and the advisability of carrying antiviral medications for either prophylaxis or treatment of influenza.

General Population

In addition to the groups for which annual influenza vaccination is recommended, physicians should administer influenza vaccine to any person who wishes to reduce the likelihood of becoming ill with influenza or transmitting influenza to others should they become infected (the vaccine can be administered to children aged ≥6 months), depending on vaccine availability (see Influenza Vaccine Supply). Persons who provide essential community services should be considered for vaccination to minimize disruption of essential activities during influenza outbreaks. Students or other persons in institutional settings (e.g., those who reside in dormitories) should be encouraged to receive vaccine to minimize the disruption of routine activities during epidemics.

Comparison of LAIV with Inactivated Influenza Vaccine

Both inactivated influenza vaccine and LAIV are available to reduce the risk for influenza infection and illness. However, the vaccines also differ in key ways (Table 3).

Major Similarities

LAIV and inactivated influenza vaccine contain strains of influenza viruses that are antigenically equivalent to the annually recommended strains: one influenza A (H3N2) virus, one A (H1N1) virus, and one B virus. Each year, one or more

TABLE 3. Live, attenuated influenza vaccine (LAIV) compared with inactivated influenza vaccine

Factor	LAIV	Inactivated influenza vaccine
Route of administration	Intranasal spray	Intramuscular injection
Type of vaccine	Live virus	Killed virus
No. of included virus strains		
	3 (2 influenza A, 1 influenza B)	Same as LAIV
Vaccine virus strains updated	Annually	Same as LAIV
Frequency of administration	Annually	Same as LAIV
Approved age and risk groups*	Healthy persons aged 5–49 yrs	Persons aged ≥ 6 mos
Can be administered to family members or close contacts of immunosuppressed persons not requiring a protected environment	Yes	Yes
Can be administered to family members or close contacts of immunosuppressed persons requiring a protected environment (e.g., hematopoietic stemcell transplant recipient)	Inactivated influenza vaccine preferred	Yes
Can be administered to family members or close contacts of persons at high risk but not severely immunosuppressed	Yes	Yes
Can be simultaneously administered with other vaccines	Yes [†]	Yes [§]
If not simultaneously administered, can be administered within 4 weeks of another live vaccine	Prudent to space 4 weeks apart	Yes
If not simultaneously administered, can be administered within 4 weeks of an inactivated vaccine	Yes	Yes

* Populations at high risk from complications of influenza infection include persons aged ≥ 65 years; residents of nursing homes and other chronic-care facilities that house persons with chronic medical conditions; adults and children with chronic disorders of the pulmonary or cardiovascular systems; adults and children with chronic metabolic diseases (including diabetes mellitus), renal dysfunction, hemoglobinopathies, or immunosuppression; children and adolescents receiving long-term aspirin therapy (at risk for developing Reye syndrome after wild-type influenza infection); pregnant women; and children aged 6–23 months.

[†] No data are available regarding effect on safety or efficacy.

[§] Inactivated influenza vaccine coadministration has been evaluated systematically only among adults with pneumococcal polysaccharide vaccine.

virus strains might be changed on the basis of global surveillance for influenza viruses and the emergence and spread of new strains. Viruses for both vaccines are grown in eggs. Both vaccines are administered annually to provide optimal protection against influenza infection (Table 3).

Major Differences

Inactivated influenza vaccine contains killed viruses, whereas LAIV contains live, attenuated viruses still capable of replication. LAIV is administered intranasally by sprayer, whereas inactivated influenza vaccine is administered intramuscularly by injection. LAIV is more expensive than inactivated influenza vaccine, although the price differential between inactivated vaccine and LAIV has decreased for the 2005–06 season. LAIV is approved for use among healthy persons aged 5–49 years; inactivated influenza vaccine is approved for use among persons aged ≥ 6 months, including those who are healthy and those with chronic medical conditions (Table 3).

Inactivated Influenza Vaccine Recommendations

Persons Who Should Not Be Vaccinated with Inactivated Influenza Vaccine

Inactivated influenza vaccine should not be administered to persons known to have anaphylactic hypersensitivity to eggs or to other components of the influenza vaccine without first consulting a physician (see Side Effects and Adverse Reactions). Prophylactic use of antiviral agents is an option for preventing influenza among such persons. However, persons who have a history of anaphylactic hypersensitivity to vaccine components but who are also at high risk for complications from influenza can benefit from vaccine after appropriate allergy evaluation and desensitization. Information regarding vaccine components is located in package inserts from each manufacturer. Persons with acute febrile illness usually should not be vaccinated until their symptoms have abated. However, minor illnesses with or without fever do not contraindicate use of influenza vaccine, particularly among children with mild upper-respiratory-tract infection or allergic rhinitis.

Dosage

Dosage recommendations vary according to age group (Table 4). Among previously unvaccinated children aged <9 years, 2 doses administered ≥ 1 month apart are recommended for satisfactory antibody responses. If possible, the second dose should be administered before December. If a child aged <9 years receiving vaccine for the first time does not receive a second dose of vaccine within the same season, only 1 dose of vaccine should be administered the following season. Two doses are not required at that time. Among adults, studies have indicated limited or no improvement in antibody response when a second dose is administered during the same season (178–180). Even when the current influenza vaccine contains one or more antigens administered in previous years, annual vaccination with the current vaccine is necessary because immunity declines during the year after vaccination (181,182). Vaccine prepared for a previous influenza season should not be administered to provide protection for the current season. Because of lack of vaccine efficacy data, ACIP does not recommend that a child receiving influenza vaccine for the first time be given the first dose of vaccine in the spring, followed by the second dose in the autumn of the same year.

Route

The intramuscular route is recommended for influenza vaccine. Adults and older children should be vaccinated in the

deltoid muscle. A needle length ≥ 1 inch can be considered for these age groups because needles <1 inch might be of insufficient length to penetrate muscle tissue in certain adults and older children (183).

Infants and young children should be vaccinated in the anterolateral aspect of the thigh (67). ACIP recommends a needle length of 7/8–1 inch for children aged <12 months for intramuscular vaccination into the anterolateral thigh. When injecting into the deltoid muscle among children with adequate deltoid muscle mass, a needle length of 7/8–1.25 inches is recommended (67).

Side Effects and Adverse Reactions

When educating patients regarding potential side effects, clinicians should emphasize that 1) inactivated influenza vaccine contains noninfectious killed viruses and cannot cause influenza; and 2) coincidental respiratory disease unrelated to influenza vaccination can occur after vaccination.

Local Reactions

In placebo-controlled studies among adults, the most frequent side effect of vaccination is soreness at the vaccination site (affecting 10%–64% of patients) that lasts <2 days (12,184–186). These local reactions typically are mild and rarely interfere with the person's ability to conduct usual daily activities. One blinded, randomized, cross-over study among 1,952 adults and children with asthma, demonstrated that only body aches were reported more frequently after inactivated influenza vaccine (25.1%) than placebo-injection (20.8%) (187). One study (83) reported 20%–28% of children with asthma aged 9 months–18 years with local pain and swelling, and another study (80) reported 23% of children aged 6 months–4 years with chronic heart or lung disease had local reactions. A different study (81) reported no difference in local reactions among 53 children aged 6 months–6 years with high-risk medical conditions or among 305 healthy children aged 3–12 years in a placebo-controlled trial of inactivated influenza vaccine. In a study of 12 children aged 5–32 months, no substantial local or systemic reactions were noted (188).

Systemic Reactions

Fever, malaise, myalgia, and other systemic symptoms can occur after vaccination with inactivated vaccine and most often affect persons who have had no previous exposure to the influenza virus antigens in the vaccine (e.g., young children) (189,190). These reactions begin 6–12 hours after vaccination and can persist for 1–2 days. Recent placebo-controlled trials demonstrate that among older persons and healthy young adults, administration of split-virus influenza vaccine is not

TABLE 4. Inactivated influenza vaccine* dosage, by age group — United States, 2005–06 season

Age group†	Dose	No. of doses	Route§
6–35 mos	0.25 mL	1 or 2¶	Intramuscular
3–8 yrs	0.50 mL	1 or 2¶	Intramuscular
≥ 9 yrs	0.50 mL	1	Intramuscular

* A 0.5-mL dose contains 15 mg each of A/California/7/2004 (H3N2)-like, A/New Caledonia/20/99 (H1N1)-like, and B/Shanghai/361/2002-like antigens. For the A/California/7/2004 (H3N2)-like antigen, manufacturers may use the antigenically equivalent A/New York/55/2004 virus, and for the B/Shanghai/361/2002-like antigen, manufacturers may use the antigenically equivalent B/Jilin/20/2003 virus or B/Jiangsu/10/2003 virus. Manufacturers include Sanofi Pasteur, Inc. (formerly Aventis Pasteur, Inc.) (FluZone® split virus); and Chiron (Fluvirin™ purified-surface-antigen vaccine). FluZone is approved by the Food and Drug Administration for use among persons aged ≥ 6 months. Fluvirin is approved for use in persons aged ≥ 4 years. For further product information, call Sanofi Pasteur at 800-822-2463 or Chiron at 800-244-7668.

† Because of their decreased potential for causing febrile reactions, only split-virus vaccines should be used for children aged <13 years. Whole-virus vaccine is not available in the United States. Split-virus vaccine might be labeled as *split*, *subvirion*, or *purified-surface-antigen* vaccine. Immunogenicity and side effects of split- and whole-virus vaccines are similar among adults when vaccines are administered at the recommended dosage.

§ For adults and older children, the recommended site of vaccination is the deltoid muscle. The preferred site for infants and young children is the anterolateral aspect of the thigh.

¶ Two doses administered at least 1 month apart are recommended for children aged <9 years who are receiving influenza vaccine for the first time.

associated with higher rates of systemic symptoms (e.g., fever, malaise, myalgia, and headache) when compared with placebo injections (12,184–186).

Less information from published studies is available for children, compared with adults. However, in a randomized crossover study among both children and adults with asthma, no increase in asthma exacerbations was reported for either age group (187). An analysis of 215,600 children aged <18 years and 8,476 children aged 6–23 months enrolled in one of five health maintenance organizations reported no increase in biologically plausible medically attended events during the 2 weeks after inactivated influenza vaccination, compared with control periods 3–4 weeks before and after vaccination (191). In a study of 791 healthy children (71), postvaccination fever was noted among 11.5% of children aged 1–5 years, 4.6% among children aged 6–10 years, and 5.1% among children aged 11–15 years. Among children with high-risk medical conditions, one study of 52 children aged 6 months–4 years reported fever among 27% and irritability and insomnia among 25% (80); and a study among 33 children aged 6–18 months reported that one child had irritability and one had a fever and seizure after vaccination (192). No placebo comparison was made in these studies. However, in pediatric trials of A/New Jersey/76 swine influenza vaccine, no difference was reported between placebo and split-virus vaccine groups in febrile reactions after injection, although the vaccine was associated with mild local tenderness or erythema (81).

Limited data regarding potential adverse events after influenza vaccination are available from the Vaccine Adverse Event Reporting System (VAERS). During January 1, 1991–June 30, 2004, VAERS received 1,895 reports of adverse events among children aged <18 years, including 479 reports of adverse events among children aged 6–23 months. The number of influenza vaccine doses received by children during this entire period is unknown (CDC, unpublished data, 2005). A recently published review of VAERS reports of trivalent inactivated influenza vaccine (TIV) in children aged 6–23 months documented that the most frequently reported adverse events were fever, rash, injection-site reactions, and seizures. The majority of the small total number of reported seizures appeared to be febrile (193). Because of the limitations of passive reporting systems, determining causality for specific types of adverse events, with the exception of injection-site reactions, is usually not possible by using VAERS data alone. A population-based study of TIV safety in children aged 6–23 months indicated no vaccine associated adverse events that had a plausible relationship to vaccination (194).

Health-care professionals should promptly report to VAERS all clinically significant adverse events after influenza vaccination of children, even if the health-care professional is not

certain that the vaccine caused the event. The Institute of Medicine has specifically recommended reporting of potential neurologic complications (e.g., demyelinating disorders such as Guillain-Barré syndrome [GBS]), although no evidence exists of a causal relationship between influenza vaccine and neurologic disorders in children.

Immediate — presumably allergic — reactions (e.g., hives, angioedema, allergic asthma, and systemic anaphylaxis) rarely occur after influenza vaccination (195). These reactions probably result from hypersensitivity to certain vaccine components; the majority of reactions probably are caused by residual egg protein. Although current influenza vaccines contain only a limited quantity of egg protein, this protein can induce immediate hypersensitivity reactions among persons who have severe egg allergy. Persons who have had hives or swelling of the lips or tongue, or who have experienced acute respiratory distress or collapse after eating eggs should consult a physician for appropriate evaluation to help determine if vaccine should be administered. Persons who have documented immunoglobulin E (IgE)-mediated hypersensitivity to eggs, including those who have had occupational asthma or other allergic responses to egg protein, might also be at increased risk for allergic reactions to influenza vaccine, and consultation with a physician should be considered. Protocols have been published for safely administering influenza vaccine to persons with egg allergies (196–198).

Hypersensitivity reactions to any vaccine component can occur. Although exposure to vaccines containing thimerosal can lead to induction of hypersensitivity, the majority of patients do not have reactions to thimerosal when it is administered as a component of vaccines, even when patch or intradermal tests for thimerosal indicate hypersensitivity (199,200). When reported, hypersensitivity to thimerosal usually has consisted of local, delayed hypersensitivity reactions (199).

Guillain-Barré Syndrome

The 1976 swine influenza vaccine was associated with an increased frequency of GBS (201,202). Among persons who received the swine influenza vaccine in 1976, the rate of GBS was <10 cases/1 million persons vaccinated. The risk for influenza vaccine-associated GBS is higher among persons aged ≥25 years than persons <25 years (201). Evidence for a causal relation of GBS with subsequent vaccines prepared from other influenza viruses is unclear. Obtaining strong epidemiologic evidence for a possible limited increase in risk is difficult for such a rare condition as GBS, which has an annual incidence of 10–20 cases/1 million adults (203). More definitive data probably will require using other methodologies (e.g., laboratory studies of the pathophysiology of GBS).

During three of four influenza seasons studied during 1977–1991, the overall relative risk estimates for GBS after influenza vaccination were slightly elevated but were not statistically significant in any of these studies (204–206). However, in a study of the 1992–93 and 1993–94 seasons, the overall relative risk for GBS was 1.7 (95% CI = 1.0–2.8; $p = 0.04$) during the 6 weeks after vaccination, representing approximately 1 additional case of GBS/1 million persons vaccinated. The combined number of GBS cases peaked 2 weeks after vaccination (207). Thus, investigations to date have not documented a substantial increase in GBS associated with influenza vaccines (other than the swine influenza vaccine in 1976), and that, if influenza vaccine does pose a risk, it is probably slightly more than one additional case/1 million persons vaccinated. Recent data from VAERS has documented decreased reporting of post influenza vaccine GBS across age groups, despite overall increased reporting for influenza vaccine (208). Cases of GBS after influenza infection have been reported, but no epidemiologic studies have documented such an association (209,210). Substantial evidence exists that multiple infectious illnesses, most notably *Campylobacter jejuni*, and upper respiratory tract infections are associated with GBS (203,211–213).

Even if GBS were a true side effect of vaccination in the years after 1976, the estimated risk for GBS of approximately 1 additional case/1 million persons vaccinated is substantially less than the risk for severe influenza, which can be prevented by vaccination among all age groups, especially persons aged ≥ 65 years and those who have medical indications for influenza vaccination (Table 1) (see Hospitalizations and Deaths from Influenza). The potential benefits of influenza vaccination in preventing serious illness, hospitalization, and death substantially outweigh the possible risks for experiencing vaccine-associated GBS. The average case fatality ratio for GBS is 6% and increases with age (203,214). No evidence indicates that the case fatality ratio for GBS differs among vaccinated persons and those not vaccinated.

The incidence of GBS among the general population is low, but persons with a history of GBS have a substantially greater likelihood of subsequently experiencing GBS than persons without such a history (204,215). Thus, the likelihood of coincidentally experiencing GBS after influenza vaccination is expected to be greater among persons with a history of GBS than among persons with no history of this syndrome. Whether influenza vaccination specifically might increase the risk for recurrence of GBS is unknown; therefore, avoiding vaccinating persons who are not at high risk for severe influenza complications and who are known to have experienced GBS within 6 weeks after a previous influenza vaccination is prudent. As an alternative, physicians might consider using influenza an-

tiviral chemoprophylaxis for these persons. Although data are limited, for the majority of persons who have a history of GBS and who are at high risk for severe complications from influenza, the established benefits of influenza vaccination justify yearly vaccination.

Live, Attenuated Influenza Vaccine Recommendations

Background

Description and Action Mechanisms. LAIVs have been in development since the 1960s in the United States, where they have been evaluated as mono-, bi-, and trivalent formulations (216–218). The LAIV licensed for use in the United States beginning in 2003 is produced by MedImmune, Inc. (Gaithersburg, Maryland; <http://www.medimmune.com>) and marketed under the name FluMist™. It is a live, trivalent, intranasally administered vaccine that is

- attenuated, producing mild or no signs or symptoms related to influenza virus infection;
- temperature-sensitive, a property that limits the replication of the vaccine viruses at 38°C–39°C, and thus restricts LAIV viruses from replicating efficiently in human lower airways; and
- cold-adapted, replicating efficiently at 25°C, a temperature that is permissive for replication of LAIV viruses, but restrictive for replication of different wild-type viruses.

In animal studies, LAIV viruses replicate in the mucosa of the nasopharynx, inducing protective immunity against viruses included in the vaccine, but replicate inefficiently in the lower airways or lungs.

The first step in developing an LAIV was the derivation of two stably attenuated master donor viruses (MDV), one for type A and one for type B influenza viruses. The two MDVs each acquired the cold-adapted, temperature-sensitive, attenuated phenotypes through serial passage in viral culture conducted at progressively lower temperatures. The vaccine viruses in LAIV are reassortant viruses containing genes from these MDVs that confer attenuation, temperature sensitivity, and cold adaptation and genes from the recommended contemporary wild-type influenza viruses, encoding the surface antigens hemagglutinin (HA) and neuraminidase (NA). Thus, MDVs provide the stably attenuated vehicles for presenting influenza HA and NA antigens, to which the protective antibody response is directed, to the immune system. The reassortant vaccine viruses are grown in embryonated hens eggs. After the vaccine is formulated and inserted into individual sprayers for nasal administration, the vaccine must be stored at -15°C or colder.

The immunogenicity of the approved LAIV has been assessed in multiple studies (102,219–224), which included approximately 100 children aged 5–17 years, and approximately 300 adults aged 18–49 years. LAIV virus strains replicate primarily in nasopharyngeal epithelial cells. The protective mechanisms induced by vaccination with LAIV are not completely understood but appear to involve both serum and nasal secretory antibodies. No single laboratory measurement closely correlates with protective immunity induced by LAIV.

Shedding and Transmission of Vaccine Viruses. Available data indicate that both children and adults vaccinated with LAIV can shed vaccine viruses for ≥ 2 days after vaccination, although in lower titers than typically occur with shedding of wild-type influenza viruses. Shedding should not be equated with person-to-person transmission of vaccine viruses, although, in rare instances, shed vaccine viruses can be transmitted from vaccinees to nonvaccinated persons.

One unpublished study in a child care center setting—assessed transmissibility of vaccine viruses from 98 vaccinated to 99 unvaccinated subjects, all aged 8–36 months. Eighty percent of vaccine recipients shed one or more virus strains, with a mean of 7.6 days' duration (225). One vaccine type influenza type B isolate was recovered from a placebo recipient and was confirmed to be vaccine-type virus. The type B isolate retained the cold-adapted, temperature-sensitive, attenuated phenotype, and it possessed the same genetic sequence as a virus shed from a vaccine recipient in the same children's play group. The placebo recipient from whom the influenza type B vaccine virus was isolated did not exhibit symptoms that were different from those experienced by vaccine recipients. The estimated probability of acquiring vaccine virus after close contact with a single LAIV recipient in this child care population was 0.58%–2.4%.

One study assessing shedding of vaccine viruses in 20 healthy vaccinated adults aged 18–49 years demonstrated that the majority of shedding occurred within the first 3 days after vaccination, although one subject was noted to shed virus on day 7 after vaccine receipt. No subject shed vaccine viruses ≥ 10 days after vaccination. Duration or type of symptoms associated with receipt of LAIV did not correlate with duration of shedding of vaccine viruses. Person-to-person transmission of vaccine viruses was not assessed in this study (226).

Another study assessing shedding of vaccine viruses in 14 healthy adults aged 18–49 years indicated that 50% of these adults had viral antigen detected by direct immunofluorescence or rapid antigen tests within 7 days of vaccination. Most viral shedding was detected on day 2 or 3. Person-to-person transmission of vaccine viruses was not assessed in this study (227).

Stability of Vaccine Viruses. In clinical trials, viruses shed by vaccine recipients have been phenotypically stable. In one study, nasal and throat swab specimens were collected from 17 study participants for 2 weeks after vaccine receipt (228). Virus isolates were analyzed by multiple genetic techniques. All isolates retained the LAIV genotype after replication in the human host, and all retained the cold-adapted and temperature-sensitive phenotypes.

Using LAIV

LAIV is an option for vaccination of healthy persons aged 5–49 years, including health-care workers and other persons in close contact with groups at high risk and those wanting to avoid influenza. During periods when inactivated vaccine is in short supply, use of LAIV is encouraged when feasible for eligible persons (including health-care workers) because use of LAIV by these persons might increase availability of inactivated vaccine for persons in groups at high risk. Possible advantages of LAIV include its potential to induce a broad mucosal and systemic immune response, its ease of administration, and the acceptability of an intranasal rather than intramuscular route of administration.

Persons Who Should Not Be Vaccinated with LAIV

The following populations should not be vaccinated with LAIV:

- persons aged < 5 years or those aged ≥ 50 years;*
- persons with asthma, reactive airways disease, or other chronic disorders of the pulmonary or cardiovascular systems; persons with other underlying medical conditions, including such metabolic diseases as diabetes, renal dysfunction, and hemoglobinopathies; or persons with known or suspected immunodeficiency diseases or who are receiving immunosuppressive therapies;*
- children or adolescents receiving aspirin or other salicylates (because of the association of Reye syndrome with wild-type influenza infection);*
- persons with a history of GBS;
- pregnant women;*
- persons with a history of hypersensitivity, including anaphylaxis, to any of the components of LAIV or to eggs.

Close Contacts of Persons at High Risk for Complications from Influenza

Close contacts of persons at high risk for complications from influenza should receive influenza vaccine to reduce transmission of wild-type influenza viruses to persons at high risk.

* These persons should receive inactivated influenza vaccine.

ACIP has not indicated a preference for inactivated influenza vaccine use by health-care workers or other persons who have close contact with persons with lesser degrees of immunosuppression (e.g., persons with diabetes, persons with asthma taking corticosteroids, or persons infected with HIV) or for inactivated influenza vaccine use by health-care workers or other healthy persons aged 5–49 years in close contact with all other groups at high risk. Use of inactivated influenza vaccine is preferred for vaccinating household members, health-care workers, and others who have close contact with severely immunosuppressed persons (e.g., patients with hematopoietic stem cell transplants) during those periods in which the immunosuppressed person requires care in a protective environment. The rationale for not using LAIV among health-care workers caring for such patients is the theoretical risk that a live, attenuated vaccine virus could be transmitted to the severely immunosuppressed person. If a health-care worker receives LAIV, that worker should refrain from contact with severely immunosuppressed patients for 7 days after vaccine receipt. Hospital visitors who have received LAIV should refrain from contact with severely immunosuppressed persons for 7 days after vaccination; however, such persons need not be excluded from visitation of patients who are not severely immunosuppressed.

Personnel Who May Administer LAIV

Low-level introduction of vaccine viruses into the environment is likely unavoidable when administering LAIV. The risk for acquiring vaccine viruses from the environment is unknown but likely to be limited. Severely immunosuppressed persons should not administer LAIV. However, other persons at high risk for influenza complications may administer LAIV. These include persons with underlying medical conditions placing them at high risk or who are likely to be at risk, including pregnant women, persons with asthma, and persons aged ≥ 50 years.

LAIV Dosage and Administration

LAIV is intended for intranasal administration only and should not be administered by the intramuscular, intradermal, or intravenous route. LAIV must be thawed before administration. This can be accomplished by holding an individual sprayer in the palm of the hand until thawed, with subsequent immediate administration. Alternatively, the vaccine can be thawed in a refrigerator and stored at 2°C–8°C for ≤ 24 hours before use. Vaccine should not be refrozen after thawing. LAIV is supplied in a prefilled single-use sprayer containing 0.5 mL of vaccine. Approximately 0.25 mL (i.e., half of the total sprayer contents) is sprayed into the first nostril while the recipient is in the upright position. An attached

dose-divider clip is removed from the sprayer to administer the second half of the dose into the other nostril. If the vaccine recipient sneezes after administration, the dose should not be repeated.

LAIV should be administered annually according to the following schedule:

- Children aged 5–8 years previously unvaccinated at any time with either LAIV or inactivated influenza vaccine should receive 2 doses[†] of LAIV separated by 6–10 weeks.
- Children aged 5–8 years previously vaccinated at any time with either LAIV or inactivated influenza vaccine should receive 1 dose of LAIV. They do not require a second dose.
- Persons aged 9–49 years should receive 1 dose of LAIV.

LAIV can be administered to persons with minor acute illnesses (e.g., diarrhea or mild upper respiratory tract infection with or without fever). However, if clinical judgment indicates nasal congestion is present that might impede delivery of the vaccine to the nasopharyngeal mucosa, deferral of administration should be considered until resolution of the illness.

Whether concurrent administration of LAIV with other vaccines affects the safety or efficacy of either LAIV or the simultaneously administered vaccine is unknown. In the absence of specific data indicating interference, following the ACIP general recommendations for immunization is prudent (67). Inactivated vaccines do not interfere with the immune response to other inactivated vaccines or to live vaccines. An inactivated vaccine can be administered either simultaneously or at any time before or after LAIV. Two live vaccines not administered on the same day should be administered ≥ 4 weeks apart when possible.

LAIV and Use of Influenza Antiviral Medications

The effect on safety and efficacy of LAIV coadministration with influenza antiviral medications has not been studied. However, because influenza antivirals reduce replication of influenza viruses, LAIV should not be administered until 48 hours after cessation of influenza antiviral therapy, and influenza antiviral medications should not be administered for 2 weeks after receipt of LAIV.

LAIV Storage

LAIV must be stored at -15°C or colder. A manufacturer-supplied freezer box was formerly required for storage of LAIV in a frost-free freezer; however, the freezer box is now optional, and LAIV may now be stored in frost-free freezers without

[†] One dose equals 0.5 mL, divided equally between each nostril.

using a freezer box. LAIV can be thawed in a refrigerator and stored at 2°C–8°C for ≤60 hours before use. It should not be refrozen after thawing.

Side Effects and Adverse Reactions

Twenty prelicensure clinical trials assessed the safety of the approved LAIV. In these combined studies, approximately 28,000 doses of the vaccine were administered to approximately 20,000 subjects. A subset of these trials were randomized, placebo-controlled studies in which an estimated 4,000 healthy children aged 5–17 years and 2,000 healthy adults aged 18–49 years were vaccinated. The incidence of adverse events possibly complicating influenza (e.g., pneumonia, bronchitis, bronchiolitis, or central nervous system events) was not statistically different among LAIV and placebo recipients aged 5–49 years. LAIV is made from attenuated viruses and does not cause influenza in vaccine recipients.

Children. In a subset of healthy children aged 60–71 months from one clinical trial (97,98), certain signs and symptoms were reported more often among LAIV recipients after the first dose ($n = 214$) than placebo recipients ($n = 95$) (e.g., runny nose, 48.1% versus 44.2%; headache, 17.8% versus 11.6%; vomiting, 4.7% versus 3.2%; and myalgias, 6.1% versus 4.2%), but these differences were not statistically significant. In other trials, signs and symptoms reported after LAIV administration have included runny nose or nasal congestion (20%–75%), headache (2%–46%), fever (0%–26%), vomiting (3%–13%), abdominal pain (2%), and myalgias (0%–21%) (219,222,224,229–231). These symptoms were associated more often with the first dose and were self-limited. Unpublished data from a study including subjects aged 1–17 years indicated an increase in asthma or reactive airways disease in the subset aged 12–59 months. Because of this, LAIV is not approved for use among children aged <60 months.

Adults. Among adults, runny nose or nasal congestion (28%–78%), headache (16%–44%), and sore throat (15%–27%) have been reported more often among vaccine recipients than placebo recipients (100,232,233). In one clinical trial (100) among a subset of healthy adults aged 18–49 years, signs and symptoms reported more frequently among LAIV recipients ($n = 2,548$) than placebo recipients ($n = 1,290$) within 7 days after each dose included cough (13.9% versus 10.8%); runny nose (44.5% versus 27.1%); sore throat (27.8% versus 17.1%); chills (8.6% versus 6.0%); and tiredness/weakness (25.7% versus 21.6%).

Safety Among Groups at High Risk from Influenza-Related Morbidity. Until additional data are acquired and analyzed, persons at high risk for experiencing complications from influenza infection (e.g., immunocompromised patients; patients with asthma, cystic fibrosis, or chronic obstructive

pulmonary disease; or persons aged ≥65 years) should not be vaccinated with LAIV. Protection from influenza among these groups should be accomplished by using inactivated influenza vaccine.

Serious Adverse Events. Serious adverse events among healthy children aged 5–17 years or healthy adults aged 18–49 years occurred at a rate of <1%. Surveillance should continue for adverse events that might not have been detected in previous studies. A preliminary review of reports to VAERS after distribution of approximately 800,000 doses during the 2003–04 influenza season did not reveal any substantial new safety concerns (234). Health-care professionals should promptly report all clinically significant adverse events after LAIV administration to VAERS, as recommended for inactivated influenza vaccine.

Recommended Vaccines for Different Age Groups

When vaccinating children aged 6 months–3 years, health-care providers should use inactivated influenza vaccine that has been approved by FDA for this age group. Inactivated influenza vaccine from Sanofi Pasteur, Inc., (FluZone split-virus) is approved for use among persons aged ≥6 months. Inactivated influenza vaccine from Chiron (Fluvirin) is labeled in the United States for use among persons aged ≥4 years because data to demonstrate efficacy among younger persons have not been provided to FDA. Live, attenuated influenza vaccine from MedImmune (FluMist) is approved for use by healthy persons aged 5–49 years (Table 5).

Timing of Annual Influenza Vaccination

The annual supply of influenza vaccine and the timing of its distribution cannot be guaranteed in any year. Information regarding the supply of 2005–06 vaccine might not be available until late summer or early fall 2005. To allow vaccine providers to plan for the upcoming vaccination season, taking into account the yearly possibility of vaccine delays or shortages and the need to ensure vaccination of persons at high risk and their contacts, ACIP recommends that inactivated influenza vaccine campaigns conducted in October focus primarily on persons at increased risk for influenza complications and their contacts, including health-care workers. Campaigns conducted in November and later should continue to vaccinate persons at high risk and their contacts, but also vaccinate other persons who wish to decrease their risk for influenza infection. Vaccination for all groups should continue into December and beyond. CDC and other public health agencies will assess the vaccine supply on a continuing

TABLE 5. Approved influenza vaccines for different age groups — United States

Vaccine	6 mos–3 yrs	4 yrs	5–49 yrs	≥50 yrs
FluZone [®] (Sanofi Pasteur, Inc.)	X*	X	X	X
Fluvirin [™] (Chiron)		X	X	X
FluMist [™] (MedImmune, Inc.)			X	

* Children aged 6–35 months should receive 0.25 mL/dose. Persons aged >35 months should receive 0.50 mL/dose.

basis throughout the manufacturing period and will make recommendations preceding the 2005–06 influenza season regarding the need for tiered timing of inactivated influenza vaccination of different risk groups. Because LAIV is approved for use in healthy persons 5–49 years, its use has not been subject to tiered timing.

Vaccination Before October

To avoid missed opportunities for vaccination of persons at high risk for serious complications, such persons should be offered vaccine beginning in September during routine health-care visits or during hospitalizations, if vaccine is available. In facilities housing older persons (e.g., nursing homes), vaccination before October typically should be avoided because antibody levels in such persons can begin to decline within a limited time after vaccination (235). In addition, children aged <9 years who have not been previously vaccinated and who need 2 doses before the start of the influenza season can receive their first dose in September so that both doses of the most up-to-date vaccine can be administered before the onset of influenza activity. For previously vaccinated children, 2 doses are needed to provide optimal protection against influenza.

Vaccination in October and November

The optimal time to vaccinate is usually during October–November. ACIP recommends that vaccine providers focus their vaccination efforts in October and earlier primarily on persons aged ≥50 years, persons aged <50 years at increased risk for influenza-related complications (including children aged 6–23 months), household contacts of persons at high risk (including out-of-home caregivers and household contacts of children aged 0–23 months), and health-care workers. Vaccination of children aged <9 years who are receiving vaccine for the first time should also begin in October or earlier because those persons need a booster dose 1 month after the initial dose. Efforts to vaccinate other persons who wish to decrease their risk for influenza infection should begin in

November; however, if such persons request vaccination in October, vaccination should not be deferred, unless vaccine supplies dictate otherwise. Materials to assist providers in prioritizing early vaccine are available at <http://www.cdc.gov/flu/professionals/vaccination/index.htm> (see also Travelers in this report).

Timing of Organized Vaccination Campaigns

Persons and institutions planning substantial organized vaccination campaigns should consider scheduling these events after mid-October because the availability of vaccine in any location cannot be ensured consistently in early fall. Scheduling campaigns after mid-October will minimize the need for cancellations because vaccine is unavailable. Campaigns conducted before November using inactivated vaccine should focus efforts on vaccination of persons aged ≥50 years, persons aged <50 years at increased risk for influenza-related complications (including children aged 6–23 months and pregnant women), health-care workers, and household contacts of persons at high-risk (including children aged 0–23 months) to the extent feasible. Campaigns using the LAIV are also optimally conducted in October and November.

Vaccination in December and Later

After November, many persons who should or want to receive influenza vaccine remain unvaccinated. In addition, substantial amounts of vaccine are often left over at the end of the influenza season. To improve vaccine coverage, influenza vaccine should continue to be offered in December and throughout the influenza season as long as vaccine supplies are available, even after influenza activity has been documented in the community. In the United States, seasonal influenza activity can begin to increase as early as October or November, but influenza activity has not reached peak levels in the majority of recent seasons until late December–early March (Table 6). Therefore, although the timing of influenza activity can vary by region, vaccine administered after November

TABLE 6. Month of peak influenza activity* during 29 influenza seasons — United States, 1976–2005

	Month						
	Nov	Dec	Jan	Feb	Mar	Apr	May
No. (%) of years with peak influenza activity	1 (3)	4 (14)	6 (21)	13 (45)	3 (10)	1 (3)	1 (3)

* The peak week of activity was defined as the week with the greatest percentage of respiratory specimens testing positive for influenza on the basis of a 3-week moving average. Laboratory data were provided by U.S. World Health Organization Collaborating Centers (CDC, National Center for Infectious Diseases, unpublished data, 1976–2005).

is likely to be beneficial in the majority of influenza seasons. Adults develop peak antibody protection against influenza infection 2 weeks after vaccination (236,237).

Strategies for Implementing Vaccination Recommendations in Health-Care Settings

Successful vaccination programs combine publicity and education for health-care workers and other potential vaccine recipients, a plan for identifying persons at high risk, use of reminder/recall systems, assessment of practice-level vaccination rates with feedback to staff, and efforts to remove administrative and financial barriers that prevent persons from receiving the vaccine, including use of standing orders programs (19,238). Using standing orders programs is recommended for long-term-care facilities (e.g., nursing homes and skilled nursing facilities), hospitals, and home health agencies to ensure the administration of recommended vaccinations for adults (239). Standing orders programs for both influenza and pneumococcal vaccination should be conducted under the supervision of a licensed practitioner according to a physician-approved facility or agency policy by health-care workers trained to screen patients for contraindications to vaccination, administer vaccine, and monitor for adverse events. The Centers for Medicare and Medicaid Services (CMS) has removed the physician signature requirement for the administration of influenza and pneumococcal vaccines to Medicare and Medicaid patients in hospitals, long-term-care facilities, and home health agencies (239). To the extent allowed by local and state law, these facilities and agencies may implement standing orders for influenza and pneumococcal vaccination of Medicare- and Medicaid-eligible patients. Other settings (e.g., outpatient facilities, managed care organizations, assisted living facilities, correctional facilities, pharmacies, and adult workplaces) are encouraged to introduce standing orders programs as well (20). In addition, physician reminders (e.g., flagging charts) and patient reminders are recommended strategies for increasing rates of influenza vaccination. Persons for whom influenza vaccine is recommended can be identified and vaccinated in the settings described in the following sections.

Outpatient Facilities Providing Ongoing Care

Staff in facilities providing ongoing medical care (e.g., physicians' offices, public health clinics, employee health clinics, hemodialysis centers, hospital specialty-care clinics, and outpatient rehabilitation programs) should identify and label the medical records of patients who should receive vaccination. Vaccine should be offered during visits beginning in Septem-

ber and throughout the influenza season. The offer of vaccination and its receipt or refusal should be documented in the medical record. Patients for whom vaccination is recommended and who do not have regularly scheduled visits during the fall should be reminded by mail, telephone, or other means of the need for vaccination.

Outpatient Facilities Providing Episodic or Acute Care

Beginning each September, acute health-care facilities (e.g., emergency departments and walk-in clinics) should offer vaccinations to persons for whom vaccination is recommended or provide written information regarding why, where, and how to obtain the vaccine. This written information should be available in languages appropriate for the populations served by the facility.

Nursing Homes and Other Residential Long-Term-Care Facilities

During October and November each year, vaccination should be routinely provided to all residents of chronic-care facilities with the concurrence of attending physicians. Consent for vaccination should be obtained from the resident or a family member at the time of admission to the facility or anytime afterwards. All residents should be vaccinated at one time, preceding the influenza season. Residents admitted through March after completion of the vaccination program at the facility should be vaccinated at the time of admission.

Acute-Care Hospitals

Persons of all ages (including children) with high-risk conditions and persons aged ≥ 50 years who are hospitalized at any time during September–March should be offered and strongly encouraged to receive influenza vaccine before they are discharged. In one study, 39%–46% of adult patients hospitalized during the winter with influenza-related diagnoses had been hospitalized during the preceding autumn (240). Thus, the hospital serves as a setting in which persons at increased risk for subsequent hospitalization can be identified and vaccinated. However, vaccination of persons at high risk during or after their hospitalizations is often not done. In a study of hospitalized Medicare patients, only 31.6% were vaccinated before admission, 1.9% during admission, and 10.6% after admission (241). Using standing orders in hospitals increases vaccination rates among hospitalized persons (242).

Visiting Nurses and Others Providing Home Care to Persons at High Risk

Beginning in September, nursing-care plans should identify patients for whom vaccination is recommended, and vac-

cine should be administered in the home, if necessary. Caregivers and other persons in the household (including children) should be referred for vaccination.

Other Facilities Providing Services to Persons Aged ≥ 50 Years

Beginning in October, such facilities as assisted living housing, retirement communities, and recreation centers should offer unvaccinated residents and attendees vaccination on-site before the influenza season. Staff education should emphasize the need for influenza vaccine.

Health-Care Workers

Beginning in October each year, health-care facilities should offer influenza vaccinations to all workers, including night and weekend staff. Particular emphasis should be placed on providing vaccinations to persons who care for members of groups at high risk. Efforts should be made to educate health-care workers regarding the benefits of vaccination and the potential health consequences of influenza illness for themselves, their family members, and their patients. All health-care workers should be provided convenient access to influenza vaccine at the work site, free of charge, as part of employee health programs (127,137).

Influenza Vaccine Supply

Influenza vaccine distribution delays or vaccine supply shortages have occurred in the United States vaccine in three of the last five influenza seasons. Influenza vaccine delivery delays or vaccine shortages remain possible in part because of the inherent critical time constraints in manufacturing the vaccine given the annual updating of the influenza vaccine strains. Steps being taken to accommodate possible future delays or vaccine shortages include identification and implementation of ways to expand the influenza vaccine supply and improvement of targeted delivery of vaccine to groups at high risk when delays or shortages are expected.

Influenza Vaccine Use During Shortages of Inactivated Vaccine

ACIP will publish additional guidance regarding the prioritized (tiered) use of inactivated influenza vaccine to be implemented only during periods when there is a shortage of influenza vaccine. Otherwise, when vaccine is in adequate supply, every effort should be made to promote and use influenza vaccine for all regularly targeted groups and for other persons who wish to reduce their risk for influenza illness. The prioritized (tiered) use of influenza vaccine during inactivated influenza vaccine shortages applies only to use of inactivated vaccine and not to LAIV. When feasible, during

shortages of inactivated influenza vaccine, LAIV should be used preferentially for all healthy persons aged 5–49 years (including health-care workers) to increase the availability of inactivated vaccine for groups at high risk.

Future Directions for Influenza Vaccine Recommendations

ACIP plans to review new vaccination strategies for improving prevention and control of influenza, including the possibility of expanding recommendations for use of influenza vaccines. In addition, strategies for regularly monitoring vaccine effectiveness will be reviewed.

Recommendations for Using Antiviral Agents for Influenza

Antiviral drugs for influenza are an adjunct to influenza vaccine for controlling and preventing influenza. However, these agents are not a substitute for vaccination. Four licensed influenza antiviral agents are available in the United States: amantadine, rimantadine, zanamivir, and oseltamivir.

Amantadine and rimantadine are chemically related antiviral drugs known as adamantanes with activity against influenza A viruses, but not influenza B viruses. Amantadine was approved in 1966 for chemoprophylaxis of influenza A (H2N2) infection and was later approved in 1976 for treatment and chemoprophylaxis of influenza type A virus infections among adults and children aged ≥ 1 year. Rimantadine was approved in 1993 for treatment and chemoprophylaxis of influenza A infection among adults and prophylaxis among children. Although rimantadine is approved only for chemoprophylaxis of influenza A infection among children, rimantadine treatment for influenza A among children can be beneficial (243).

Zanamivir and oseltamivir are chemically related antiviral drugs known as neuraminidase inhibitors that have activity against both influenza A and B viruses. Both zanamivir and oseltamivir were approved in 1999 for treating uncomplicated influenza infections. Zanamivir is approved for treating persons aged ≥ 7 years, and oseltamivir is approved for treatment of persons aged ≥ 1 year. In 2000, oseltamivir was approved for chemoprophylaxis of influenza among persons aged ≥ 13 years.

The four drugs differ in pharmacokinetics, side effects, routes of administration, approved age groups, dosages, and costs. An overview of the indications, use, administration, and known primary side effects of these medications is presented in the following sections. Information contained in this report might not represent FDA approval or approved labeling for the anti-

viral agents described. Package inserts should be consulted for additional information.

Role of Laboratory Diagnosis

Appropriate treatment of patients with respiratory illness depends on accurate and timely diagnosis. Early diagnosis of influenza can reduce the inappropriate use of antibiotics and provide the option of using antiviral therapy. However, because certain bacterial infections can produce symptoms similar to influenza, bacterial infections should be considered and appropriately treated, if suspected. In addition, bacterial infections can occur as a complication of influenza.

Influenza surveillance information and diagnostic testing can aid clinical judgment and help guide treatment decisions. The accuracy of clinical diagnosis of influenza on the basis of symptoms alone is limited because symptoms from illness caused by other pathogens can overlap considerably with influenza (30,34,35). Influenza surveillance by state and local health departments and CDC can provide information regarding the presence of influenza viruses in the community. Surveillance can also identify the predominant circulating types, influenza A subtypes, and strains of influenza.

Diagnostic tests available for influenza include viral culture, serology, rapid antigen testing, polymerase chain reaction (PCR), and immunofluorescence assays (25). Sensitivity and specificity of any test for influenza might vary by the laboratory that performs the test, the type of test used, and the type of specimen tested. Among respiratory specimens for viral isolation or rapid detection, nasopharyngeal specimens are typically more effective than throat swab specimens (244). As with any diagnostic test, results should be evaluated in the context of other clinical and epidemiologic information available to health-care providers.

Commercial rapid diagnostic tests are available that can detect influenza viruses within 30 minutes (25,245). Some tests are approved for use in any outpatient setting, whereas others must be used in a moderately complex clinical laboratory. These rapid tests differ in the types of influenza viruses they can detect and whether they can distinguish between influenza types. Different tests can detect 1) only influenza A viruses; 2) both influenza A and B viruses, but not distinguish between the two types; or 3) both influenza A and B and distinguish between the two.

None of the tests provide any information about influenza A subtypes. The types of specimens acceptable for use (i.e., throat, nasopharyngeal, or nasal aspirates, swabs, or washes) also vary by test. The specificity and, in particular, the sensitivity of rapid tests are lower than for viral culture and vary by test (246,247). Because of the lower sensitivity of the rapid tests, physicians should consider confirming negative tests with

viral culture or other means because of the possibility of false-negative rapid test results, especially during periods of peak community influenza activity. In contrast, false-positive rapid test results are less likely, but can occur during periods of low influenza activity. Therefore, when interpreting results of a rapid influenza test, physicians should consider the positive and negative predictive values of the test in the context of the level of influenza activity in their community. Package inserts and the laboratory performing the test should be consulted for more details regarding use of rapid diagnostic tests. Additional information concerning diagnostic testing is available at <http://www.cdc.gov/flu/professionals/labdiagnosis.htm>.

Despite the availability of rapid diagnostic tests, collecting clinical specimens for viral culture is critical, because only culture isolates can provide specific information regarding circulating strains and subtypes of influenza viruses. This information is needed to compare current circulating influenza strains with vaccine strains, to guide decisions regarding influenza treatment and chemoprophylaxis, and to formulate vaccine for the coming year. Virus isolates also are needed to monitor the emergence of antiviral resistance and the emergence of novel influenza A subtypes that might pose a pandemic threat.

Indications for Use

Treatment

When administered within 2 days of illness onset to otherwise healthy adults, amantadine and rimantadine can reduce the duration of uncomplicated influenza A illness, and zanamivir and oseltamivir can reduce the duration of uncomplicated influenza A and B illness by approximately 1 day, compared with placebo (75,248–265). More clinical data are available concerning the efficacy of zanamivir and oseltamivir for treatment of influenza A infection than for treatment of influenza B infection (253,266–281). However, *in vitro* data and studies of treatment among mice and ferrets (282–289), in addition to clinical studies, have documented that zanamivir and oseltamivir have activity against influenza B viruses (254,258–260,290,291).

Data are limited regarding the effectiveness of the four antiviral agents in preventing serious influenza-related complications (e.g., bacterial or viral pneumonia or exacerbation of chronic diseases). Evidence for the effectiveness of these four antiviral drugs is principally based on studies of patients with uncomplicated influenza (292). Data are limited and inconclusive concerning the effectiveness of amantadine, rimantadine, zanamivir, and oseltamivir for treatment of influenza among persons at high risk for serious complications of influenza (28,248,250,251,253,254,261,266–270). One

study assessing oseltamivir treatment primarily among adults reported a reduction in complications, necessitating antibiotic therapy compared with placebo (271). Fewer studies of the efficacy of influenza antivirals have been conducted among pediatric populations (248,251,257,258,267,272,273). One study of oseltamivir treatment documented a decreased incidence of otitis media among children (258). Inadequate data exist regarding the safety and efficacy of any of the influenza antiviral drugs for use among children aged <1 year (247).

To reduce the emergence of antiviral drug-resistant viruses, amantadine or rimantadine therapy for persons with influenza A illness should be discontinued as soon as clinically warranted, typically after 3–5 days of treatment or within 24–48 hours after the disappearance of signs and symptoms. The recommended duration of treatment with either zanamivir or oseltamivir is 5 days.

Chemoprophylaxis

Chemoprophylactic drugs are not a substitute for vaccination, although they are critical adjuncts in preventing and controlling influenza. Both amantadine and rimantadine are indicated for chemoprophylaxis of influenza A infection, but not influenza B. Both drugs are approximately 60%–90% effective in preventing illness from influenza A infection (75,248,267). When used as prophylaxis, these antiviral agents can prevent illness while permitting subclinical infection and development of protective antibody against circulating influenza viruses. Therefore, certain persons who take these drugs will develop protective immune responses to circulating influenza viruses. Amantadine and rimantadine do not interfere with the antibody response to the vaccine (248). Both drugs have been studied extensively among nursing home populations as a component of influenza outbreak-control programs, which can limit the spread of influenza within chronic-care institutions (248,266,274–276).

Among the neuraminidase inhibitor antivirals zanamivir and oseltamivir, only oseltamivir has been approved for prophylaxis, but community studies of healthy adults indicate that both drugs are similarly effective in preventing febrile, laboratory-confirmed influenza illness (efficacy: zanamivir, 84%; oseltamivir, 82%) (253,277,293). Both antiviral agents have also been reported to prevent influenza illness among persons administered chemoprophylaxis after a household member had influenza diagnosed (278,290,293). Experience with prophylactic use of these agents in institutional settings or among patients with chronic medical conditions is limited in comparison with the adamantanes (260,269,270,279–281). One 6-week study of oseltamivir prophylaxis among nursing home residents reported a 92% reduction in influenza illness (260,294). Use of zanamivir has not been reported to impair

the immunologic response to influenza vaccine (259,295). Data are not available regarding the efficacy of any of the four antiviral agents in preventing influenza among severely immunocompromised persons.

When determining the timing and duration for administering influenza antiviral medications for prophylaxis, factors related to cost, compliance, and potential side effects should be considered. To be maximally effective as prophylaxis, the drug must be taken each day for the duration of influenza activity in the community. However, to be most cost-effective, one study of amantadine or rimantadine prophylaxis reported that the drugs should be taken only during the period of peak influenza activity in a community (296).

Persons at High Risk Who Are Vaccinated After Influenza Activity Has Begun. Persons at high risk for complications of influenza still can be vaccinated after an outbreak of influenza has begun in a community. However, development of antibodies in adults after vaccination takes approximately 2 weeks (236,237). When influenza vaccine is administered while influenza viruses are circulating, chemoprophylaxis should be considered for persons at high risk during the time from vaccination until immunity has developed. Children aged <9 years who receive influenza vaccine for the first time can require 6 weeks of prophylaxis (i.e., prophylaxis for 4 weeks after the first dose of vaccine and an additional 2 weeks of prophylaxis after the second dose).

Persons Who Provide Care to Those at High Risk. To reduce the spread of virus to persons at high risk during community or institutional outbreaks, chemoprophylaxis during peak influenza activity can be considered for unvaccinated persons who have frequent contact with persons at high risk. Persons with frequent contact include employees of hospitals, clinics, and chronic-care facilities, household members, visiting nurses, and volunteer workers. If an outbreak is caused by a variant strain of influenza that might not be controlled by the vaccine, chemoprophylaxis should be considered for all such persons, regardless of their vaccination status.

Persons Who Have Immune Deficiencies. Chemoprophylaxis can be considered for persons at high risk who are expected to have an inadequate antibody response to influenza vaccine. This category includes persons infected with HIV, chiefly those with advanced HIV disease. No published data are available concerning possible efficacy of chemoprophylaxis among persons with HIV infection or interactions with other drugs used to manage HIV infection. Such patients should be monitored closely if chemoprophylaxis is administered.

Other Persons. Chemoprophylaxis throughout the influenza season or during peak influenza activity might be appropriate for persons at high risk who should not be vaccinated.

Chemoprophylaxis can also be offered to persons who wish to avoid influenza illness. Health-care providers and patients should make this decision on an individual basis.

Control of Influenza Outbreaks in Institutions

Using antiviral drugs for treatment and prophylaxis of influenza is a key component of influenza outbreak control in institutions. In addition to antiviral medications, other outbreak-control measures include instituting droplet precautions and establishing cohorts of patients with confirmed or suspected influenza, re-offering influenza vaccinations to unvaccinated staff and patients, restricting staff movement between wards or buildings, and restricting contact between ill staff or visitors and patients (297–299) (for additional information regarding outbreak control in specific settings, see Additional Information Regarding Influenza Infection Control Among Specific Populations).

The majority of published reports concerning use of antiviral agents to control influenza outbreaks in institutions are based on studies of influenza A outbreaks among nursing home populations where amantadine or rimantadine were used (248,266,274–276,296). Less information is available concerning use of neuraminidase inhibitors in influenza A or B institutional outbreaks (269,270,281,294,300). When confirmed or suspected outbreaks of influenza occur in institutions that house persons at high risk, chemoprophylaxis should be started as early as possible to reduce the spread of the virus. In these situations, having preapproved orders from physicians or plans to obtain orders for antiviral medications on short notice can substantially expedite administration of antiviral medications.

When outbreaks occur in institutions, chemoprophylaxis should be administered to all residents, regardless of whether they received influenza vaccinations during the previous fall, and should continue for a minimum of 2 weeks. If surveillance indicates that new cases continue to occur, chemoprophylaxis should be continued until approximately 1 week after the end of the outbreak. The dosage for each resident should be determined individually. Chemoprophylaxis also can be offered to unvaccinated staff who provide care to persons at high risk. Prophylaxis should be considered for all employees, regardless of their vaccination status, if the outbreak is caused by a variant strain of influenza that is not well-matched by the vaccine.

In addition to nursing homes, chemoprophylaxis also can be considered for controlling influenza outbreaks in other closed or semiclosed settings (e.g., dormitories or other settings in which persons live in close proximity). For example, chemoprophylaxis with rimantadine has been used successfully to control an influenza A outbreak aboard a large cruise ship (177).

To limit the potential transmission of drug-resistant virus during outbreaks in institutions, whether in chronic or acute-care settings or other closed settings, measures should be taken to reduce contact as much as possible between persons taking antiviral drugs for treatment and other persons, including those taking chemoprophylaxis (see Antiviral Drug-Resistant Strains of Influenza).

Dosage

Dosage recommendations vary by age group and medical conditions (Table 7).

Children

Amantadine. Use of amantadine among children aged <1 year has not been adequately evaluated. The FDA-approved dosage for children aged 1–9 years for treatment and prophylaxis is 4.4–8.8 mg/kg body weight/day, not to exceed 150 mg/day. Although further studies are needed to determine the optimal dosage for children aged 1–9 years, physicians should consider prescribing only 5 mg/kg body weight/day (not to exceed 150 mg/day) to reduce the risk for toxicity. The approved dosage for children aged ≥10 years is 200 mg/day (100 mg twice a day); however, for children weighing <40 kg, prescribing 5 mg/kg body weight/day, regardless of age, is advisable (268).

Rimantadine. Rimantadine is approved for prophylaxis among children aged ≥1 year and for treatment and prophylaxis among adults. Although rimantadine is approved only for prophylaxis of infection among children, certain specialists in the management of influenza consider it appropriate for treatment among children (243). Use of rimantadine among children aged <1 year has not been adequately evaluated. Rimantadine should be administered in 1 or 2 divided doses at a dosage of 5 mg/kg body weight/day, not to exceed 150 mg/day for children aged 1–9 years. The approved dosage for children aged ≥10 years is 200 mg/day (100 mg twice a day); however, for children weighing <40 kg, prescribing 5 mg/kg body weight/day, regardless of age, is recommended (301).

Zanamivir. Zanamivir is approved for treatment among children aged ≥7 years. The recommended dosage of zanamivir for treatment of influenza is two inhalations (one 5-mg blister per inhalation for a total dose of 10 mg) twice daily (approximately 12 hours apart) (259).

Oseltamivir. Oseltamivir is approved for treatment among persons aged ≥1 year and for chemoprophylaxis among persons aged ≥13 years. Recommended treatment dosages for children vary by the weight of the child: the dosage recommendation for children who weigh ≤15 kg is 30 mg twice a day; for children weighing >15–23 kg, the dosage is 45 mg

TABLE 7. Recommended daily dosage of influenza antiviral medications for treatment and prophylaxis — United States

Antiviral agent	Age group (yrs)				
	1–6	7–9	10–12	13–64	≥65
Amantadine*					
Treatment, influenza A	5 mg/kg body weight/day up to 150 mg in 2 divided doses [†]	5 mg/kg body weight/day up to 150 mg in 2 divided doses [†]	100 mg twice daily [§]	100 mg twice daily [§]	≤100 mg/day
Prophylaxis, influenza A	5 mg/kg body weight/day up to 150 mg in 2 divided doses [†]	5 mg/kg body weight/day up to 150 mg in 2 divided doses [†]	100 mg twice daily [§]	100 mg twice daily [§]	≤100 mg/day
Rimantadine[¶]					
Treatment,** influenza A	NA ^{††}	NA	NA	100 mg twice daily ^{§ §§}	100 mg/day
Prophylaxis, influenza A	5 mg/kg body weight/day up to 150 mg in 2 divided doses [†]	5 mg/kg body weight/day up to 150 mg in 2 divided doses [†]	100 mg twice daily [§]	100 mg twice daily [§]	100 mg/day ^{¶¶}
Zanamivir*** †††					
Treatment, influenza A and B	NA	10 mg twice daily	10 mg twice daily	10 mg twice daily	10 mg twice daily
Oseltamivir					
Treatment, ^{§§§} influenza A and B	Dose varies by child's weight ^{¶¶¶}	Dose varies by child's weight ^{¶¶¶}	Dose varies by child's weight ^{¶¶¶}	75 mg twice daily	75 mg twice daily
Prophylaxis, influenza A and B	NA	NA	NA	75 mg/day	75 mg/day

NOTE: Amantadine manufacturers include Endo Pharmaceuticals (Symmetrel[®] — tablet and syrup); Geneva Pharms Tech (Amantadine HCL — capsule); USL Pharma (Amantadine HCL — capsule and tablet); and Alpharma, Carolina Medical, Copley Pharmaceutical, HiTech Pharma, Mikart, Morton Grove, and Pharmaceutical Associates (Amantadine HCL — syrup), and Sandoz. Rimantadine is manufactured by Forest Laboratories (Flumadine[®] — tablet and syrup); Corepharma, Impax Labs (Rimantadine HCL — tablet), and Amide Pharmaceuticals (Rimantadine HCL — tablet). Zanamivir is manufactured by GlaxoSmithKline (Relenza[®] — inhaled powder). Oseltamivir is manufactured by Roche Pharmaceuticals (Tamiflu[®] — tablet). This information is based on data published by the Food and Drug Administration (FDA), which is available at <http://www.fda.gov>.

* The drug package insert should be consulted for dosage recommendations for administering amantadine to persons with creatinine clearance ≤50 mL/min/1.73m².

[†] 5 mg/kg body weight of amantadine or rimantadine syrup = 1 tsp/2.2 lbs.

[§] Children aged ≥10 years who weigh <40 kg should be administered amantadine or rimantadine at a dosage of 5 mg/kg body weight/day.

[¶] A reduction in dosage to 100 mg/day of rimantadine is recommended for persons who have severe hepatic dysfunction or those with creatinine clearance ≤10 mL/min. Other persons with less severe hepatic or renal dysfunction taking 100 mg/day of rimantadine should be observed closely, and the dosage should be reduced or the drug discontinued, if necessary.

** Only approved by FDA for treatment among adults.

†† Not applicable.

^{§§} Rimantadine is approved by FDA for treatment among adults. However, certain specialists in the management of influenza consider rimantadine appropriate for treatment among children. Studies evaluating the efficacy of amantadine and rimantadine in children are limited, but they indicate that treatment with either drug diminishes the severity of influenza A infection when administered within 48 hours of illness onset (243).

^{¶¶} Older nursing-home residents should be administered only 100 mg/day of rimantadine. A reduction in dosage to 100 mg/day should be considered for all persons aged ≥65 years, if they experience possible side effects when taking 200 mg/day.

*** Zanamivir administered through inhalation by using a plastic device included in the medication package. Patients will benefit from instruction and demonstration of the correct use of the device.

††† Zanamivir is not approved for prophylaxis.

^{§§§} A reduction in the dose of oseltamivir is recommended for persons with creatinine clearance <30 mL/min.

^{¶¶¶} The dose recommendation for children who weigh ≤15 kg is 30 mg twice a day. For children who weigh >15–23 kg, the dose is 45 mg twice a day. For children who weigh >23–40 kg, the dose is 60 mg twice a day. And, for children who weigh >40 kg, the dose is 75 mg twice a day.

twice a day; for those weighing >23–40 kg, the dosage is 60 mg twice a day; and for children weighing >40 kg, the dosage is 75 mg twice a day. The treatment dosage for persons aged ≥13 years is 75 mg twice daily. For children aged ≥13 years, the recommended dose for prophylaxis is 75 mg once a day (260).

Persons Aged ≥65 Years

Amantadine. The daily dosage of amantadine for persons aged ≥65 years should not exceed 100 mg for prophylaxis or

treatment, because renal function declines with increasing age. For certain older persons, the dose should be further reduced.

Rimantadine. Among older persons, the incidence and severity of central nervous system (CNS) side effects are substantially lower among those taking rimantadine at a dosage of 100 mg/day than among those taking amantadine at dosages adjusted for estimated renal clearance (302). However, chronically ill older persons have had a higher incidence of

CNS and gastrointestinal symptoms and serum concentrations 2–4 times higher than among healthy, younger persons when rimantadine has been administered at a dosage of 200 mg/day (248).

For prophylaxis among persons aged ≥ 65 years, the recommended dosage is 100 mg/day. For treatment of older persons in the community, a reduction in dosage to 100 mg/day should be considered if they experience side effects when taking a dosage of 200 mg/day. For treatment of older nursing home residents, the dosage of rimantadine should be reduced to 100 mg/day (301).

Zanamivir and Oseltamivir. No reduction in dosage is recommended on the basis of age alone.

Persons with Impaired Renal Function

Amantadine. A reduction in dosage is recommended for patients with creatinine clearance ≤ 50 mL/min. Guidelines for amantadine dosage on the basis of creatinine clearance are located in the package insert. Because recommended dosages on the basis of creatinine clearance might provide only an approximation of the optimal dose for a given patient, such persons should be observed carefully for adverse reactions. If necessary, further reduction in the dose or discontinuation of the drug might be indicated because of side effects. Hemodialysis contributes minimally to amantadine clearance (303,304).

Rimantadine. A reduction in dosage to 100 mg/day is recommended for persons with creatinine clearance < 10 mL/min. Because of the potential for accumulation of rimantadine and its metabolites, patients with any degree of renal insufficiency, including older persons, should be monitored for adverse effects, and either the dosage should be reduced or the drug should be discontinued, if necessary. Hemodialysis contributes minimally to drug clearance (305).

Zanamivir. Limited data are available regarding the safety and efficacy of zanamivir for patients with impaired renal function. Among patients with renal failure who were administered a single intravenous dose of zanamivir, decreases in renal clearance, increases in half-life, and increased systemic exposure to zanamivir were observed (259,306). However, a limited number of healthy volunteers who were administered high doses of intravenous zanamivir tolerated systemic levels of zanamivir that were substantially higher than those resulting from administration of zanamivir by oral inhalation at the recommended dose (307,308). On the basis of these considerations, the manufacturer recommends no dose adjustment for inhaled zanamivir for a 5-day course of treatment for patients with either mild-to-moderate or severe impairment in renal function (259).

Oseltamivir. Serum concentrations of oseltamivir carboxylate (GS4071), the active metabolite of oseltamivir, increase with declining renal function (260,309). For patients with creatinine clearance of 10–30 mL/min (260), a reduction of the treatment dosage of oseltamivir to 75 mg once daily and in the prophylaxis dosage to 75 mg every other day is recommended. No treatment or prophylaxis dosing recommendations are available for patients undergoing routine renal dialysis treatment.

Persons with Liver Disease

Amantadine. No increase in adverse reactions to amantadine has been observed among persons with liver disease. Rare instances of reversible elevation of liver enzymes among patients receiving amantadine have been reported, although a specific relation between the drug and such changes has not been established (310).

Rimantadine. A reduction in dosage to 100 mg/day is recommended for persons with severe hepatic dysfunction.

Zanamivir and Oseltamivir. Neither of these medications has been studied among persons with hepatic dysfunction.

Persons with Seizure Disorders

Amantadine. An increased incidence of seizures has been reported among patients with a history of seizure disorders who have received amantadine (311). Patients with seizure disorders should be observed closely for possible increased seizure activity when taking amantadine.

Rimantadine. Seizures (or seizure-like activity) have been reported among persons with a history of seizures who were not receiving anticonvulsant medication while taking rimantadine (312). The extent to which rimantadine might increase the incidence of seizures among persons with seizure disorders has not been adequately evaluated.

Zanamivir and Oseltamivir. Seizure events have been reported during postmarketing use of zanamivir and oseltamivir, although no epidemiologic studies have reported any increased risk for seizures with either zanamivir or oseltamivir use.

Route

Amantadine, rimantadine, and oseltamivir are administered orally. Amantadine and rimantadine are available in tablet or syrup form, and oseltamivir is available in capsule or oral suspension form. Zanamivir is available as a dry powder that is self-administered via oral inhalation by using a plastic device included in the package with the medication. Patients will benefit from instruction and demonstration of correct use of this device.

Pharmacokinetics

Amantadine

Approximately 90% of amantadine is excreted unchanged in the urine by glomerular filtration and tubular secretion (274,313–316). Thus, renal clearance of amantadine is reduced substantially among persons with renal insufficiency, and dosages might need to be decreased (see Dosage) (Table 7).

Rimantadine

Approximately 75% of rimantadine is metabolized by the liver (267). The safety and pharmacokinetics of rimantadine among persons with liver disease have been evaluated only after single-dose administration (267,317). In a study of persons with chronic liver disease (the majority with stabilized cirrhosis), no alterations in liver function were observed after a single dose. However, for persons with severe liver dysfunction, the apparent clearance of rimantadine was 50% lower than that reported for persons without liver disease (302).

Rimantadine and its metabolites are excreted by the kidneys. The safety and pharmacokinetics of rimantadine among patients with renal insufficiency have been evaluated only after single-dose administration (267,305). Further studies are needed to determine multiple-dose pharmacokinetics and the most appropriate dosages for patients with renal insufficiency. In a single-dose study of patients with anuric renal failure, the apparent clearance of rimantadine was approximately 40% lower, and the elimination half-life was approximately 1.6-fold greater than that among healthy persons of the same age (305). Hemodialysis did not contribute to drug clearance. In studies of persons with less severe renal disease, drug clearance was also reduced, and plasma concentrations were higher than those among control patients without renal disease who were the same weight, age, and sex (301,317).

Zanamivir

In studies of healthy volunteers, approximately 7%–21% of the orally inhaled zanamivir dose reached the lungs, and 70%–87% was deposited in the oropharynx (259,318). Approximately 4%–17% of the total amount of orally inhaled zanamivir is systemically absorbed. Systemically absorbed zanamivir has a half-life of 2.5–5.1 hours and is excreted unchanged in the urine. Unabsorbed drug is excreted in the feces (259,308).

Oseltamivir

Approximately 80% of orally administered oseltamivir is absorbed systemically (309). Absorbed oseltamivir is metabolized to oseltamivir carboxylate, the active neuraminidase inhibitor, primarily by hepatic esterases. Oseltamivir carboxylate

has a half-life of 6–10 hours and is excreted in the urine by glomerular filtration and tubular secretion via the anionic pathway (260,319). Unmetabolized oseltamivir also is excreted in the urine by glomerular filtration and tubular secretion (320).

Side Effects and Adverse Reactions

When considering use of influenza antiviral medications (i.e., choice of antiviral drug, dosage, and duration of therapy), clinicians must consider the patient's age, weight, and renal function (Table 7); presence of other medical conditions; indications for use (i.e., prophylaxis or therapy); and the potential for interaction with other medications.

Amantadine and Rimantadine

Both amantadine and rimantadine can cause CNS and gastrointestinal side effects when administered to young, healthy adults at equivalent dosages of 200 mg/day. However, incidence of CNS side effects (e.g., nervousness, anxiety, insomnia, difficulty concentrating, and lightheadedness) is higher among persons taking amantadine than among those taking rimantadine (320). In a 6-week study of prophylaxis among healthy adults, approximately 6% of participants taking rimantadine at a dosage of 200 mg/day experienced one or more CNS symptoms, compared with approximately 13% of those taking the same dosage of amantadine and 4% of those taking placebo (320). A study of older persons also demonstrated fewer CNS side effects associated with rimantadine compared with amantadine (302). Gastrointestinal side effects (e.g., nausea and anorexia) occur among approximately 1%–3% of persons taking either drug, compared with 1% of persons receiving the placebo (320).

Side effects associated with amantadine and rimantadine are usually mild and cease soon after discontinuing the drug. Side effects can diminish or disappear after the first week, despite continued drug ingestion. However, serious side effects have been observed (e.g., marked behavioral changes, delirium, hallucinations, agitation, and seizures) (303,311). These more severe side effects have been associated with high plasma drug concentrations and have been observed most often among persons who have renal insufficiency, seizure disorders, or certain psychiatric disorders and among older persons who have been taking amantadine as prophylaxis at a dosage of 200 mg/day (274). Clinical observations and studies have indicated that lowering the dosage of amantadine among these persons reduces the incidence and severity of such side effects (Table 7). In acute overdosage of amantadine, CNS, renal, respiratory, and cardiac toxicity, including arrhythmias, have been reported (303). Because rimantadine has been marketed for a shorter period than amantadine, its

safety among certain patient populations (e.g., chronically ill and older persons) has been evaluated less frequently. Because amantadine has anticholinergic effects and might cause mydriasis, it should not be used among patients with untreated angle closure glaucoma (303).

Zanamivir

In a study of zanamivir treatment of ILI among persons with asthma or chronic obstructive pulmonary disease where study medication was administered after use of a B₂-agonist, 13% of patients receiving zanamivir and 14% of patients who received placebo (inhaled powdered lactose vehicle) experienced a >20% decline in forced expiratory volume in 1 second (FEV₁) after treatment (259,261). However, in a phase-I study of persons with mild or moderate asthma who did not have ILI, one of 13 patients experienced bronchospasm after administration of zanamivir (259). In addition, during postmarketing surveillance, cases of respiratory function deterioration after inhalation of zanamivir have been reported. Certain patients had underlying airway disease (e.g., asthma or chronic obstructive pulmonary disease). Because of the risk for serious adverse events and because the efficacy has not been demonstrated among this population, zanamivir is not recommended for treatment for patients with underlying airway disease (259). If physicians decide to prescribe zanamivir to patients with underlying chronic respiratory disease after carefully considering potential risks and benefits, the drug should be used with caution under conditions of appropriate monitoring and supportive care, including the availability of short-acting bronchodilators (292). Patients with asthma or chronic obstructive pulmonary disease who use zanamivir are advised to 1) have a fast-acting inhaled bronchodilator available when inhaling zanamivir and 2) stop using zanamivir and contact their physician if they experience difficulty breathing (259). No definitive evidence is available regarding the safety or efficacy of zanamivir for persons with underlying respiratory or cardiac disease or for persons with complications of acute influenza (292). Allergic reactions, including oropharyngeal or facial edema, have also been reported during postmarketing surveillance (259,269).

In clinical treatment studies of persons with uncomplicated influenza, the frequencies of adverse events were similar for persons receiving inhaled zanamivir and for those receiving placebo (i.e., inhaled lactose vehicle alone) (249–254,269). The most common adverse events reported by both groups were diarrhea; nausea; sinusitis; nasal signs and symptoms; bronchitis; cough; headache; dizziness; and ear, nose, and throat infections. Each of these symptoms was reported by <5% of persons in the clinical treatment studies combined (259).

Oseltamivir

Nausea and vomiting were reported more frequently among adults receiving oseltamivir for treatment (nausea without vomiting, approximately 10%; vomiting, approximately 9%) than among persons receiving placebo (nausea without vomiting, approximately 6%; vomiting, approximately 3%) (255,256,260,321). Among children treated with oseltamivir, 14.3% had vomiting, compared with 8.5% of placebo recipients. Overall, 1% discontinued the drug secondary to this side effect (258), whereas a limited number of adults who were enrolled in clinical treatment trials of oseltamivir discontinued treatment because of these symptoms (260). Similar types and rates of adverse events were reported in studies of oseltamivir prophylaxis (260). Nausea and vomiting might be less severe if oseltamivir is taken with food (260,321).

Use During Pregnancy

No clinical studies have been conducted regarding the safety or efficacy of amantadine, rimantadine, zanamivir, or oseltamivir for pregnant women; only two cases of amantadine use for severe influenza illness during the third trimester have been reported (144,145). However, both amantadine and rimantadine have been demonstrated in animal studies to be teratogenic and embryotoxic when administered at substantially high doses (301,303). Because of the unknown effects of influenza antiviral drugs on pregnant women and their fetuses, these four drugs should be used during pregnancy only if the potential benefit justifies the potential risk to the embryo or fetus (see manufacturers' package inserts) (259,260,301,303).

Drug Interactions

Careful observation is advised when amantadine is administered concurrently with drugs that affect CNS, including CNS stimulants. Concomitant administration of antihistamines or anticholinergic drugs can increase the incidence of adverse CNS reactions (248). No clinically substantial interactions between rimantadine and other drugs have been identified.

Clinical data are limited regarding drug interactions with zanamivir. However, no known drug interactions have been reported, and no clinically critical drug interactions have been predicted on the basis of *in vitro* data and data from studies using rats (259,322).

Limited clinical data are available regarding drug interactions with oseltamivir. Because oseltamivir and oseltamivir carboxylate are excreted in the urine by glomerular filtration and tubular secretion via the anionic pathway, a potential exists for interaction with other agents excreted by this pathway.

For example, coadministration of oseltamivir and probenecid resulted in reduced clearance of oseltamivir carboxylate by approximately 50% and a corresponding approximate two-fold increase in the plasma levels of oseltamivir carboxylate (260,319).

No published data are available concerning the safety or efficacy of using combinations of any of these four influenza antiviral drugs. For more detailed information concerning potential drug interactions for any of these influenza antiviral drugs, package inserts should be consulted.

Antiviral Drug-Resistant Strains of Influenza

Amantadine-resistant viruses are cross-resistant to rimantadine and vice versa (323). Drug-resistant viruses can appear in approximately one third of patients when either amantadine or rimantadine is used for therapy (273,323,324). During the course of amantadine or rimantadine therapy, resistant influenza strains can replace susceptible strains within 2–3 days of starting therapy (325,326). Resistant viruses have been isolated from persons who live at home or in an institution where other residents are taking or have recently taken amantadine or rimantadine as therapy (327,328); however, the frequency with which resistant viruses are transmitted and their effect on efforts to control influenza are unknown. Amantadine- and rimantadine-resistant viruses are not more virulent or transmissible than susceptible viruses (329). The screening of epidemic strains of influenza A has rarely detected amantadine- and rimantadine-resistant viruses (325,330,331).

Persons who have influenza A infection and who are treated with either amantadine or rimantadine can shed susceptible viruses early in the course of treatment and later shed drug-resistant viruses, including after 5–7 days of therapy (273). Such persons can benefit from therapy even when resistant viruses emerge.

Resistance to zanamivir and oseltamivir can be induced in influenza A and B viruses in vitro (332–339), but induction of resistance usually requires multiple passages in cell culture. By contrast, resistance to amantadine and rimantadine in vitro can be induced with fewer passages in cell culture (340,341). Development of viral resistance to zanamivir and oseltamivir during treatment has been identified but does not appear to be frequent (260,342–345). In one pediatric study, 5.5% of patients treated with oseltamivir had posttreatment isolates that were resistant to neuraminidase inhibitors. One limited study of Japanese children treated with oseltamivir reported a high frequency of resistant viruses (346). However, no transmission of neuraminidase inhibitor resistant viruses in humans has been documented to date. No isolates with reduced

susceptibility to zanamivir have been reported from clinical trials, although the number of posttreatment isolates tested is limited (347), and the risk for emergence of zanamivir-resistant isolates cannot be quantified (259). Only one clinical isolate with reduced susceptibility to zanamivir, obtained from an immunocompromised child on prolonged therapy, has been reported (343). Available diagnostic tests are not optimal for detecting clinical resistance to the neuraminidase inhibitor antiviral drugs, and additional tests are being developed (347,348). Postmarketing surveillance for neuraminidase inhibitor-resistant influenza viruses is being conducted (349).

Sources of Information Regarding Influenza and Its Surveillance

Information regarding influenza surveillance, prevention, detection, and control is available at <http://www.cdc.gov/flu/weekly/fluactivity.htm>. Surveillance information is available through the CDC Voice Information System (influenza update) at 888-232-3228 or CDC Fax Information Service at 888-232-3299. During October–May, surveillance information is updated at least every other week. In addition, periodic updates regarding influenza are published in the *MMWR Weekly Report* (<http://www.cdc.gov/mmwr>). Additional information regarding influenza vaccine can be obtained by calling 800-CDC-INFO (800-232-4636). State and local health departments should be consulted concerning availability of influenza vaccine, access to vaccination programs, information related to state or local influenza activity, and for reporting influenza outbreaks and receiving advice concerning outbreak control.

Additional Information Regarding Influenza Infection Control Among Specific Populations

Each year, ACIP provides general, annually updated information regarding control and prevention of influenza. Other reports related to controlling and preventing influenza among specific populations (e.g., immunocompromised persons, health-care workers, hospitals, and travelers) are also available in the following publications:

- CDC. Recommended adult immunization schedule—United States, 2004–05. *MMWR* 2004;53:Q1–4.
- Garner JS, Hospital Infection Control Practices Advisory Committee. Guideline for isolation precautions in hospitals. *Infect Control Hosp Epidemiol* 1996;17:53–80.

- CDC. Guidelines for preventing health-care-associated pneumonia, 2003: recommendations of CDC and the Health-care Infection Control Practices Advisory Committee. *MMWR* 2003;53(No. RR-3):1–36.
- Bolyard EA, Tablan OC, Williams WW, et al., Hospital Infection Control Practices Advisory Committee. Guideline for infection control in health-care personnel. *Am J Infect Control* 1998;26:289–354.
- CDC. Respiratory hygiene/cough etiquette in health-care settings. Atlanta, GA: US Department of Health and Human Services, CDC, 2003. Available at <http://www.cdc.gov/flu/professionals/infectioncontrol/resphygiene.htm>.
- Bradley SF. The Long-Term-Care Committee of the Society for Health-care Epidemiology of America. Prevention of influenza in long-term care facilities. *Infect Control Hosp Epidemiol* 1999;20:629–37.
- Sneller V-P, Izurieta H, Bridges C, et al. Prevention and control of vaccine-preventable diseases in long-term care facilities. *Journal of the American Medical Directors Association* 2000;1(Suppl):S2–37.
- American Academy of Pediatrics. 2003 red book: report of the Committee on Infectious Diseases. 26th ed. Elk Grove Village, IL: American Academy of Pediatrics; 2003.
- CDC. General recommendations on immunization: recommendations of the Advisory Committee on Immunization Practices (ACIP) and the American Academy of Family Practitioners (AAFP). *MMWR* 2002;51(No. RR-2):1–35.
- Bodnar UR, Maloney SA, Fielding KL, et al. Preliminary guidelines for the prevention and control of influenza-like illness among passengers and crew members on cruise ships. Atlanta, GA: US Department of Health and Human Services, CDC, National Center for Infectious Diseases; 1999.
- CDC. General recommendations for preventing influenza A infection among travelers. Atlanta, GA: US Department of Health and Human Services, CDC; 2003. Available at <http://www2.ncid.cdc.gov/travel/yb/utills/ybGet.asp?section=dis&obj=influenza.htm>.
- US Public Health Service (USPHS) and Infectious Diseases Society of America (IDSA). USPHS/IDSA Prevention of Opportunistic Infections Working Group. 2001 USPHS/IDSA guidelines for the prevention of opportunistic infections in persons infected with human immunodeficiency virus. Final November 28, 2001:1–65. Available at <http://www.aidsinfo.nih.gov>.
- CDC. Detection and control of influenza outbreaks in acute care facilities. Atlanta, GA: US Department of

Health and Human Services, CDC, National Center for Infectious Diseases; 2001. Available at <http://www.cdc.gov/ncidod/hip/INFECT/FluBook2001.pdf>.

References

1. Thompson WW, Shay DK, Weintraub E, et al. Mortality associated with influenza and respiratory syncytial virus in the United States. *JAMA* 2003;289:179–86.
2. Monto AS, Kiumeh F. The Tecumseh Study of Respiratory Illness. IX. Occurrence of influenza in the community, 1966–1971. *Am J Epidemiol* 1975;102:553–63.
3. Glezen WP, Couch RB. Interpandemic influenza in the Houston area, 1974–76. *N Engl J Med* 1978;298:587–92.
4. Glezen WP, Greenberg SB, Atmar RL, et al. Impact of respiratory virus infections on persons with chronic underlying conditions. *JAMA* 2000;283:499–505.
5. Barker WH. Excess pneumonia and influenza associated hospitalization during influenza epidemics in the United States, 1970–78. *Am J Public Health* 1986;76:761–5.
6. Barker WH, Mullooly JP. Impact of epidemic type A influenza in a defined adult population. *Am J Epidemiol* 1980;112:798–811.
7. Glezen WP. Serious morbidity and mortality associated with influenza epidemics. *Epidemiol Rev* 1982;4:25–44.
8. Office of Technology Assessment. Cost effectiveness of influenza vaccination. In: U.S. Congress. Washington, DC: Office of Technology Assessment; 1981.
9. Wilde JA, McMillan JA, Serwint J, et al. Effectiveness of influenza vaccine in health care professionals: a randomized trial. *JAMA* 1999;281:908–13.
10. Nichol KL, Lind A, Margolis KL, et al. The effectiveness of vaccination against influenza in healthy, working adults. *N Engl J Med* 1995;333:889–93.
11. Campbell DS, Rumley MH. Cost-effectiveness of the influenza vaccine in a healthy, working-age population. *J Occup Environ Med* 1997;39:408–14.
12. Bridges CB, Thompson WW, Meltzer MI, et al. Effectiveness and cost-benefit of influenza vaccination of healthy working adults: a randomized controlled trial. *JAMA* 2000;284:1655–63.
13. Patriarca PA, Weber JA, Parker RA, et al. Risk factors for outbreaks of influenza in nursing homes. A case-control study. *Am J Epidemiol* 1986;124:114–9.
14. Gross PA, Hermogenes AW, Sacks HS, et al. The efficacy of influenza vaccine in elderly persons. A meta-analysis and review of the literature. *Ann Intern Med* 1995;123:518–27.
15. Mullooly JP, Bennett MD, Hornbrook MC, et al. Influenza vaccination programs for elderly persons: cost-effectiveness in a health maintenance organization. *Ann Intern Med* 1994;121:947–52.
16. Clements DA, Langdon L, Bland C, et al. Influenza A vaccine decreases the incidence of otitis media in 6- to 30-month-old children in day care. *Arch Pediatr Adolesc Med* 1995;149:1113–7.
17. Heikkinen T, Ruuskanen O, Waris M, et al. Influenza vaccination in the prevention of acute otitis media in children. *Am J Dis Child* 1991;145:445–8.
18. Nordin J, Mullooly J, Poblete S, et al. Influenza vaccine effectiveness in preventing hospitalizations and deaths in persons 65 years or older in Minnesota, New York, and Oregon: data from 3 health plans. *J Infect Dis* 2001;184:665–70.

19. CDC. Vaccine-preventable diseases: improving vaccination coverage in children, adolescents, and adults: a report on recommendations from the Task Force on Community Preventive Services. *MMWR* 1999;48(No. RR-8).
20. CDC. Use of standing orders programs to increase adult vaccination rates. *MMWR* 2000;49(No. RR-1):15–6.
21. CDC. Improving influenza, pneumococcal polysaccharide, and hepatitis B vaccination coverage among adults aged <65 years at high risk: a report on recommendations of the Task Force on Community Preventive Services. *MMWR* 2005;54(No. RR-5).
22. Murphy BR WR. Orthomyxoviruses. In: Fields KD, Howley PM, eds. *Fields virology*. Philadelphia, PA: Lippincott; 1996:1397–445.
23. Clements ML, Betts RF, Tierney EL, et al. Serum and nasal wash antibodies associated with resistance to experimental challenge with influenza A wild-type virus. *J Clin Microbiol* 1986;24:157–60.
24. Couch RB, Kasel JA. Immunity to influenza in man. *Annu Rev Microbiol* 1983;37:529–49.
25. Cox NJ, Subbarao K. Influenza. *Lancet* 1999;354(9186):1277–82.
26. Frank AL, Taber LH, Wells CR, et al. Patterns of shedding of myxoviruses and paramyxoviruses in children. *J Infect Dis* 1981;144:433–41.
27. Klimov AI, Rocha E, Hayden FG, et al. Prolonged shedding of amantadine-resistant influenza A viruses by immunodeficient patients: detection by polymerase chain reaction-restriction analysis. *J Infect Dis* 1995;172:1352–5.
28. Englund JA, Champlin RE, Wyde PR, et al. Common emergence of amantadine- and rimantadine-resistant influenza A viruses in symptomatic immunocompromised adults. *Clin Infect Dis* 1998;26:1418–24.
29. Boivin G, Goyette N, Bernatchez H. Prolonged excretion of amantadine-resistant influenza a virus quasi species after cessation of antiviral therapy in an immunocompromised patient. *Clin Infect Dis* 2002;34:E23–5.
30. Nicholson KG. Clinical features of influenza. *Semin Respir Infect* 1992;7:26–37.
31. Ryan-Poirier K. Influenza virus infection in children. *Adv Pediatr Infect Dis* 1995;10:125–56.
32. Peltola V, Ziegler T, Ruuskanen O. Influenza A and B virus infections in children. *Clin Infect Dis* 2003;36:299–305.
33. Neuzil KM, Zhu Y, Griffin MR, et al. Burden of interpandemic influenza in children younger than 5 years: a 25-year prospective study. *J Infect Dis* 2002;185:147–52.
34. Boivin G, Hardy I, Tellier G, et al. Predicting influenza infections during epidemics with use of a clinical case definition. *Clin Infect Dis* 2000;31:1166–9.
35. Monto AS, Gravenstein S, Elliott M, et al. Clinical signs and symptoms predicting influenza infection. *Arch Intern Med* 2000;160:3243–7.
36. Orenstein WA, Bernier RH, Hinman AR. Assessing vaccine efficacy in the field. Further observations. *Epidemiol Rev* 1988;10:212–41.
37. Govaert TM, Dinant GJ, Aretz K, et al. The predictive value of influenza symptomatology in elderly people. *Fam Pract* 1998;15:16–22.
38. Walsh EE, Cox C, Falsey AR. Clinical features of influenza A virus infection in older hospitalized persons. *J Am Geriatr Soc* 2002;50:1498–503.
39. Neuzil KM, O'Connor TZ, Gorse GJ, et al. Recognizing influenza in older patients with chronic obstructive pulmonary disease who have received influenza vaccine. *Clin Infect Dis* 2003;36:169–74.
40. Douglas R Jr. Influenza in man. In: Kilbourne ED, ed. *Influenza viruses and influenza*. New York, NY: Academic Press, Inc.; 1975:395–418.
41. Dagan R, Hall CB. Influenza A virus infection imitating bacterial sepsis in early infancy. *Pediatr Infect Dis* 1984;3:218–21.
42. Meibalane R, Sedmak GV, Sasidharan P, et al. Outbreak of influenza in a neonatal intensive care unit. *J Pediatr* 1977;91:974–6.
43. Chiu SS, Tse CY, Lau YL, et al. Influenza A infection is an important cause of febrile seizures. *Pediatrics* 2001;108:E63.
44. McCullers JA, Facchini S, Chesney PJ, et al. Influenza B virus encephalitis. *Clin Infect Dis* 1999;28:898–900.
45. Morishima T, Togashi T, Yokota S, et al. Encephalitis and encephalopathy associated with an influenza epidemic in Japan. *Clin Infect Dis* 2002;35:512–7.
46. Thompson WW, Shay DK, Weintraub E, et al. Influenza-associated hospitalizations in the United States. *JAMA* 2004;292:1333–40.
47. Simonsen L, Schonberger LB, Stroup DF, et al. Impact of influenza on mortality in the USA. In: *Proceedings of the 3rd International Conference on Options for the Control of Influenza*, Cairns, Australia. Brown LE, Webster RG, eds. 1996, Amsterdam: Elsevier Science; 1996:26–32.
48. Lui KJ, Kendal AP. Impact of influenza epidemics on mortality in the United States from October 1972 to May 1985. *Am J Public Health* 1987;77:712–6.
49. Noble G. Epidemiological and clinical aspects of influenza. In: Beare AS, ed. *Basic and applied influenza research*. Boca Raton, FL: CRC Press; 1982:11–50.
50. Eickhoff TC, Sherman IL, Serfling RE. Observations on excess mortality associated with epidemic influenza. *JAMA* 1961;176:776–82.
51. Barker WH, Mullooly JP. Pneumonia and influenza deaths during epidemics: implications for prevention. *Arch Intern Med* 1982;142:85–9.
52. Simonsen L, Clarke MJ, Schonberger LB, et al. Pandemic versus epidemic influenza mortality: a pattern of changing age distribution. *J Infect Dis* 1998;178:53–60.
53. Glezen WP, Decker M, Perrotta DM. Survey of underlying conditions of persons hospitalized with acute respiratory disease during influenza epidemics in Houston, 1978–1981. *Am Rev Respir Dis* 1987;136:550–5.
54. Neuzil KM, Wright PF, Mitchel EF Jr, et al. The burden of influenza illness in children with asthma and other chronic medical conditions. *J Pediatr* 2000;137:856–64.
55. Izurieta HS, Thompson WW, Kramarz P, et al. Influenza and the rates of hospitalization for respiratory disease among infants and young children. *N Engl J Med* 2000;342:232–9.
56. Neuzil KM, Mellen BG, Wright PF, et al. The effect of influenza on hospitalizations, outpatient visits, and courses of antibiotics in children. *N Engl J Med* 2000;342:225–31.
57. Thompson WW, Shay DK, Weintraub E, et al. Influenza-associated hospitalizations in the United States. *JAMA* 2004;292:1333–40.
58. Simonsen L, Fukuda K, Schonberger LB, et al. The impact of influenza epidemics on hospitalizations. *J Infect Dis* 2000;181:831–7.
59. National Center for Health Statistics. *Health, United States, 1998*. Hyattsville, MD: US Department of Health and Human Services, CDC; 1998.
60. Simonsen L, Clarke MJ, Williamson GD, et al. The impact of influenza epidemics on mortality: introducing a severity index. *Am J Public Health* 1997;87:1944–50.
61. CDC. Update: influenza-associated deaths reported among children aged <18 years—United States, 2003–04 influenza season. *MMWR* 2004;52:1286–8.

62. CDC. Severe morbidity and mortality associated with influenza in children and young adults—Michigan, 2003. *MMWR* 2003;52:837–40.
63. Kilbourne E. *Influenza*. New York, NY: Plenum Medical Book Company; 1987.
64. CDC. Recommendations regarding the use of vaccines that contain thimerosal as a preservative. *MMWR* 1999;48:996–8.
65. Pichichero ME, Cernichiari E, Lopreiato J, et al. Mercury concentrations and metabolism in infants receiving vaccines containing thiomersal: a descriptive study. *Lancet* 2002;360(9347):1737–41.
66. Stratton K, Gable A, McCormick MC, eds. *Immunization safety review: thimerosal-containing vaccines and neurodevelopmental disorders*. Washington, DC: National Academy Press; 2001.
67. CDC. General recommendations on immunization: recommendations of the Advisory Committee on Immunization Practices (ACIP) and the American Academy of Family Physicians (AAFP). *MMWR* 2002;51(No. RR-2):1–35.
68. CDC. Summary of the joint statement on thimerosal in vaccines. American Academy of Family Physicians, American Academy of Pediatrics, Advisory Committee on Immunization Practices, Public Health Service. *MMWR* 2000;49:622, 631.
69. La Montagne JR, Noble GR, Quinnan GV, et al. Summary of clinical trials of inactivated influenza vaccine—1978. *Rev Infect Dis* 1983;5:723–36.
70. Oxford JS, Schild GC, Potter CW, et al. The specificity of the anti-haemagglutinin antibody response induced in man by inactivated influenza vaccines and by natural infection. *J Hyg (Lond)* 1979;82:51–61.
71. Neuzil KM, Dupont WD, Wright PF, et al. Efficacy of inactivated and cold-adapted vaccines against influenza A infection, 1985 to 1990: the pediatric experience. *Pediatr Infect Dis J* 2001;20:733–40.
72. Potter CW, Oxford JS. Determinants of immunity to influenza infection in man. *Br Med Bull* 1979;35:69–75.
73. Hirota Y, Kaji M, Ide S, et al. Antibody efficacy as a keen index to evaluate influenza vaccine effectiveness. *Vaccine* 1997;15:962–7.
74. Palache AM. Influenza vaccines. A reappraisal of their use. *Drugs* 1997;54:841–56.
75. Demicheli V, Jefferson T, Rivetti D, et al. Prevention and early treatment of influenza in healthy adults. *Vaccine* 2000;18:957–1030.
76. Smith JW, Pollard R. Vaccination against influenza: a five-year study in the Post Office. *J Hyg (Lond)* 1979;83:157–70.
77. CDC. Assessment of the effectiveness of the 2003–04 influenza vaccine among children and adults—Colorado, 2003. *MMWR* 2004;53:707–10.
78. Gonzalez M, Pirez MC, Ward E, et al. Safety and immunogenicity of a paediatric presentation of an influenza vaccine. *Arch Dis Child* 2000;83:488–91.
79. Wright PF, Cherry JD, Foy HM, et al. Antigenicity and reactogenicity of influenza A/USSR/77 virus vaccine in children—a multicentered evaluation of dosage and safety. *Rev Infect Dis* 1983;5:758–64.
80. Daubeney P, Taylor CJ, McGaw J, et al. Immunogenicity and tolerability of a trivalent influenza subunit vaccine (Influvac) in high-risk children aged 6 months to 4 years. *Br J Clin Pract* 1997;51:87–90.
81. Wright PF, Thompson J, Vaughn WK, et al. Trials of influenza A/New Jersey/76 virus vaccine in normal children: an overview of age-related antigenicity and reactogenicity. *J Infect Dis* 1977;136(Suppl):S731–41.
82. Groothuis JR, Lehr MV, Levin MJ. Safety and immunogenicity of a purified haemagglutinin antigen in very young high-risk children. *Vaccine* 1994;12:139–41.
83. Park CL, Frank AL, Sullivan M, et al. Influenza vaccination of children during acute asthma exacerbation and concurrent prednisone therapy. *Pediatrics* 1996;98(2 Pt 1):196–200.
84. Clover RD, Crawford S, Glezen WP, et al. Comparison of heterotypic protection against influenza A/Taiwan/86 (H1N1) by attenuated and inactivated vaccines to A/Chile/83-like viruses. *J Infect Dis* 1991;163:300–4.
85. Sugaya N, Nerome K, Ishida M, et al. Efficacy of inactivated vaccine in preventing antigenically drifted influenza type A and well-matched type B. *JAMA* 1994;272:1122–6.
86. Hoberman A, Greenberg DP, Paradise JL, et al. Effectiveness of inactivated influenza vaccine in preventing acute otitis media in young children: a randomized controlled trial. *JAMA* 2003;290:1608–16.
87. Ritzwoller DP, Brides CB, Sheetterly S, et al. Effectiveness of the 2003–04 influenza vaccine among children 6 months to 8 years for 1 versus 2 doses. *Pediatrics* 2005; In press.
88. Blumberg EA, Albano C, Pruett T, et al. The immunogenicity of influenza virus vaccine in solid organ transplant recipients. *Clin Infect Dis* 1996;22:295–302.
89. Dorrell L, Hassan I, Marshall S, et al. Clinical and serological responses to an inactivated influenza vaccine in adults with HIV infection, diabetes, obstructive airways disease, elderly adults and healthy volunteers. *Int J STD AIDS* 1997;8:776–9.
90. McElhane JE, Beattie BL, Devine R, et al. Age-related decline in interleukin 2 production in response to influenza vaccine. *J Am Geriatr Soc* 1990;38:652–8.
91. Govaert TM, Thijs CT, Masarel N, et al. The efficacy of influenza vaccination in elderly individuals. A randomized double-blind placebo-controlled trial. *JAMA* 1994;272:1661–5.
92. Hak E, Nordin J, Wei F, et al. Influence of high-risk medical conditions on the effectiveness of influenza vaccination among elderly members of 3 large managed-care organizations. *Clin Infect Dis* 2002;35:370–7.
93. Nichol KL, Wuorenma J, von Sternberg T. Benefits of influenza vaccination for low-, intermediate-, and high-risk senior citizens. *Arch Intern Med* 1998;158:1769–76.
94. Patriarca PA, Weber JA, Parker RA, et al. Efficacy of influenza vaccine in nursing homes. Reduction in illness and complications during an influenza A (H3N2) epidemic. *JAMA* 1985;253:1136–9.
95. Arden NH, PA Patriarcha, Kendal AP. Experiences in the use and efficacy of inactivated influenza vaccine in nursing homes. In: *Options for the Control of Influenza*. New York, NY: Alan R. Liss, Inc.; 1986.
96. Monto AS, Hornbuckle K, Ohmit SE. Influenza vaccine effectiveness among elderly nursing home residents: a cohort study. *Am J Epidemiol* 2001;154:155–60.
97. Belshe RB, Mendelman PM, Treanor J, et al. The efficacy of live attenuated, cold-adapted, trivalent, intranasal influenza virus vaccine in children. *N Engl J Med* 1998;338:1405–12.
98. Belshe RB, Gruber WC, Mendelman PM, et al. Efficacy of vaccination with live attenuated, cold-adapted, trivalent, intranasal influenza virus vaccine against a variant (A/Sydney) not contained in the vaccine. *J Pediatr* 2000;136:168–75.
99. Halloran ME, Longini IM Jr, Gaglani MJ, et al. Estimating efficacy of trivalent, cold-adapted, influenza virus vaccine (CAIV-T) against influenza A (H1N1) and B using surveillance cultures. *Am J Epidemiol* 2003;158:305–11.

100. Nichol KL, Mendelman PM, Mallon KP, et al. Effectiveness of live, attenuated intranasal influenza virus vaccine in healthy, working adults: a randomized controlled trial. *JAMA* 1999;282:137–44.
101. Treanor JJ, Kotloff K, Betts RF, et al. Evaluation of trivalent, live, cold-adapted (CAIV-T) and inactivated (TIV) influenza vaccines in prevention of virus infection and illness following challenge of adults with wild-type influenza A (H1N1), A (H3N2), and B viruses. *Vaccine* 1999;18:899–906.
102. Riddiough MA, Sisk JE, Bell JC. Influenza vaccination. *JAMA* 1983;249:3189–95.
103. Mixeu MA, Vespa GN, Forleo-Neto E, et al. Impact of influenza vaccination on civilian aircrew illness and absenteeism. *Aviat Space Environ Med* 2002;73:876–80.
104. Nichol KL, Mallon KP, Mendelman PM. Cost benefit of influenza vaccination in healthy, working adults: an economic analysis based on the results of a clinical trial of trivalent live attenuated influenza virus vaccine. *Vaccine* 2003;21:2207–17.
105. Nichol KL, Mendelman P. Influence of clinical case definitions with differing levels of sensitivity and specificity on estimates of the relative and absolute health benefits of influenza vaccination among healthy working adults and implications for economic analyses. *Virus Res* 2004;103:3–8.
106. Nichol KL. Cost-benefit analysis of a strategy to vaccinate healthy working adults against influenza. *Arch Intern Med* 2001;161:749–59.
107. Cohen GM, Nettleman MD. Economic impact of influenza vaccination in preschool children. *Pediatrics* 2000;106:973–6.
108. White T, Lavoie S, Nettleman MD. Potential cost savings attributable to influenza vaccination of school-aged children. *Pediatrics* 1999;103:e73.
109. Dayan GH, Nguyen VH, Debbag R, et al. Cost-effectiveness of influenza vaccination in high-risk children in Argentina. *Vaccine* 2001;19:4204–13.
110. Luce BR, Zangwill KM, Palmer CS, et al. Cost-effectiveness analysis of an intranasal influenza vaccine for the prevention of influenza in healthy children. *Pediatrics* 2001;108:E24.
111. US Department of Health and Human Services. *Healthy People 2010* (conference ed., in 2 vols). Washington, DC: US Department of Health and Human Services; 2000.
112. CDC. Influenza and pneumococcal vaccination coverage levels among persons aged ≥ 65 years—United States, 1973–1993. *MMWR* 1995;44:506–7, 513–5.
113. CDC. National Health Interview Survey—2002. Hyattsville, MD: US Department of Health and Human Services, CDC, National Center for Health Statistics; 2002. Available at http://www.cdc.gov/nchs/about/major/nhis/quest_data_related_1997_forward.htm.
114. US Department of Health and Human Services. *Healthy People 2000: national health promotion and disease prevention objectives—full report, with commentary*. Washington, DC: US Department of Health and Human Services, Public Health Service; 1991.
115. CDC. Implementation of the Medicare influenza vaccination benefit—United States, 1993. *MMWR* 1994;43:771–3.
116. Singleton JA, Greby SM, Wooten KG, et al. Influenza, pneumococcal, and tetanus toxoid vaccination of adults—United States, 1993–1997. In: *Surveillance Summaries*, September 22, 2000. *MMWR* 2000;49(No. SS-9):39–62.
117. CDC. Racial/ethnic disparities in influenza and pneumococcal vaccination levels among persons aged ≥ 65 Years—United States, 1989–2001. *MMWR* 2003;52:958–62.
118. Buikema AR SJ, Sneller VP, Strikas RA. Influenza vaccination in nursing homes, United States, 1995 and 1997 [Abstract P2-49]. Presented at the Options for the Control of Influenza IV Conference, Crete, Greece; September 23–28, 2000.
119. Zadeh MM, Buxton Bridges C, Thompson WW, et al. Influenza outbreak detection and control measures in nursing homes in the United States. *J Am Geriatr Soc* 2000;48:1310–5.
120. Kramarz P, DeStefano F, Gargiullo PM, et al. Influenza vaccination in children with asthma in health maintenance organizations. Vaccine Safety Datalink Team. *Vaccine* 2000;18:2288–94.
121. Chung EK, Casey R, Pinto-Martin JA, et al. Routine and influenza vaccination rates in children with asthma. *Ann Allergy Asthma Immunol* 1998;80:318–22.
122. Gaglani M, Riggs M, Kamenicky C, et al. A computerized reminder strategy is effective for annual influenza immunization of children with asthma or reactive airway disease. *Pediatr Infect Dis J* 2001;20:1155–60.
123. Marshall BC, Henshaw C, Evans DA, et al. Influenza vaccination coverage level at a cystic fibrosis center. *Pediatrics* 2002;109:E80–0.
124. CDC. Estimated influenza vaccination coverage among adults and children—United States, September 1, 2004–January 31, 2005. *MMWR* 2005;54:304–7.
125. Potter J, Stott DJ, Roberts MA, et al. Influenza vaccination of health care workers in long-term-care hospitals reduces the mortality of elderly patients. *J Infect Dis* 1997;175:1–6.
126. Carman WF, Elder AG, Wallace LA, et al. Effects of influenza vaccination of health-care personnel on mortality of elderly people in long-term care: a randomised controlled trial. *Lancet* 2000;355(9198):93–7.
127. National Foundation for Infectious Diseases. *Call to action: influenza immunization among health-care personnel, 2003*. Bethesda, MD: National Foundation for Infectious Diseases; 2003. Available at <http://www.nfid.org>.
128. Poland GA, Tosh P, Jacobson RM. Requiring influenza vaccination for health care workers: seven truths we must accept. *Vaccine* 2005;23:2251–5.
129. Lu P, Singleton J. Influenza vaccination of pregnant women: Behavioral Risk Factor Surveillance System (BRFSS), 1997–2001. Presented at the Annual Behavioral Risk Factor Surveillance System Conference, St. Louis, Missouri; 2003.
130. Yeager DP, Toy EC, Baker B III. Influenza vaccination in pregnancy. *Am J Perinatol* 1999;16:283–6.
131. Gonik B, Jones T, Contreras D, et al. The obstetrician-gynecologist's role in vaccine-preventable diseases and immunization. *Obstet Gynecol* 2000;96:81–4.
132. Zimmerman RK, Raymund M, Janosky JE, et al. Sensitivity and specificity of patient self-report of influenza and pneumococcal polysaccharide vaccinations among elderly outpatients in diverse patient care strata. *Vaccine* 2003;21:1486–91.
133. O'Mara D, Fukuda K, Singleton JA. Influenza vaccine: ensuring timely and adequate supply. *Infect Med* 2003;20:548–54.
134. CDC. Assessing adult vaccination status at age 50 years. *MMWR* 1995;44:561–3.
135. Fedson DS. Adult immunization. Summary of the National Vaccine Advisory Committee Report. *JAMA* 1994;272:1133–7.
136. Piedra PA, Gaglani MJ, Kozinetz CA, et al. Herd immunity in adults against influenza-related illnesses with use of the trivalent-live attenuated influenza vaccine (CAIV-T) in children. *Vaccine* 2005;23:1540–8.

137. CDC. Interventions to increase influenza vaccination of health-care personnel—California and Minnesota. *MMWR* 2005;54:196–9.
138. Stewart A, Cox M, Rosenbaum S. The epidemiology of U.S. immunization law: immunization requirements for staff and residents of long-term care facilities under state laws/regulations. Available at <http://www.gwumc.edu/sphhs/healthpolicy/immunization/EUSIL-LTC-report.pdf>.
139. Harris JW. Influenza occurring pregnant women: a statistical study of thirteen hundred and fifty cases. *JAMA* 1919;72:978–80.
140. Widelock D, Csizmas L, Klein S. Influenza, pregnancy, and fetal outcome. *Public Health Rep* 1963;78:1–11.
141. Freeman DW, Barno A. Deaths from Asian influenza associated with pregnancy. *Am J Obstet Gynecol* 1959;78:1172–5.
142. Shahab SZ, Glezen WP. Influenza virus. In: Gonik B, ed. *Viral diseases in pregnancy*. New York, NY: Springer-Verlag; 1994:215–23.
143. Schoenbaum SC, Weinstein L. Respiratory infection in pregnancy. *Clin Obstet Gynecol* 1979;22:293–300.
144. Kirshon B, Faro S, Zurawin RK, et al. Favorable outcome after treatment with amantadine and ribavirin in a pregnancy complicated by influenza pneumonia. A case report. *J Reprod Med* 1988;33:399–401.
145. Kort BA, Cefalo RC, Baker VV. Fatal influenza A pneumonia in pregnancy. *Am J Perinatol* 1986;3:179–82.
146. Irving WL, James DK, Stephenson T, et al. Influenza virus infection in the second and third trimesters of pregnancy: a clinical and seroepidemiological study. *BJOG* 2000;107:1282–9.
147. Neuzil KM, Reed GW, Mitchel EF, et al. Impact of influenza on acute cardiopulmonary hospitalizations in pregnant women. *Am J Epidemiol* 1998;148:1094–102.
148. Heinonen OP, Shapiro S, Monson RR, et al. Immunization during pregnancy against poliomyelitis and influenza in relation to childhood malignancy. *Int J Epidemiol* 1973;2:229–35.
149. Mullooly JP, Barker WH. Impact of type A influenza on children: a retrospective study. *Am J Public Health* 1982;72:1008–16.
150. Glezen WP, Decker M, Joseph SW, et al. Acute respiratory disease associated with influenza epidemics in Houston, 1981–1983. *J Infect Dis* 1987;155:1119–26.
151. Cooney MK, Fox JP, Hall CE. The Seattle Virus Watch. VI. Observations of infections with and illness due to parainfluenza, mumps and respiratory syncytial viruses and *Mycoplasma pneumoniae*. *Am J Epidemiol* 1975;101:532–51.
152. Glezen WP, Taber LH, Frank AL, et al. Risk of primary infection and reinfection with respiratory syncytial virus. *Am J Dis Child* 1986;140:543–6.
153. Glezen WP. Morbidity associated with the major respiratory viruses. *Pediatr Ann* 1990;19:535–6, 538, 540, passim.
154. CDC. Prevention and control of influenza: recommendations of the Advisory Committee on Immunization Practices (ACIP). *MMWR* 2004;53(No. RR-6):1–40.
155. CDC. Expansion of availability for influenza vaccines through the Vaccines for Children Program. *MMWR* 2002;51:864, 875.
156. Couch RB. Influenza, influenza virus vaccine, and human immunodeficiency virus infection. *Clin Infect Dis* 1999;28:548–51.
157. Tasker SA, O'Brien WA, Treanor JJ, et al. Effects of influenza vaccination in HIV-infected adults: a double-blind, placebo-controlled trial. *Vaccine* 1998;16:1039–42.
158. Neuzil KM, Reed GW, Mitchel EF Jr., et al. Influenza-associated morbidity and mortality in young and middle-aged women. *JAMA* 1999;281:901–7.
159. Lin JC, Nichol KL. Excess mortality due to pneumonia or influenza during influenza seasons among persons with acquired immunodeficiency syndrome. *Arch Intern Med* 2001;161:441–6.
160. Safrin S, Rush JD, Mills J. Influenza in patients with human immunodeficiency virus infection. *Chest* 1990;98:33–7.
161. Radwan HM, Cheeseman SH, Lai KK, et al. Influenza in human immunodeficiency virus-infected patients during the 1997–1998 influenza season. *Clin Infect Dis* 2000;31:604–6.
162. Fine AD, Bridges CB, De Guzman AM, et al. Influenza A among patients with human immunodeficiency virus: an outbreak of infection at a residential facility in New York City. *Clin Infect Dis* 2001;32:1784–91.
163. Chadwick EG, Chang G, Decker MD, et al. Serologic response to standard inactivated influenza vaccine in human immunodeficiency virus-infected children. *Pediatr Infect Dis J* 1994;13:206–11.
164. Huang KL, Ruben FL, Rinaldo CR Jr., et al. Antibody responses after influenza and pneumococcal immunization in HIV-infected homosexual men. *JAMA* 1987;257:2047–50.
165. Staprans SI, Hamilton BL, Follansbee SE, et al. Activation of virus replication after vaccination of HIV-1-infected individuals. *J Exp Med* 1995;182:1727–37.
166. Kroon FP, van Dissel JT, de Jong JC, et al. Antibody response after influenza vaccination in HIV-infected individuals: a consecutive 3-year study. *Vaccine* 2000;18:3040–9.
167. Miotti PG, Nelson KE, Dallabetta GA, et al. The influence of HIV infection on antibody responses to a two-dose regimen of influenza vaccine. *JAMA* 1989;262:779–83.
168. Ho DD. HIV-1 viraemia and influenza. *Lancet* 1992;339(8808):1549.
169. O'Brien WA, Grovit-Ferbas K, Namazi A, et al. Human immunodeficiency virus-type 1 replication can be increased in peripheral blood of seropositive patients after influenza vaccination. *Blood* 1995;86:1082–9.
170. Glesby MJ, Hoover DR, Farzadegan H, et al. The effect of influenza vaccination on human immunodeficiency virus type 1 load: a randomized, double-blind, placebo-controlled study. *J Infect Dis* 1996;174:1332–6.
171. Fowke KR, D'Amico R, Chernoff DN, et al. Immunologic and virologic evaluation after influenza vaccination of HIV-1-infected patients. *AIDS* 1997;11:1013–21.
172. Fuller JD, Craven DE, Steger KA, et al. Influenza vaccination of human immunodeficiency virus (HIV)-infected adults: impact on plasma levels of HIV type 1 RNA and determinants of antibody response. *Clin Infect Dis* 1999;28:541–7.
173. Amendola A, Boschini A, Colzani D, et al. Influenza vaccination of HIV-1-positive and HIV-1-negative former intravenous drug users. *J Med Virol* 2001;65:644–8.
174. Sullivan PS, Hanson DL, Dworkin MS, et al. Effect of influenza vaccination on disease progression among HIV-infected persons. *AIDS* 2000;14:2781–5.
175. Gunthard HF, Wong JK, Spina CA, et al. Effect of influenza vaccination on viral replication and immune response in persons infected with human immunodeficiency virus receiving potent antiretroviral therapy. *J Infect Dis* 2000;181:522–31.
176. Miller JM, Tam TW, Maloney S, et al. Cruise ships: high-risk passengers and the global spread of new influenza viruses. *Clin Infect Dis* 2000;31:433–8.

177. Uyeki TM, Zane SB, Bodnar UR, et al. Large summertime influenza A outbreak among tourists in Alaska and the Yukon Territory. *Clin Infect Dis* 2003;36:1095–102.
178. Gross PA, Weksler ME, Quinnan GV Jr., et al. Immunization of elderly people with two doses of influenza vaccine. *J Clin Microbiol* 1987;25:1763–5.
179. Feery BJ, Cheyne IM, Hampson AW, et al. Antibody response to one and two doses of influenza virus subunit vaccine. *Med J Aust* 1976;1:186, 188–9.
180. Levine M, Beattie BL, McLean DM. Comparison of one- and two-dose regimens of influenza vaccine for elderly men. *CMAJ* 1987;137:722–6.
181. Cate TR, Couch RB, Parker D, et al. Reactogenicity, immunogenicity, and antibody persistence in adults given inactivated influenza virus vaccines—1978. *Rev Infect Dis* 1983;5:737–47.
182. Kunzel W, Glathe H, Engelmann H, et al. Kinetics of humoral antibody response to trivalent inactivated split influenza vaccine in subjects previously vaccinated or vaccinated for the first time. *Vaccine* 1996;14:1108–10.
183. Poland GA, Borrud A, Jacobson RM, et al. Determination of deltoid fat pad thickness. Implications for needle length in adult immunization. *JAMA* 1997;277:1709–11.
184. Govaert TM, Dinant GJ, Aretz K, et al. Adverse reactions to influenza vaccine in elderly people: randomised double blind placebo controlled trial. *BMJ* 1993;307:988–90.
185. Margolis KL, Nichol KL, Poland GA, et al. Frequency of adverse reactions to influenza vaccine in the elderly. A randomized, placebo-controlled trial. *JAMA* 1990;264:1139–41.
186. Nichol KL, Margolis KL, Lind A, et al. Side effects associated with influenza vaccination in healthy working adults. A randomized, placebo-controlled trial. *Arch Intern Med* 1996;156:1546–50.
187. American Lung Association Asthma Clinical Research Centers. The safety of inactivated influenza vaccine in adults and children with asthma. *N Engl J Med* 2001;345:1529–36.
188. Piedra PA, Glezen WP, Mbawuikie I, et al. Studies on reactogenicity and immunogenicity of attenuated bivalent cold recombinant influenza type A (CRA) and inactivated trivalent influenza virus (TI) vaccines in infants and young children. *Vaccine* 1993;11:718–24.
189. Scheifele DW, Bjornson G, Johnston J. Evaluation of adverse events after influenza vaccination in hospital personnel. *CMAJ* 1990;142:127–30.
190. Barry DW, Mayner RE, Hochstein HD, et al. Comparative trial of influenza vaccines. II. Adverse reactions in children and adults. *Am J Epidemiol* 1976;104:47–59.
191. France EK, Jackson L, Vaccine Safety Datalink Team. Safety of the trivalent inactivated influenza vaccine among children: a population-based study [Abstract 76]. Presented at the National Immunization Conference, Chicago, Illinois; 2003.
192. Groothuis JR, Levin MJ, Rabalais GP, et al. Immunization of high-risk infants younger than 18 months of age with split-product influenza vaccine. *Pediatrics* 1991;87:823–8.
193. McMahon AW, Iskander J, Haber P, et al. Adverse events after inactivated influenza vaccination among children less than 2 years of age: analysis of reports from the vaccine adverse event reporting system, 1990–2003. *Pediatrics* 2005;115:453–60.
194. France EK, Glanz JM, Xu S, et al. Safety of the trivalent inactivated influenza vaccine among children: a population-based study. *Arch Pediatr Adolesc Med* 2004;158:1031–6.
195. Bierman CW, Shapiro GG, Pierson WE, et al. Safety of influenza vaccination in allergic children. *J Infect Dis* 1977;136(Suppl):S652–5.
196. James JM, Zeiger RS, Lester MR, et al. Safe administration of influenza vaccine to patients with egg allergy. *J Pediatr* 1998;133:624–8.
197. Murphy KR, Strunk RC. Safe administration of influenza vaccine in asthmatic children hypersensitive to egg proteins. *J Pediatr* 1985;106:931–3.
198. Zeiger RS. Current issues with influenza vaccination in egg allergy. *J Allergy Clin Immunol* 2002;110:834–40.
199. Aberer W. Vaccination despite thimerosal sensitivity. *Contact Dermatitis* 1991;24:6–10.
200. Kirkland LR. Ocular sensitivity to thimerosal: a problem with hepatitis B vaccine? *South Med J* 1990;83:497–9.
201. Schonberger LB, Bregman DJ, Sullivan-Bolyai JZ, et al. Guillain-Barre syndrome following vaccination in the National Influenza Immunization Program, United States, 1976–1977. *Am J Epidemiol* 1979;110:105–23.
202. Safranek TJ, Lawrence DN, Kurland LT, et al. Reassessment of the association between Guillain-Barre syndrome and receipt of swine influenza vaccine in 1976–1977: results of a two-state study. *Expert Neurology Group. Am J Epidemiol* 1991;133:940–51.
203. Ropper AH. The Guillain-Barre syndrome. *N Engl J Med* 1992;326:1130–6.
204. Hurwitz ES, Schonberger LB, Nelson DB, et al. Guillain-Barre syndrome and the 1978–1979 influenza vaccine. *N Engl J Med* 1981;304:1557–61.
205. Kaplan JE, Katona P, Hurwitz ES, et al. Guillain-Barre syndrome in the United States, 1979–1980 and 1980–1981. Lack of an association with influenza vaccination. *JAMA* 1982;248:698–700.
206. Chen R, Kent J, Rhodes P, et al. Investigations of a possible association between influenza vaccination and Guillain-Barre syndrome in the United States, 1990–1991 [Abstract 040]. *Post Marketing Surveillance* 1992;6:5–6.
207. Lasky T, Terracciano GJ, Magder L, et al. The Guillain-Barre syndrome and the 1992–1993 and 1993–1994 influenza vaccines. *N Engl J Med* 1998;339:1797–802.
208. Haber P, DeStefano F, Angulo FJ, et al. Guillain-Barre syndrome following influenza vaccination. *JAMA* 2004;292:2478–81.
209. Flewett TH, Hoult JG. Influenzal encephalopathy and postinfluenzal encephalitis. *Lancet* 1958;2(7036):11–5.
210. Horner FA. Neurologic disorders after Asian influenza. *N Engl J Med* 1958;258:983–5.
211. Jacobs BC, Rothbarth PH, van der Meche FG, et al. The spectrum of antecedent infections in Guillain-Barre syndrome: a case-control study. *Neurology* 1998;51:1110–5.
212. Guarino M, Casmiro M, D'Alessandro R. *Campylobacter jejuni* infection and Guillain-Barre syndrome: a case-control study. Emilia-Romagna Study Group on Clinical and Epidemiological problems in neurology. *Neuroepidemiology* 1998;17:296–302.
213. Sheikh KA, Nachamkin I, Ho TW, et al. *Campylobacter jejuni* lipopolysaccharides in Guillain-Barre syndrome: molecular mimicry and host susceptibility. *Neurology* 1998;51:371–8.
214. Prevots DR, Sutter RW. Assessment of Guillain-Barre syndrome mortality and morbidity in the United States: implications for acute flaccid paralysis surveillance. *J Infect Dis* 1997;175(Suppl 1):S151–5.
215. Barohn RJ, Saperstein DS. Guillain-Barre syndrome and chronic inflammatory demyelinating polyneuropathy. *Semin Neurol* 1998;18:49–61.

216. Maassab HF, Bryant ML. The development of live attenuated cold-adapted influenza virus vaccine for humans. *Rev Med Virol* 1999;9:237-44.
217. Maassab HF, DeBorde DC. Development and characterization of cold-adapted viruses for use as live virus vaccines. *Vaccine* 1985;3:355-69.
218. Clements ML, Stephens I. New and improved vaccines against influenza. In: Levine MM, Woodrow GC, and Kasper JB, eds. *New Generation Vaccines*. New York, NY: Marcel Dekker, Inc.; 1997:645-70.
219. King JC Jr., Lagos R, Bernstein DI, et al. Safety and immunogenicity of low and high doses of trivalent live cold-adapted influenza vaccine administered intranasally as drops or spray to healthy children. *J Infect Dis* 1998;177:1394-7.
220. Belshe RB, Gruber WC, Mendelman PM, et al. Correlates of immune protection induced by live, attenuated, cold-adapted, trivalent, intranasal influenza virus vaccine. *J Infect Dis* 2000;181:1133-7.
221. Boyce TG, Gruber WC, Coleman-Dockery SD, et al. Mucosal immune response to trivalent live attenuated intranasal influenza vaccine in children. *Vaccine* 1999;18:82-8.
222. Zangwill KM, Droge J, Mendelman P, et al. Prospective, randomized, placebo-controlled evaluation of the safety and immunogenicity of three lots of intranasal trivalent influenza vaccine among young children. *Pediatr Infect Dis J* 2001;20:740-6.
223. Bernstein DI, Yan L, Treanor J, et al. Effect of yearly vaccinations with live, attenuated, cold-adapted, trivalent, intranasal influenza vaccines on antibody responses in children. *Pediatr Infect Dis J* 2003;22:28-34.
224. Nolan T, Lee MS, Cordova JM, et al. Safety and immunogenicity of a live-attenuated influenza vaccine blended and filled at two manufacturing facilities. *Vaccine* 2003;21:1224-31.
225. Vesikari T. Randomized, double-blind, placebo-controlled trial of the safety, transmissibility and phenotypic stability of a live, attenuated, cold-adapted influenza virus vaccine (CAIV-T) in children attending day care [Abstract G-450]. Presented at the 41st Annual Interscience Conference on Antimicrobial Agents and Chemotherapy (ICAAC); 2001.
226. Talbot TR, Crocker DD, Peters J. Degree and duration of mucosal shedding following use of trivalent intranasal live attenuated influenza vaccine in adults [Abstract]. Presented at 14th Annual Meeting, Society for Health-care Epidemiology in America, Philadelphia, Pennsylvania; 2004.
227. Ali T, Scott N, Kallas W, et al. Detection of influenza antigen with rapid antibody-based tests after intranasal influenza vaccination (FluMist). *Clin Infect Dis* 2004;38:760-2.
228. Cha TA, Kao K, Zhao J, et al. Genotypic stability of cold-adapted influenza virus vaccine in an efficacy clinical trial. *J Clin Microbiol* 2000;38:839-45.
229. King JC Jr., Fast PE, Zangwill KM, et al. Safety, vaccine virus shedding and immunogenicity of trivalent, cold-adapted, live attenuated influenza vaccine administered to human immunodeficiency virus-infected and noninfected children. *Pediatr Infect Dis J* 2001;20:1124-31.
230. Redding G, Walker RE, Hessel C, et al. Safety and tolerability of cold-adapted influenza virus vaccine in children and adolescents with asthma. *Pediatr Infect Dis J* 2002;21:44-8.
231. Piedra PA, Yan L, Kotloff K, et al. Safety of the trivalent, cold-adapted influenza vaccine in preschool-aged children. *Pediatrics* 2002;110:662-72.
232. Jackson LA, Holmes SJ, Mendelman PM, et al. Safety of a trivalent live attenuated intranasal influenza vaccine, FluMist, administered in addition to parenteral trivalent inactivated influenza vaccine to seniors with chronic medical conditions. *Vaccine* 1999;17:1905-9.
233. King JC Jr., Treanor J, Fast PE, et al. Comparison of the safety, vaccine virus shedding, and immunogenicity of influenza virus vaccine, trivalent, types A and B, live cold-adapted, administered to human immunodeficiency virus (HIV)-infected and non-HIV-infected adults. *J Infect Dis* 2000;181:725-8.
234. Izurieta HS, Haber P, Ball R, et al. Post-licensure surveillance of the first live, cold-adapted influenza vaccine in the U.S. [Abstract]. *Pharmacoepidemiology and Drug Safety* 2004;13:S145.
235. McElhaney JE, Gravenstein S, Upshaw CM, et al. Immune response to influenza vaccination in institutionalized elderly: effect on different T-cell subsets. *Vaccine* 1998;16:403-9.
236. Gross PA, Russo C, Dran S, et al. Time to earliest peak serum antibody response to influenza vaccine in the elderly. *Clin Diagn Lab Immunol* 1997;4:491-2.
237. Brokstad KA, Cox RJ, Olofsson J, et al. Parenteral influenza vaccination induces a rapid systemic and local immune response. *J Infect Dis* 1995;171:198-203.
238. Lawson F, Baker V, Au D, et al. Standing orders for influenza vaccination increased vaccination rates in inpatient settings compared with community rates. *J Gerontol A Biol Sci Med Sci* 2000;55:M522-6.
239. Centers for Medicare and Medicaid Services. Medicare and Medicaid programs; conditions of participation: immunization standards for hospitals, long-term care facilities, and home health agencies. Final rule with comment period. *Federal Register* 2002;67:61808-14. Available at <http://www.cms.hhs.gov/preventiveservices/2h.asp>.
240. Fedson DS, Wajda A, Nicol JP, et al. Disparity between influenza vaccination rates and risks for influenza-associated hospital discharge and death in Manitoba in 1982-1983. *Ann Intern Med* 1992;116:550-5.
241. Bratzler DW, Houck PM, Jiang H, et al. Failure to vaccinate Medicare inpatients: a missed opportunity. *Arch Intern Med* 2002;162:2349-56.
242. Fedson DS, Houck P, Bratzler D. Hospital-based influenza and pneumococcal vaccination: Sutton's Law applied to prevention. *Infect Control Hosp Epidemiol* 2000;21:692-9.
243. American Academy of Pediatrics. Influenza. In: Pickering LK, ed. *Red book: 2003 Report of the Committee on Infectious Diseases*. 26th ed. Elk Grove Village, IL: American Academy of Pediatrics; 2003.
244. Schmid ML, Kudesia G, Wake S, et al. Prospective comparative study of culture specimens and methods in diagnosing influenza in adults. *BMJ* 1998;316:275.
245. Letter M. Rapid diagnostic tests for influenza. *Medical Letter* 1999;41:121-22.
246. Storch GA. Rapid diagnostic tests for influenza. *Curr Opin Pediatr* 2003;15:77-84.
247. Uyeki TM. Influenza diagnosis and treatment in children: a review of studies on clinically useful tests and antiviral treatment for influenza. *Pediatr Infect Dis J* 2003;22:164-77.
248. Tominack RL, Hayden FG. Rimantadine hydrochloride and amantadine hydrochloride use in influenza A virus infections. *Infect Dis Clin North Am* 1987;1:459-78.
249. Hayden FG, Osterhaus AD, Treanor JJ, et al. Efficacy and safety of the neuraminidase inhibitor zanamivir in the treatment of influenza virus infections. GG167 Influenza Study Group. *N Engl J Med* 1997;337:874-80.

250. MIST (Management of Influenza in the Southern Hemisphere Trialists). Randomised trial of efficacy and safety of inhaled zanamivir in treatment of influenza A and B virus infections. The MIST (Management of Influenza in the Southern Hemisphere Trialists) Study Group. *Lancet* 1998;352(9144):1877–81.
251. Makela MJ, Pauksens K, Rostila T, et al. Clinical efficacy and safety of the orally inhaled neuraminidase inhibitor zanamivir in the treatment of influenza: a randomized, double-blind, placebo-controlled European study. *J Infect* 2000;40:42–8.
252. Matsumoto K, Ogawa N, Nerome K, et al. Safety and efficacy of the neuraminidase inhibitor zanamivir in treating influenza virus infection in adults: results from Japan. GG167 Group. *Antivir Ther* 1999;4:61–8.
253. Monto AS, Fleming DM, Henry D, et al. Efficacy and safety of the neuraminidase inhibitor zanamivir in the treatment of influenza A and B virus infections. *J Infect Dis* 1999;180:254–61.
254. Lalezari J, Champion K, Keene O, et al. Zanamivir for the treatment of influenza A and B infection in high-risk patients: a pooled analysis of randomized controlled trials. *Arch Intern Med* 2001;161:212–7.
255. Treanor JJ, Hayden FG, Vrooman PS, et al. Efficacy and safety of the oral neuraminidase inhibitor oseltamivir in treating acute influenza: a randomized controlled trial. US Oral Neuraminidase Study Group. *JAMA* 2000;283:1016–24.
256. Nicholson KG, Aoki FY, Osterhaus AD, et al. Efficacy and safety of oseltamivir in treatment of acute influenza: a randomised controlled trial. Neuraminidase Inhibitor Flu Treatment Investigator Group. *Lancet* 2000;355(9218):1845–50.
257. Hedrick JA, Barzilai A, Behre U, et al. Zanamivir for treatment of symptomatic influenza A and B infection in children five to twelve years of age: a randomized controlled trial. *Pediatr Infect Dis J* 2000;19:410–7.
258. Whitley RJ, Hayden FG, Reisinger KS, et al. Oral oseltamivir treatment of influenza in children. *Pediatr Infect Dis J* 2001;20:127–33.
259. Glaxo Wellcome. Relenza (zanamivir for inhalation) [Product information]. Research Triangle Park, NC: Glaxo Wellcome, Inc.; 2001.
260. Roche Laboratories I. Tamiflu (oseltamivir phosphate) capsules [Product information]. Nutley, NJ: Roche Laboratories, Inc.; 2000.
261. Murphy KR, Eivindson A, Pauksens K. Efficacy and safety of inhaled zanamivir for the treatment of influenza in patients with asthma or chronic obstructive pulmonary disease: a double-blind, randomised, placebo-controlled, multicentre study. *Clin Drug Invest* 2000;20:337–49.
262. Uyeki T, Winquist A. Influenza. *Clin Evid* 2002;(7):645–51.
263. Cooper NJ, Sutton AJ, Abrams KR, et al. Effectiveness of neuraminidase inhibitors in treatment and prevention of influenza A and B: systematic review and meta-analyses of randomised controlled trials. *BMJ* 2003;326:1235.
264. Jefferson T, Deeks JJ, Demicheli V, et al. Amantadine and rimantadine for preventing and treating influenza A in adults. *Cochrane Database Syst Rev* 2004;(3):CD001169.
265. Jefferson T, Demicheli V, Deeks J, et al. Neuraminidase inhibitors for preventing and treating influenza in healthy adults. *Cochrane Database Syst Rev* 2000;(2):CD001265.
266. Nicholson KG. Use of antivirals in influenza in the elderly: prophylaxis and therapy. *Gerontology* 1996;42:280–9.
267. Wintermeyer SM, Nahata MC. Rimantadine: a clinical perspective. *Ann Pharmacother* 1995;29:299–310.
268. Martin C, Mahoney P, Ward P. Oral oseltamivir reduces febrile illness in patients considered at high risk of influenza complications [Abstract W22-7]. In: Options for the control of influenza IV. New York, NY: Excerpta Medica;2001:807–11.
269. Gravenstein S, Johnston SL, Loeschel E, et al. Zanamivir: a review of clinical safety in individuals at high risk of developing influenza-related complications. *Drug Saf* 2001;24:1113–25.
270. Bowles SK, Lee W, Simor AE, et al. Use of oseltamivir during influenza outbreaks in Ontario nursing homes, 1999–2000. *J Am Geriatr Soc* 2002;50:608–16.
271. Kaiser L, Wat C, Mills T, et al. Impact of oseltamivir treatment on influenza-related lower respiratory tract complications and hospitalizations. *Arch Intern Med* 2003;163:1667–72.
272. Thompson J, Fleet W, Lawrence E, et al. A comparison of acetaminophen and rimantadine in the treatment of influenza A infection in children. *J Med Virol* 1987;21:249–55.
273. Hall CB, Dolin R, Gala CL, et al. Children with influenza A infection: treatment with rimantadine. *Pediatrics* 1987;80:275–82.
274. Guay DR. Amantadine and rimantadine prophylaxis of influenza A in nursing homes. A tolerability perspective. *Drugs Aging* 1994;5:8–19.
275. Patriarca PA, Kater NA, Kendal AP, et al. Safety of prolonged administration of rimantadine hydrochloride in the prophylaxis of influenza A virus infections in nursing homes. *Antimicrob Agents Chemother* 1984;26:101–3.
276. Arden NH, Patriarca PA, Fasano MB, et al. The roles of vaccination and amantadine prophylaxis in controlling an outbreak of influenza A (H3N2) in a nursing home. *Arch Intern Med* 1988;148:865–8.
277. Hayden FG, Atmar RL, Schilling M, et al. Use of the selective oral neuraminidase inhibitor oseltamivir to prevent influenza. *N Engl J Med* 1999;341:1336–43.
278. Hayden FG, Gubareva LV, Monto AS, et al. Inhaled zanamivir for the prevention of influenza in families. Zanamivir Family Study Group. *N Engl J Med* 2000;343:1282–9.
279. Schilling M, Povinelli L, Krause P, et al. Efficacy of zanamivir for chemoprophylaxis of nursing home influenza outbreaks. *Vaccine* 1998;16:1771–4.
280. Lee C, Loeb M, Phillips A, et al. Zanamivir use during transmission of amantadine-resistant influenza A in a nursing home. *Infect Control Hosp Epidemiol* 2000;21:700–4.
281. Parker R, Loewen N, Skowronski D. Experience with oseltamivir in the control of a nursing home influenza B outbreak. *Can Commun Dis Rep* 2001;27:37–40.
282. Woods JM, Bethell RC, Coates JA, et al. 4-Guanidino-2,4-dideoxy-2,3-dehydro-N-acetylneuraminic acid is a highly effective inhibitor both of the sialidase (neuraminidase) and of growth of a wide range of influenza A and B viruses in vitro. *Antimicrob Agents Chemother* 1993;37:1473–9.
283. Hayden FG, Rollins BS, Madren LK. Anti-influenza virus activity of the neuraminidase inhibitor 4-guanidino-Neu5Ac2en in cell culture and in human respiratory epithelium. *Antiviral Res* 1994;25:123–31.
284. Mendel DB, Tai CY, Escarpe PA, et al. Oral administration of a prodrug of the influenza virus neuraminidase inhibitor GS 4071 protects mice and ferrets against influenza infection. *Antimicrob Agents Chemother* 1998;42:640–6.
285. Sidwell RW, Huffman JH, Barnard DL, et al. Inhibition of influenza virus infections in mice by GS4104, an orally effective influenza virus neuraminidase inhibitor. *Antiviral Res* 1998;37:107–20.

286. Hayden FG, Rollins BS. In vitro activity of the neuraminidase inhibitor GS4071 against influenza viruses [Abstract 159]. *Antiviral Res* 1997;34:A86.
287. Mendel DB, Tai CY, Escarpe PA. GS4071 is a potent and selective inhibitor of the growth and neuraminidase activity of influenza A and B viruses in vitro [Abstract 111]. *Antiviral Res* 1997;34:A73.
288. Ryan DM, Ticehurst J, Dempsey MH, et al. Inhibition of influenza virus replication in mice by GG167 (4-guanidino-2,4-dideoxy-2,3-dehydro-N-acetylneuraminic acid) is consistent with extracellular activity of viral neuraminidase (sialidase). *Antimicrob Agents Chemother* 1994;38:2270–5.
289. Ryan DM, Ticehurst J, Dempsey MH. GG167 (4-guanidino-2,4-dideoxy-2,3-dehydro-N-acetylneuraminic acid) is a potent inhibitor of influenza virus in ferrets. *Antimicrob Agents Chemother* 1995;39:2583–4.
290. Welliver R, Monto AS, Carewicz O, et al. Effectiveness of oseltamivir in preventing influenza in household contacts: a randomized controlled trial. *JAMA* 2001;285:748–54.
291. Hayden FG, Jennings L, Robson R, et al. Oral oseltamivir in human experimental influenza B infection. *Antivir Ther* 2000;5:205–13.
292. Food and Drug Administration. Subject: safe and appropriate use of influenza drugs [Public Health Advisory]. Rockville, MD: US Department of Health and Human Services, Food and Drug Administration; 2000.
293. Monto AS, Pichichero ME, Blanckenberg SJ, et al. Zanamivir prophylaxis: an effective strategy for the prevention of influenza types A and B within households. *J Infect Dis* 2002;186:1582–8.
294. Peters PH Jr., Gravenstein S, Norwood P, et al. Long-term use of oseltamivir for the prophylaxis of influenza in a vaccinated frail older population. *J Am Geriatr Soc* 2001;49:1025–31.
295. Webster A, Boyce M, Edmundson S, et al. Coadministration of orally inhaled zanamivir with inactivated trivalent influenza vaccine does not adversely affect the production of antihaemagglutinin antibodies in the serum of healthy volunteers. *Clin Pharmacokinet* 1999;36(Suppl 1):51–8.
296. Patriarca PA, Arden NH, Koplan JP, et al. Prevention and control of type A influenza infections in nursing homes. Benefits and costs of four approaches using vaccination and amantadine. *Ann Intern Med* 1987;107:732–40.
297. Gomolin IH, Leib HB, Arden NH, et al. Control of influenza outbreaks in the nursing home: guidelines for diagnosis and management. *J Am Geriatr Soc* 1995;43:71–4.
298. Garner JS. Guideline for isolation precautions in hospitals. The Hospital Infection Control Practices Advisory Committee. *Infect Control Hosp Epidemiol* 1996;17:53–80.
299. Bradley SF. Prevention of influenza in long-term-care facilities. Long-Term-Care Committee of the Society for Health-care Epidemiology of America. *Infect Control Hosp Epidemiol* 1999;20:629–37.
300. Shijubo N, Yamada G, Takahashi M, et al. Experience with oseltamivir in the control of nursing home influenza A outbreak. *Intern Med* 2002;41:366–70.
301. Forest Pharmaceuticals. Flumadine syrup (rimantadine hydrochloride syrup) [package insert]. St. Louis, MO: Forest Pharmaceuticals; 2001.
302. Keyser LA, Karl M, Nafziger AN, et al. Comparison of central nervous system adverse effects of amantadine and rimantadine used as sequential prophylaxis of influenza A in elderly nursing home patients. *Arch Intern Med* 2000;160:1485–8.
303. Endo Pharmaceuticals. Symmetrel [package insert]. Dayton, NJ: Endo Pharmaceuticals, Inc; 2000.
304. Soung LS, Ing TS, Daugirdas JT, et al. Amantadine hydrochloride pharmacokinetics in hemodialysis patients. *Ann Intern Med* 1980;93:46–9.
305. Capparelli EV, Stevens RC, Chow MS, et al. Rimantadine pharmacokinetics in healthy subjects and patients with end-stage renal failure. *Clin Pharmacol Ther* 1988;43:536–41.
306. Cass LM, Efthymiopoulos C, Marsh J, et al. Effect of renal impairment on the pharmacokinetics of intravenous zanamivir. *Clin Pharmacokinet* 1999;36(Suppl 1):13–9.
307. Calfee DP, Peng AW, Cass LM, et al. Safety and efficacy of intravenous zanamivir in preventing experimental human influenza A virus infection. *Antimicrob Agents Chemother* 1999;43:1616–20.
308. Cass LM, Efthymiopoulos C, Bye A. Pharmacokinetics of zanamivir after intravenous, oral, inhaled or intranasal administration to healthy volunteers. *Clin Pharmacokinet* 1999;36(Suppl 1):1–11.
309. Bardsley-Elliot A, Noble S. Oseltamivir. *Drugs* 1999;58:851–60; discussion: 861–2.
310. Schnack H, Wewalka F, Guttman G. Liver function during amantadine hydrochloride medication in compensated liver disease. *Int Z Klin Pharmakol Ther Toxikol* 1969;2:185–7.
311. Atkinson WL, Arden NH, Patriarca PA, et al. Amantadine prophylaxis during an institutional outbreak of type A (H1N1) influenza. *Arch Intern Med* 1986;146:1751–6.
312. Soo W. Adverse effects of rimantadine: summary from clinical trials. *J Respir Dis* 1989;10(Suppl):S26–31.
313. Bleidner WE, Harmon JB, Hewes WE, et al. Absorption, distribution and excretion of amantadine hydrochloride. *J Pharmacol Exp Ther* 1965;150:484–90.
314. Douglas RG Jr. Prophylaxis and treatment of influenza. *N Engl J Med* 1990;322:443–50.
315. Aoki FY, Sitar DS. Amantadine kinetics in healthy elderly men: implications for influenza prevention. *Clin Pharmacol Ther* 1985;37:137–44.
316. Aoki FY, Sitar DS. Clinical pharmacokinetics of amantadine hydrochloride. *Clin Pharmacokinet* 1988;14:35–51.
317. Wills RJ, Belshe R, Tomlinsin D, et al. Pharmacokinetics of rimantadine hydrochloride in patients with chronic liver disease. *Clin Pharmacol Ther* 1987;42:449–54.
318. Cass LM, Brown J, Pickford M, et al. Pharmacoscintigraphic evaluation of lung deposition of inhaled zanamivir in healthy volunteers. *Clin Pharmacokinet* 1999;36(Suppl 1):21–31.
319. He G, Massarella J, Ward P. Clinical pharmacokinetics of the prodrug oseltamivir and its active metabolite Ro 64-0802. *Clin Pharmacokinet* 1999;37:471–84.
320. Dolin R, Reichman RC, Madore HP, et al. A controlled trial of amantadine and rimantadine in the prophylaxis of influenza A infection. *N Engl J Med* 1982;307:580–4.
321. Hayden FG, Treanor JJ, Fritz RS, et al. Use of the oral neuraminidase inhibitor oseltamivir in experimental human influenza: randomized controlled trials for prevention and treatment. *JAMA* 1999;282:1240–6.
322. Daniel MJ, Barnett JM, Pearson BA. The low potential for drug interactions with zanamivir. *Clin Pharmacokinet* 1999;36(Suppl 1):41–50.
323. Belshe RB, Smith MH, Hall CB, et al. Genetic basis of resistance to rimantadine emerging during treatment of influenza virus infection. *J Virol* 1988;62:1508–12.

324. Saito R, Oshitani H, Masuda H, et al. Detection of amantadine-resistant influenza A virus strains in nursing homes by PCR-restriction fragment length polymorphism analysis with nasopharyngeal swabs. *J Clin Microbiol* 2002;40:84–8.
325. Hayden FG, Sperber SJ, Belshe RB, et al. Recovery of drug-resistant influenza A virus during therapeutic use of rimantadine. *Antimicrob Agents Chemother* 1991;35:1741–7.
326. Houck P, Hemphill M, LaCroix S, et al. Amantadine-resistant influenza A in nursing homes. Identification of a resistant virus prior to drug use. *Arch Intern Med* 1995;155:533–7.
327. Hayden FG, Belshe RB, Clover RD, et al. Emergence and apparent transmission of rimantadine-resistant influenza A virus in families. *N Engl J Med* 1989;321:1696–702.
328. Mast EE, Harmon MW, Gravenstein S, et al. Emergence and possible transmission of amantadine-resistant viruses during nursing home outbreaks of influenza A (H3N2). *Am J Epidemiol* 1991;134:988–97.
329. Hayden FG, Hay AJ. Emergence and transmission of influenza A viruses resistant to amantadine and rimantadine. *Curr Top Microbiol Immunol* 1992;176:119–30.
330. Degelau J, Somani SK, Cooper SL, et al. Amantadine-resistant influenza A in a nursing facility. *Arch Intern Med* 1992;152:390–2.
331. Ziegler T, Hemphill ML, Ziegler ML, et al. Low incidence of rimantadine resistance in field isolates of influenza A viruses. *J Infect Dis* 1999;180:935–9.
332. Gubareva LV, Robinson MJ, Bethell RC, et al. Catalytic and framework mutations in the neuraminidase active site of influenza viruses that are resistant to 4-guanidino-Neu5Ac2en. *J Virol* 1997;71:3385–90.
333. Colacino JM, Laver WG, Air GM. Selection of influenza A and B viruses for resistance to 4-guanidino-Neu5Ac2en in cell culture. *J Infect Dis* 1997;176(Suppl 1):S66–8.
334. Gubareva LV, Bethell R, Hart GJ, et al. Characterization of mutants of influenza A virus selected with the neuraminidase inhibitor 4-guanidino-Neu5Ac2en. *J Virol* 1996;70:1818–27.
335. Blick TJ, Tiong T, Sahasrabudhe A, et al. Generation and characterization of an influenza virus neuraminidase variant with decreased sensitivity to the neuraminidase-specific inhibitor 4-guanidino-Neu5Ac2en. *Virology* 1995;214:475–84.
336. McKimm-Breschkin JL, Blick TJ, Sahasrabudhe A, et al. Generation and characterization of variants of NWS/G70C influenza virus after in vitro passage in 4-amino-Neu5Ac2en and 4-guanidino-Neu5Ac2en. *Antimicrob Agents Chemother* 1996;40:40–6.
337. Staschke KA, Colacino JM, Baxter AJ, et al. Molecular basis for the resistance of influenza viruses to 4-guanidino-Neu5Ac2en. *Virology* 1995;214:642–6.
338. McKimm-Breschkin JL, Sahasrabudhe A, Blick TJ, et al. Mutations in a conserved residue in the influenza virus neuraminidase active site decreases sensitivity to Neu5Ac2en-derived inhibitors. *J Virol* 1998;72:2456–62.
339. Tai CY, Escarpe PA, Sidwell RW, et al. Characterization of human influenza virus variants selected in vitro in the presence of the neuraminidase inhibitor GS 4071. *Antimicrob Agents Chemother* 1998;42:3234–41.
340. Hay AJ, Wolstenholme AJ, Skehel JJ, et al. The molecular basis of the specific anti-influenza action of amantadine. *Embo J* 1985;4:3021–4.
341. Appleyard G. Amantadine-resistance as a genetic marker for influenza viruses. *J Gen Virol* 1977;36:249–55.
342. Barnett JM, Cadman A, Gor D, et al. Zanamivir susceptibility monitoring and characterization of influenza virus clinical isolates obtained during phase II clinical efficacy studies. *Antimicrob Agents Chemother* 2000;44:78–87.
343. Gubareva LV, Matrosovich MN, Brenner MK, et al. Evidence for zanamivir resistance in an immunocompromised child infected with influenza B virus. *J Infect Dis* 1998;178:1257–62.
344. Gubareva LV, Kaiser L, Matrosovich MN, et al. Selection of influenza virus mutants in experimentally infected volunteers treated with oseltamivir. *J Infect Dis* 2001;183:523–31.
345. Jackson HC, Roberts N, Wang ZM, et al. Management of influenza: use of new antivirals and resistance in perspective. *Clin Drug Invest* 2000;20:447–54.
346. Kiso M, Mitamura K, Sakai-Tagawa Y, et al. Resistant influenza A viruses in children treated with oseltamivir: descriptive study. *Lancet* 2004;364(9436):759–65.
347. Tisdale M. Monitoring of viral susceptibility: new challenges with the development of influenza NA inhibitors. *Rev Med Virol* 2000;10:45–55.
348. Gubareva LV, Webster RG, Hayden FG. Detection of influenza virus resistance to neuraminidase inhibitors by an enzyme inhibition assay. *Antiviral Res* 2002;53:47–61.
349. Zambon M, Hayden FG. Position statement: global neuraminidase inhibitor susceptibility network. *Antiviral Res* 2001;49:147–56.

Advisory Committee on Immunization Practices Membership List, January 2005

Chairman: Myron J. Levin, MD, Professor of Pediatrics and Medicine, University of Colorado Health Sciences Center, Denver, Colorado

Executive Secretary: Larry Pickering, MD, National Immunization Program, Centers for Disease Control and Prevention, Atlanta, Georgia

Members: Jon S. Abramson, MD, Wake Forest University School of Medicine, Winston-Salem, North Carolina; Ban Mishu Allos, MD, Vanderbilt University School of Medicine, Nashville, Tennessee; Guthrie S. Birkhead, MD, New York State Department of Health, Albany, New York; Judith Campbell, MD, Baylor College of Medicine, Houston, Texas; Reginald Finger, MD, Focus on the Family, Colorado Springs, Colorado; Janet Gildsford, MD, University of Michigan, Ann Arbor, Michigan; Tracy Lieu, MD, Harvard Pilgrim Health Care and Harvard Medical School, Boston, Massachusetts; Edgar Marcuse, MD, Children's Hospital and Regional Medical Center, Seattle, Washington; Julia Morita, MD, Chicago Department of Health, Chicago, Illinois; Gregory Poland, MD, Mayo Clinic College of Medicine, Rochester, Minnesota; John B. Salamone, National Italian American Foundation, Washington, DC; Patricia Stinchfield, Children's Hospital and Clinics, St. Paul, Minnesota; John J. Treanor, MD, University of Rochester School of Medicine and Dentistry, Rochester, New York; Robin Womeodu, MD, University of Tennessee Health Sciences Center, Memphis, Tennessee.

Ex-Officio Members: James E. Cheek, MD, Indian Health Service, Albuquerque, New Mexico; Stephen Phillips, DO, Department of Defense, Falls Church, Virginia; Geoffrey S. Evans, MD, Health Resources and Services Administration, Rockville, Maryland; Bruce Gellin, MD, National Vaccine Program Office, Washington, DC; Linda Murphy, Centers for Medicare and Medicaid Services, Baltimore, Maryland; George T. Curlin, MD, National Institutes of Health, Bethesda, Maryland; Norman Baylor, MD, Food and Drug Administration, Bethesda, Maryland; Kristin Lee Nichol, MD, Department of Veterans Affairs, Minneapolis, Minnesota.

Liaison Representatives: American Academy of Family Physicians, Jonathan Temte, MD, Clarence, New York, and Richard Clover, MD, Louisville, Kentucky; American Academy of Pediatrics, Margaret Rennels, MD, Baltimore, Maryland, and Carol Baker, MD, Houston, Texas; American Association of Health Plans, Robert Scalettar, MD, North Haven, Connecticut; American College Health Association, James C. Turner, MD, Charlottesville, Virginia; American College of Obstetricians and Gynecologists, Stanley Gall, MD, Louisville, Kentucky; American College of Physicians, Kathleen Neuzil, MD, Seattle, Washington; American Medical Association, Litjen Tan, PhD, Chicago, Illinois; American Pharmacists Association, Stephan L. Foster, PharmD, Memphis, Tennessee; Association of Teachers of Preventive Medicine, W. Paul McKinney, MD, Louisville, Kentucky; Biotechnology Industry Organization, Clement Lewin, PhD, Cambridge, Massachusetts; Canadian National Advisory Committee on Immunization, Monica Naus, MD, Vancouver, British Columbia; Health-care Infection Control Practices Advisory Committee, Steve Gordon, MD, Cleveland, Ohio; Infectious Diseases Society of America, Samuel L. Katz, MD, Durham, North Carolina, and William Schaffner, MD, Nashville, Tennessee; London Department of Health, David M. Salisbury, MD, London, United Kingdom; National Association of County and City Health Officials, Nancy Bennett, MD, Rochester, New York; National Coalition for Adult Immunization, David A. Neumann, PhD, Bethesda, Maryland; National Immunization Council and Child Health Program, Mexico, Romeo Rodriguez, Mexico City, Mexico; National Medical Association, Dennis A. Brooks, MD, Baltimore, Maryland; National Vaccine Advisory Committee, Charles Helms, MD, PhD, Iowa City, Iowa; Pharmaceutical Research and Manufacturers of America, Damian A. Braga, Swiftwater, Pennsylvania, and Peter Paradiso, PhD, Collegeville, Pennsylvania; and Society for Adolescent Medicine, Amy Middleman, MD, Houston, Texas.

ACIP Influenza Working Group

Chair: Ban Mishu Allos, MD, Nashville, Tennessee.

Members: Jon Abramson, MD, Winston-Salem, North Carolina; Carolyn Bridges, MD, Atlanta, Georgia; Robert Chen, MD, Atlanta, Georgia; Richard Clover, MD, Louisville, Kentucky; Nancy Cox, PhD, Atlanta, Georgia; Keiji Fukuda, MD, Atlanta, Georgia; Stanley Gall, MD, Louisville, Kentucky; Randolph Graydon, Baltimore, Maryland; Celine Hanson, MD, Houston, Texas; Scott Harper, MD, Atlanta, Georgia; J. Henry Hershey, MD, Christiansburg, Virginia; John Iskander, MD, Atlanta, Georgia; Marika Iwane, PhD, Atlanta, Georgia; Susan Lett, MD, Boston, Massachusetts; Roland Levandowski, MD, Bethesda, Maryland; Alison Mawle, MD, Atlanta, Georgia; Kathleen Neuzil, MD, Seattle, Washington; Kristin Lee Nichol, MD, Minneapolis, Minnesota; Dennis O'Mara, Atlanta, Georgia; Margaret Rennels, MD, Baltimore, Maryland; William Schaffner, MD, Nashville, Tennessee; Jane Seward, MBBS, Atlanta, Georgia; and Litjen Tan, PhD, Chicago, Illinois.

The *Morbidity and Mortality Weekly Report (MMWR)* Series is prepared by the Centers for Disease Control and Prevention (CDC) and is available free of charge in electronic format and on a paid subscription basis for paper copy. To receive an electronic copy each week, send an e-mail message to listserv@listserv.cdc.gov. The body content should read *SUBscribe mmwr-toc*. Electronic copy also is available from CDC's World-Wide Web server at <http://www.cdc.gov/mmwr> or from CDC's file transfer protocol server at <ftp://ftp.cdc.gov/pub/publications/mmwr>. To subscribe for paper copy, contact Superintendent of Documents, U.S. Government Printing Office, Washington, DC 20402; telephone 202-512-1800.

Data in the weekly *MMWR* are provisional, based on weekly reports to CDC by state health departments. The reporting week concludes at close of business on Friday; compiled data on a national basis are officially released to the public on the following Friday. Address inquiries about the *MMWR* Series, including material to be considered for publication, to Editor, *MMWR* Series, Mailstop K-95, CDC, 1600 Clifton Rd., N.E., Atlanta, GA 30333; telephone 888-232-3228.

All material in the *MMWR* Series is in the public domain and may be used and reprinted without permission; citation as to source, however, is appreciated.

All *MMWR* references are available on the Internet at <http://www.cdc.gov/mmwr>. Use the search function to find specific articles.

Use of trade names and commercial sources is for identification only and does not imply endorsement by the U.S. Department of Health and Human Services.

References to non-CDC sites on the Internet are provided as a service to *MMWR* readers and do not constitute or imply endorsement of these organizations or their programs by CDC or the U.S. Department of Health and Human Services. CDC is not responsible for the content of these sites. URL addresses listed in *MMWR* were current as of the date of publication.

Appendix C

CDC Adult Immunization Schedule, 2004-2005

[2005-2006 version to be released in fall 2005]

For published updates during the flu season visit: <http://www.cdc.gov/nip/recs/adult-schedule.htm>

Recommended Adult Immunization Schedule

UNITED STATES · OCTOBER 2004 – SEPTEMBER 2005

Summary of Recommendations Published by

The Advisory Committee on Immunization Practices



**DEPARTMENT OF HEALTH AND HUMAN SERVICES
CENTERS FOR DISEASE CONTROL AND PREVENTION**



Recommended Adult Immunization Schedule by Vaccine and Age Group

UNITED STATES · OCTOBER 2004–SEPTEMBER 2005

Age group (yrs) ▶	19–49	50–64	≥65
Vaccine ▼			
Tetanus, Diphtheria (Td)*	1 dose booster every 10 years ¹		
Influenza	1 dose annually ²	1 dose annually ²	
Pneumococcal (polysaccharide)	1 dose ^{3,4}		1 dose ^{3,4}
Hepatitis B*	3 doses (0, 1–2, 4–6 months) ⁵		
Hepatitis A*	2 doses (0, 6–12 months) ⁶		
Measles, Mumps, Rubella (MMR)*	1 or 2 doses ⁷		
Varicella*	2 doses (0, 4–8 weeks) ⁸		
Meningococcal (polysaccharide)		1 dose ⁹	

*Covered by the Vaccine Injury Compensation Program.
See Footnotes for Recommended Adult Immunization Schedule on back cover.

 For all persons in this group

 For persons lacking documentation of vaccination or evidence of disease

 For persons at risk (i.e., with medical/exposure indications)

The Recommended Adult Immunization Schedule is Approved by the Advisory Committee on Immunization Practices (ACIP), the American College of Obstetricians and Gynecologists (ACOG), and the American Academy of Family Physicians (AAFP)

This schedule indicates the recommended age groups for routine administration of currently licensed vaccines for persons aged ≥19 years. Licensed combination vaccines may be used whenever any components of the combination are indicated and when the vaccine's other components are not contraindicated. Providers should consult manufacturers' package inserts for detailed recommendations.

Report all clinically significant postvaccination reactions to the Vaccine Adverse Event Reporting System (VAERS). Reporting forms and instructions on filing a VAERS report are available by telephone, 800-822-7967, or from the VAERS website at <http://www.vaers.org>.

Information on how to file a Vaccine Injury Compensation Program claim is available at <http://www.hrsa.gov/osp/vicp> or by telephone, 800-338-2382. To file a claim for vaccine injury, contact the U.S. Court of Federal Claims, 717 Madison Place, N.W., Washington, DC 20005, telephone 202-219-9657.

Additional information about the vaccines listed above and contraindications for immunization is available at <http://www.cdc.gov/nip> or 800-CDC-INFO [800-232-4636] (English and Spanish).

Recommended Adult Immunization Schedule by Vaccine and Medical and Other Indications

UNITED STATES · OCTOBER 2004–SEPTEMBER 2005

Indication ▶ Vaccine ▼	Pregnancy	Diabetes, heart disease, chronic pulmonary disease, chronic liver disease (including chronic alcoholism)	Congenital immunodeficiency, cochlear implants leukemia, lymphoma, generalized malignancy, therapy with alkylating agents, antimetabolites, CSF** leaks, radiation or large amounts of corticosteroids	Renal failure/end stage renal disease, recipients of hemodialysis or clotting factor concentrates	Asplenia (including elective splenectomy and terminal complement component deficiencies)	HIV*** infection	Health-care workers
Tetanus, Diphtheria (Td)*.1							
Influenza ²		A, B			C		
Pneumococcal (polysaccharide) ^{3,4}		B		D	D, E, F	D, G	
Hepatitis B*.5				H			
Hepatitis A*.6		I					L
Measles, Mumps, Rubella (MMR)*.7	////		////			J	
Varicella*.8	////		//// K			////	

*Covered by the Vaccine Injury Compensation Program.

**Cerebrospinal fluid.

***Human immunodeficiency virus.

See Special Notes for Medical and Other Indications below. Also see Footnotes for Recommended Adult Immunization Schedule on back cover.

 For all persons in this group

 For persons lacking documentation of vaccination or evidence of disease

 For persons at risk (i.e., with medical/exposure indications)

 Contraindicated

Special Notes for Medical and Other Indications

A. Although chronic liver disease and alcoholism are not indications for influenza vaccination, administer 1 dose annually if the patient is aged ≥ 50 years, has other indications for influenza vaccine, or requests vaccination.

B. Asthma is an indication for influenza vaccination but not for pneumococcal vaccination.

C. No data exist specifically on the risk for severe or complicated influenza infections among persons with asplenia. However, influenza is a risk factor for secondary bacterial infections that can cause severe disease among persons with asplenia.

D. For persons aged < 65 years, revaccinate once after ≥ 5 years have elapsed since initial vaccination.

E. Administer meningococcal vaccine and consider *Haemophilus influenzae* type b vaccine.

F. For persons undergoing elective splenectomy, vaccinate ≥ 2 weeks before surgery.

G. Vaccinate as soon after diagnosis as possible.

H. For hemodialysis patients, use special formulation of vaccine (40 $\mu\text{g}/\text{mL}$) or two 20 $\mu\text{g}/\text{mL}$ doses administered at one body site. Vaccinate early in the course of renal disease. Assess antibody titers to hepatitis B surface antigen (anti-HB) levels annually. Administer additional doses if anti-HB levels decline to < 10 mIU/mL.

I. For all persons with chronic liver disease.

J. Withhold MMR or other measles-containing vaccines from HIV-infected persons with evidence of severe immunosuppression (see *MMWR* 1998;47 [No. RR-8]:21–2 and *MMWR* 2002;51 [No. RR-2]:22–4).

K. Persons with impaired humoral immunity but intact cellular immunity may be vaccinated (see *MMWR* 1999;48[No. RR-6]).

L. No data to support a recommendation.

Footnotes

Recommended Adult Immunization Schedule • UNITED STATES • OCTOBER 2004–SEPTEMBER 2005

- 1. Tetanus and diphtheria (Td).** Adults, including pregnant women with uncertain history of a complete primary vaccination series, should receive a primary series of Td. A primary series for adults is 3 doses; administer the first 2 doses at least 4 weeks apart and the 3rd dose 6–12 months after the second. Administer 1 dose if the person received the primary series and if the last vaccination was received ≥ 10 years previously. Consult recommendations for administering Td as prophylaxis in wound management (see *MMWR* 1991;40[No. RR-10]). The American College of Physicians Task Force on Adult Immunization supports a second option for Td use in adults: a single Td booster at age 50 years for persons who have completed the full pediatric series, including the teenage/young adult booster.
- 2. Influenza vaccination.** The Advisory Committee on Immunization Practices (ACIP) recommends inactivated influenza vaccination for the following indications, when vaccine is available. *Medical indications:* chronic disorders of the cardiovascular or pulmonary systems, including asthma; chronic metabolic diseases, including diabetes mellitus, renal dysfunction, hemoglobinopathies, or immunosuppression (including immunosuppression caused by medications or by human immunodeficiency virus [HIV]); and pregnancy during the influenza season. *Occupational indications:* health-care workers and employees of long-term-care and assisted living facilities. *Other indications:* residents of nursing homes and other long-term-care facilities; persons likely to transmit influenza to persons at high risk (i.e., in-home caregivers to persons with medical indications, household/close contacts and out-of-home caregivers of children aged 0–23 months, household members and caregivers of elderly persons and adults with high-risk conditions); and anyone who wishes to be vaccinated. For healthy persons aged 5–49 years without high-risk conditions who are not contacts of severely immunocompromised persons in special care units, either the inactivated vaccine or the intranasally administered influenza vaccine (FluMist®) may be administered (see *MMWR* 2004;53[No. RR-6]).
Note: Because of the influenza vaccine shortage in 2004, the Centers for Disease Control and Prevention (CDC) and the ACIP had modified the influenza vaccine recommendations (see *MMWR* 2004;53[39];923-924). These were further modified on December 17, 2004, and this schedule reflects those changes (i.e., routinely recommending influenza vaccination of persons 50-64 years of age) (see *MMWR* 2004;53[50];1183-1184).
- 3. Pneumococcal polysaccharide vaccination.** *Medical indications:* chronic disorders of the pulmonary system (excluding asthma); cardiovascular diseases; diabetes mellitus; chronic liver diseases, including liver disease as a result of alcohol abuse (e.g., cirrhosis); chronic renal failure or nephrotic syndrome; functional or anatomic asplenia (e.g., sickle cell disease or splenectomy); immunosuppressive conditions (e.g., congenital immunodeficiency, HIV infection, leukemia, lymphoma, multiple myeloma, Hodgkins disease, generalized malignancy, or organ or bone marrow transplantation); chemotherapy with alkylating agents, antimetabolites, or long-term systemic corticosteroids; or cochlear implants. *Geographic/other indications:* Alaska Natives and certain American Indian populations. *Other indications:* residents of nursing homes and other long-term-care facilities (see *MMWR* 1997;46[No. RR-8] and *MMWR* 2003;52:739–40).
- 4. Revaccination with pneumococcal polysaccharide vaccine.** One-time revaccination after 5 years for persons with chronic renal failure or nephrotic syndrome; functional or anatomic asplenia (e.g., sickle cell disease or splenectomy); immunosuppressive conditions (e.g., congenital immunodeficiency, HIV infection, leukemia, lymphoma, multiple myeloma, Hodgkins disease, generalized malignancy, or organ or bone marrow transplantation); or chemotherapy with alkylating agents, antimetabolites, or long-term systemic corticosteroids. For persons aged ≥ 65 years, one-time revaccination if they were vaccinated ≥ 5 years previously and were aged < 65 years at the time of primary vaccination (see *MMWR* 1997;46[No. RR-8]).
- 5. Hepatitis B vaccination.** *Medical indications:* hemodialysis patients or patients who receive clotting factor concentrates. *Occupational indications:* health-care workers and public-safety workers who have exposure to blood in the workplace; and persons in training in schools of medicine, dentistry, nursing, laboratory technology, and other allied health professions. *Behavioral indications:* injection-drug users; persons with more than one sex partner during the previous 6 months; persons with a recently acquired sexually transmitted disease (STD); all clients in STD clinics; and men who have sex with men. *Other indications:* household contacts and sex partners of persons with chronic hepatitis B virus (HBV) infection; clients and staff members of institutions for the developmentally disabled; inmates of correctional facilities; or international travelers who will be in countries with high or intermediate prevalence of chronic HBV infection for > 6 months (<http://www.cdc.gov/travel/diseases/hbv.htm>) (see *MMWR* 1991;40[No. RR-13]).
- 6. Hepatitis A vaccination.** *Medical indications:* persons with clotting factor disorders or chronic liver disease. *Behavioral indications:* men who have sex with men or users of illegal drugs. *Occupational indications:* persons working with hepatitis A virus (HAV)-infected primates or with HAV in a research laboratory setting. *Other indications:* persons traveling to or working in countries that have high or intermediate endemicity of hepatitis A. If the combined Hepatitis A and Hepatitis B vaccine is used, administer 3 doses at 0, 1, and 6 months (<http://www.cdc.gov/travel/diseases/hav.htm>) (see *MMWR* 1999;48[No. RR-12]).
- 7. Measles, mumps, rubella (MMR) vaccination.** *Measles component:* adults born before 1957 can be considered immune to measles. Adults born during or after 1957 should receive ≥ 1 dose of MMR unless they have a medical contraindication, documentation of ≥ 1 dose, or other acceptable evidence of immunity. A second dose of MMR is recommended for adults who 1) were recently exposed to measles or in an outbreak setting, 2) were previously vaccinated with killed measles vaccine, 3) were vaccinated with an unknown vaccine during 1963–1967, 4) are students in postsecondary educational institutions, 5) work in health-care facilities, or 6) plan to travel internationally. *Mumps component:* 1 dose of MMR vaccine should be adequate for protection. *Rubella component:* Administer 1 dose of MMR vaccine to women whose rubella vaccination history is unreliable and counsel women to avoid becoming pregnant for 4 weeks after vaccination. For women of childbearing age, regardless of birth year, routinely determine rubella immunity and counsel women regarding congenital rubella syndrome. Do not vaccinate pregnant women or those planning to become pregnant during the next 4 weeks. For women who are pregnant and susceptible, vaccinate as early in the postpartum period as possible (see *MMWR* 1998;47[No. RR-8] and *MMWR* 2001;50:1117).
- 8. Varicella vaccination.** Recommended for all persons lacking a reliable clinical history of varicella infection or serologic evidence of varicella zoster virus (VZV) infection who might be at high risk for exposure or transmission. This includes health-care workers and family contacts of immunocompromised persons; persons who live or work in environments where transmission is likely (e.g., teachers of young children, child care employees, and residents and staff members in institutional settings); persons who live or work in environments where VZV transmission can occur (e.g., college students, inmates, and staff members of correctional institutions, and military personnel); adolescents aged 11–18 years and adults living in households with children; women who are not pregnant but who might become pregnant; and international travelers who are not immune to infection.
Note: Approximately 95% of U.S.-born adults are immune to VZV. Do not vaccinate pregnant women or those planning to become pregnant during the next 4 weeks. For women who are pregnant and susceptible, vaccinate as early in the postpartum period as possible (see *MMWR* 1999;48[No. RR-6]).
- 9. Meningococcal vaccine (quadrivalent polysaccharide for serogroups A, C, Y, and W 135).** *Medical indications:* adults with terminal complement component deficiencies or those with anatomic or functional asplenia. *Other indications:* travelers to countries in which meningococcal disease is hyperendemic or epidemic (e.g., the "meningitis belt" of sub-Saharan Africa and Mecca, Saudi Arabia). Revaccination after 3–5 years might be indicated for persons at high risk for infection (e.g., persons residing in areas where disease is epidemic). Counsel college freshmen, especially those who live in dormitories, regarding meningococcal disease and availability of the vaccine to enable them to make an educated decision about receiving the vaccination (see *MMWR* 2000;49[No. RR-7]). The American Academy of Family Physicians recommends that colleges should take the lead on providing education on meningococcal infection and availability of vaccination and offer it to students who are interested. Physicians need not initiate discussion of meningococcal quadrivalent polysaccharide vaccine as part of routine medical care.

Appendix D

Prevention and Treatment of Influenza with Antiviral Drugs*

People at high risk for serious influenza illness

- ✓ 65 years or older
- ✓ Nursing home patients
- ✓ Long-term health problems. Examples:
 - Heart disease
 - Lung disease, including asthma
 - Metabolic disease, like diabetes mellitus
 - Blood diseases (e.g. hemoglobinopathies like sickle cell, thalassemia)
 - End-stage renal disease
 - Cognitive, neurologic, or other conditions that lead to increased risk for aspiration of secretions
- ✓ Weakened immune system. Examples:
 - Cancer or cancer treatment
 - HIV/AIDS
 - Steroid therapy
- ✓ Pregnant (or plan to be pregnant during the influenza season—through spring 2005)

When to use drugs to *treat* influenza

- ✓ Potentially life-threatening influenza-related illness
OR
 - ✓ At high risk for influenza complications (see box)
AND
- Within the first two days of illness onset
- ✓ *Note: pregnant women should consult their primary health care provider*

When to use drugs to *prevent* influenza during periods of influenza activity in the community

- ✓ Unvaccinated staff and patients
 - In a hospital or nursing home outbreak
OR
 - At high risk for influenza complications
- ✓ Vaccinated patients
 - Immunosuppressed AND not expected to respond well to the vaccine (e.g. advanced HIV disease)
OR
 - Got the vaccine recently BUT are waiting the 2 weeks for it to give protection

During an institutional outbreak, chemoprophylaxis should be administered to all residents regardless of prior vaccinations, and should continue for a minimum of 2 weeks.

Health-care workers and children

- ✓ Special recommendations apply
- ✓ VA staff should see occupational health for work-related illness or their own health care providers

Which drugs for *prevention*?

- ✓ Amantadine (Symmetrel®) 100 mg bid is preferred, but
 - Has more GI & CNS effects
 - Important to reduce dose for renal insufficiency or age 65 or older
 - Reduce dose to 100 mg po qd for age \geq 65. Consult drug package insert for dosing when creatinine clearance \leq 50 mL/min.
- ✓ Alternate: Rimantadine (Flumadine®) 100 mg bid
 - Fewer side effects than amantadine
 - Reduce dose to 100 mg po qd for age \geq 65, creatinine clearance \leq 10 mL/min., or severe liver disease

Which drug for *treatment*?

- ✓ Oseltamivir (Tamiflu®) 75 mg po bid x 5 days
- ✓ Reduce dose if creatinine clearance $<$ 30 mL/min,
- ✓ May cause nausea, vomiting
- ✓ Contact your local pharmacy or Pharmacy Benefits Management if oseltamivir is needed
- ✓ Amantadine and rimantadine are also approved for treatment of influenza illness, but use of these drugs has been associated with the development of resistant influenza virus. See CDC recommendations for additional advice and dosing.

*Document adapted from Ineck, B., Rubingh, C. and Bittner, M.J. Therapeutic Bulletin, Special Report, *Anti-Influenza Drug Update*. November, 2004. VA Nebraska-Western Iowa Health Care System. Used with permission.

Reviewed by Matthew Goetz, M.D., VA Greater Los Angeles Health Care System and Member of the Pharmacy Benefits Management Medical Advisory Panel.

The CDC recommendations are the authoritative source for this document and may be found at (<http://www.cdc.gov/flu/professionals/treatment/0405antiviralguide.htm>)

Appendix E

Resources, References, and Web sites

Resources

This VA Influenza Toolkit 2005-2006 is available on the VA Internet sites www.publichealth.va.gov/flu and www.publichealth.va.gov/InfectionDontPassItOn, and the VA Intranet sites <http://vaww.vhaco.va.gov/phshcg/Flu> and <http://vaww.vhaco.va.gov/phshcg/InfectionDontPassItOn>).

In addition, the following VA resources and more are available at <http://www.nchpdp.med.va.gov/2004FluToolkitDownloads.asp>

- VA Influenza/Pneumococcal Resource Toolkit 2004-2005
- Toolkit Durham Immunization Reminder Dialogs
- Sample Generic Policy – Administration of Vaccines
- Sample Generic Policy – Outpatient Administration of Vaccines
- Sample VHA Pneumococcal Vaccine Protocol Checklist
- Sample VHA Influenza Vaccine Information and Protocol
- Staff Reminder Notices
- Telephone Script

National Influenza Vaccine Summit's "Influenza Vaccination Pocket Information Guide" is available at <http://www.immunize.org/fluguide/pocketguide.pdf>

References

Guidance on influenza immunization

Prevention and Control of Influenza: Recommendations of the Advisory Committee on Immunization Practices (ACIP), MMWR, July 29, 2005. Vol. 54. Available at <http://www.cdc.gov/mmwr/preview/mmwrhtml/rr5408a1.htm>

CDC. Update: Influenza activity--United States and worldwide, 2004-05 season. MMWR Morb Mortal Wkly Rep. 2005 Jul 1;54(25):631-4. Available at <http://www.cdc.gov/mmwr/preview/mmwrhtml/mm5425a3.htm>.

VHA. VHA Directive: Influenza Vaccine-Recommendations for 2005-2006, publication pending.

VHA. VA Influenza Advisory #1 for the 2005-2006 Influenza Season, May 9, 2005. Available at <http://www.publichealth.va.gov/flu> and <http://www.publichealth.va.gov/infectiondontpassiton>.

Guidance on immunization/vaccination in general

CDC. Prevention and Control of Influenza: Recommendations of the Advisory Committee on Immunization Practices (ACIP), MMWR, July 29, 2005. Vol. 54. Available at <http://www.cdc.gov/mmwr/preview/mmwrhtml/rr5408a1.htm>



CDC. Recommended Adult Immunization Schedule --- United States, October 2004--September 2005. November 19, 2004 / 53(45);Q1-Q4
<http://www.cdc.gov/mmwr/preview/mmwrhtml/mm5345-Immunizational1.htm>

Vaccine information statements (VISs)

CDC. Inactivated Influenza Vaccine information statement (VIS). Atlanta, GA: US Department of Health and Human Services, CDC, 2005 (interim version as of July 18, 2005). Available at: <http://www.cdc.gov/nip/publications/VIS/default.htm#flu>

CDC. Live, Intranasal Influenza Vaccine information statement (VIS). Atlanta, GA: US Department of Health and Human Services, CDC, 2005 [interim version as of July 18, 2005). Available at <http://www.cdc.gov/nip/publications/VIS/default.htm#flu>.

NOTE FROM CDC: The two Influenza VISs above are "interim" editions. Because influenza was recently added to the National Vaccine Injury Compensation Program, federal law requires a notice and comment process to develop "final" versions of both documents. These may not be available until flu season has begun. Pending completion of the final editions, providers are encouraged to use these interim versions.

CDC. Pneumococcal Polysaccharide (PPV23) Vaccine information statement (VIS). Atlanta, GA: US Department of Health and Human Services, CDC, 7/29/97
<http://www.cdc.gov/nip/publications/VIS/default.htm#ppv23>

Vaccination of health care workers

Carman WF, et al. "Effects of influenza vaccination of health-care workers on mortality of elderly people in long-term care: a randomized controlled trial." Lancet 2000;355:93-97.

CDC. "Immunization of Health Care Workers: Recommendations of the Advisory Committee on Immunization Practices (ACIP) and the Hospital Infection Control Practices Advisory Committee (HICPAC)," MMWR, December 26, 1997/46: 1-42.
<http://www.cdc.gov/mmwr/preview/mmwrhtml/00050577.htm> (web version) and
<http://www.cdc.gov/mmwr/PDF/rr/rr4618.pdf> (PDF File)

D'Heilly SJ, Nichol KL. Work-site-based influenza vaccination in health care and non-health care settings. Infect Control Hosp Epidemiol. 2004 Nov;25(11):941-5.

National Foundation for Infectious Diseases. "Influenza immunization among health care workers: Call to action," 2004. Available at <http://www.nfid.org/>

National Foundation for Infectious Diseases. "Improving influenza vaccination rates in health care workers," 2004. Available at <http://www.nfid.org/>

Vaccination of veterans, patients and the public

VA. “Maximizing Vaccination Rates for Veterans with SCI&D, VA QUERI Quarterly Newsletter. Vol 3:No 4: March 2002. http://www.hsrdr.research.va.gov/publications/queri_quarterly or PDF for this issue <http://www.hsrdr.research.va.gov/publications/internal/QUERIvol3no4Spring.pdf>

CDC. “Vaccine-Preventable Diseases: Improving Vaccination Coverage in Children, Adolescents, and Adults (TFCPS),” MMWR, June 18, 1999/48: 1-15.
<http://www.cdc.gov/mmwr/preview/mmwrhtml/rr4808a1.htm> (web version) and
<http://www.cdc.gov/mmwr/PDF/rr/rr4808.pdf> (PDF File)

CDC. “Adult Immunization Programs in Nontraditional Settings: Quality Standards and Guidance for Program Evaluation” and “Use of Standing Orders Programs to Increase Adult Vaccination Rates (APIC),” MMWR, March 24, 2000/49: 1-26.
<http://www.cdc.gov/mmwr/preview/mmwrhtml/rr4901a1.htm> (web version) and
<http://www.cdc.gov/mmwr/preview/mmwrhtml/rr4901a1.htm> (PDF File)

CDC. “Influenza vaccination coverage among adults aged >50 years and pneumococcal vaccination coverage among adults aged >65 years—United States, 2002,” MMWR 2003;52(41):987-992.
Immunization Action Coalition. “Adults only vaccination: a step-by-step guide,” 2004. Available at www.immunize.org

Jacobson VJ, Szilagyi P. Patient reminder and patient recall systems to improve immunization rates. Cochrane Database Syst Rev. 2005 Jul 20;(3):CD003941. Abstract and synopsis available at <http://www.mrw.interscience.wiley.com/cochrane/clsysrev/articles/CD003941/frame.html>

Nichol KL. “Benefits of Influenza Vaccination for Low-, Intermediate-, and High-Risk Senior Citizens.” Archives of Internal Medicine, 1998;158: 1769.

Nichol KL. “Ten-Year Durability and Success of an Organized Program to Increase Influenza and Pneumococcal Vaccination Rates Among High-Risk Adults.” American Journal of Medicine, 1998;105:385-92.

Nichol KL. “Influenza Vaccination for Healthy Working Adults.” Minnesota Medicine, November 1999; Volume 82 <http://www.mnmed.org/publications/MnMed1999/November/Nichol.cfm>

Nichol KL. “Influenza Vaccination and Reduction in Hospitalizations for Cardiac Disease and stroke Among the Elderly,” New Engl J Med, 2003;348:1322-1332.

Nichol KL, D’Heilly S, Ehlinger, E. Colds and influenza-like illnesses in university students: impact on health, academic and work performance, and health care use.
Clin Infect Dis. 2005 May 1;40(9):1263-70. Epub 2005 Mar 31.

Polarid GA, et al. “Standards for Adult Immunization Practices,” Am J Prev Med 2003;25:144-150.

Szilagyi PG, et al. “Effect of Patient Reminder/Recall Interventions on Immunization Rates,” JAMA 2000;284 1820-1827. <http://jama.ama-assn.org/issues/v284n14/rfull/jrv00030.html> (web version) or
<http://jama.ama-assn.org/issues/v284n14/rpdf/jrv00030.pdf> (PDF File)

Stone E. et al. “Interventions That Increase Use of Adult Immunization and Cancer Screening Services: A Meta-Analysis,” Annals of Internal Medicine, 2002;136:641-651.
<http://www.annals.org/issues/v136n9/full/200205070-00006.html> (web version) or
<http://www.annals.org/issues/v136n9/pdf/200205070-00006.pdf> (PDF File)

Cost effectiveness of influenza vaccination

Nichol KL. The efficacy, effectiveness and cost-effectiveness of inactivated influenza virus vaccines. Vaccine. 2003 May 1;21(16):1769-75.

Nichol KL, Goodman M. Cost effectiveness of influenza vaccination for healthy persons between ages 65 and 74 years. Vaccine. 2002 May 15;20 Suppl 2:S21-4.

Nichol KL. Cost-benefit analysis of a strategy to vaccinate healthy working adults against influenza. Arch Intern Med. 2001 Mar 12;161(5):749-59.

Nichol KL. Influenza vaccination in the elderly: impact on hospitalisation and mortality. Drugs Aging. 2005;22(6):495-515.

Pneumococcal vaccination

CDC. “Guidelines for preventing health-care-associated pneumonia, 2003: recommendations of CDC and the Health care Infection Control Practices Advisory Committee,” MMWR/March 26, 2004; 53(RR03):1-35. <http://www.cdc.gov/mmwr/preview/mmwrhtml/rr5303a1.htm> (web version) and <http://www.cdc.gov/mmwr/PDF/rr/rr5303.pdf> (PDF File)

CDC. “Prevention of Pneumococcal Disease: Recommendations of the Advisory Committee on Immunization Practices (ACIP),” MMWR, April 4, 1997/46: 1-23.
<http://www.cdc.gov/mmwr/preview/mmwrhtml/00047135.htm> (web version) and
<http://www.cdc.gov/mmwr/PDF/rr/rr4608.pdf> (PDF File)

CDC. “Influenza vaccination coverage among adults aged >50 years and pneumococcal vaccination coverage among adults aged >65 years—United States, 2002,” MMWR 2003;52(41):987-992.

CDC. Questions About the Pneumococcal Vaccine brochure. Available at <http://www.cdc.gov/nip/vaccine/pneumo/pneumo-pubs.htm#brochure>.

Lifson AR, Aitchison-Olson R, & Ramesh A. “New Threats from an Old Enemy: A Physician Update on Pneumococcus,” Minnesota Medicine, November 1999;82.
<http://www.mnmed.org/publications/MnMed1999/November/Lifson.cfm>

Sisk J. et al. Cost-effectiveness of vaccination against pneumococcal bacteremia among elderly people. JAMA, 1997;278:1333.

Web sites

<http://www.publichealth.va.gov/flu/> (VA Intranet vaww.publichealth.va.gov/flu/) - This is the influenza Web site for the Department of Veterans Affairs. It includes links on the flu and flu vaccine, VA advisories on influenza vaccine as information evolves during flu vaccine season, and the VA influenza toolkit, a set of materials to enable VA facilities to put on flu immunization campaign.

<http://www.publichealth.va.gov/infectiondontpassiton> (VA Intranet <http://vaww.vhaco.va.gov/phshcg/InfectionDontPassItOn>) - This is the Web site for the VA public health campaign “Infection: Don’t Pass It On,” which focuses on prevention of infection in the VA medical system through hand and respiratory hygiene, material for infection emergencies, and vaccination against influenza and pneumonia.

<http://www.nchpdp.med.va.gov/> - This Web site of the VA National Center for Health Promotion and Disease Prevention includes the 2004-2005 influenza toolkit.

<http://www.cdc.gov/nip> - This is the Web site for the National Immunization Program of the Centers for Disease Control and Prevention (CDC) and has a great deal of information for both the public and health care providers on all immunization topics.

<http://www.cdc.gov/nip/publications/ACIP-list.htm> - This page on the NIP site lists all recommendations of the ACIP (Advisory Committee for Immunization Practices).

<http://www.cdc.gov/nip/recs/adult-schedule.htm> - This page includes a printable schedule of adult immunization recommendations, a list of vaccines for adults, and an adult vaccination screening form.

<http://www.cdc.gov/nip/publications/adultstrat.htm> - This page includes strategies for Increasing Adult Vaccination Rates (NIP), March 8, 2002.

<http://www.cdc.gov/flu/> - This is the main influenza Web page of the CDC. It includes extensive information about the disease of influenza and its prevention and control, for both patients and health care professionals.

<http://www.immunize.org> - This is the Web site for the Immunization Action Coalition (IAC) with a wide variety of information about immunizations, including Vaccine Information Statements in many languages. The Directory of Immunization Resources is full of useful information on organizations, Web sites, hotlines, and agencies that are immunization resources.

<http://www.vaccineinformation.org> - This page from the IAC is comprehensive, organized, and easy to access. For each vaccine-preventable disease, there are answers to many questions about the disease and the vaccine, as well as sections containing photos, case histories, recommendations, references, and links to useful resources. Also included is material about vaccine safety, travel, bioterrorism, state laws—and much more.

<http://www.acponline.org/aij> - This site from the American College of Physicians provides resources and tools to support physicians in their immunization efforts, with the goal of improving adult immunization rates. It includes physician education, patient education, and practice management tools for immunization and reimbursement.

<http://www.nfid.org/> - This is the Web site for the National Foundation for Infectious Diseases and contains a call to action and strategies for increasing influenza immunization among health care workers.

<http://www.nfid.org/ncai> - This is the Web site for the National Coalition for Adult Immunization and includes fact sheets, immunization schedules, recommendations, and report cards, and an order form for a “National Adult Immunization Awareness Week” campaign kit.

<http://www.vaccines.org> - This Web site provides access to up-to-the-minute news about vaccines and an annotated database of vaccine resources on the Internet.

<http://www.cms.hhs.gov/preventiveservices/2.asp> - This is a Web page of the Centers for Medicare and Medicaid Services (CMS) that describes the CMS Adult Immunization Project, which focuses on increasing rates of preventive vaccination against influenza and pneumococcal disease for Medicare beneficiaries across all health settings.

<http://www.ImmunizationEd.org> - This is a Web page from the Society of Teachers of Family Medicine that provides news and reports to keep family physicians up-to-date on vaccines for children and adults, links to the most current immunization schedules and vaccine information, downloadable slide presentations and photographs of diseases.

<http://www.atpm.org> - This Web site of the Association of Teachers of Preventive Medicine has several educational resources available for download or purchase for training health care professionals and students about immunization issues.

<http://www.naccho.org> - This is the Web site of the National Association of County and City Health Officials and has several pages of vaccine information, with links to training and resources pages.

<http://www.partnersforimmunization.org> - This Web site of the National Partnership for Immunization, a non-profit organization dedicated to reducing the nationwide incidence of vaccine-preventable diseases through increased use of licensed vaccines, funded, in part, by the Centers for Disease Control and Prevention, is a good source for immunization resources.

<http://www.nlm.nih.gov/medlineplus/influenza.html> - This is the influenza Web page of Medline Plus, a service of the National Library of Medicine, National Institutes of Health (NIH). It includes sections on news, diagnosis, treatment, prevention, disease management, clinical trials and other research, and information focused on audiences ranging from children to the elderly.

<http://www.mayoclinic.com/invoke.cfm?objectid=5CB89570-8B46-4961-8BFE66D06D5BDD1B> - This is the Mayo Clinic patient information page on influenza.

<http://www.health.state.mn.us/divs/idepc/diseases/flu/index.html> - This is the influenza section of the Minnesota Department of Health.

Appendix F Acknowledgements

The team of VHA offices and staff that created the 2005-2006 Influenza Toolkit and Toolkit Manual would like to acknowledge the VA National Center for Health Promotion and Disease Prevention and Dr. Kristin Nichol, Minneapolis VA Medical Center, for their leadership of past toolkits and their expertise and dedication to flu prevention efforts in the VA medical system.

Coordinating Offices

Office of Public Health & Environmental Hazards

Victoria Davey, Deputy Chief Consultant, Public Health Strategic Health Care Group
 Pamela Hirsch, Clinical Program Manager, Occupational Health Program
 V. Troy Knighton, Director, Training & Education, Public Health Strategic Health Care Group
 Connie Raab, Director, Public Health Communications
 Donna Wells, National Prevention Coordinator, Public Health Strategic Health Care Group



VA Employee Education System

Pam Hebert, Patient Health Education Coordinator, EES Birmingham




VA Infection Control Professionals

Kathleen DeRoos, Nurse Practitioner/Infection Control, Omaha
 Beverly V. Johnson, Clinical Programs Coordinator, Martinsburg
 Vicki Macks, Infection Control Nurse, Memphis
 Gina Oda, Microbiologist/Infection Control Practitioner, Palo Alto
 Mary Standridge, Infection Control Nurse, Memphis




VA National Center for Patient Safety

Noel Eldridge, Executive Assistant



Infectious Diseases Program Office

Linda Danko, Clinical Program Coordinator/Infectious Diseases Program, VA Central Office



VA National Center for Health Promotion and Disease Prevention

Linda Kinsinger, Assistant Director, Policy, Programs, Training, and Education

Kristy Straits-Troster, Assistant Director, Infectious Disease Prevention

Rosemary Strickland, Acting Assistant Director, Field Operations

