

LPN Dosage Calculation Study Guide



Minneapolis VA Medical Center



Unit 1
Metric Conversions
Study sheet RN and LPN

Medications are measured in one of these 6 metric units:

Measures of weight:

kg (kilogram)
gm or g (gram)
mg (milligram)
mcg or μg (microgram)

Measures of Volume:

ml (milliliters) or cc (cubic centimeters)
these are interchangeable
L (liter)

Note: the unit mEq (milliequivalents) is a chemical, not a metric, unit of measure.

Metric conversion is used when the 2 parts of a problem are stated in different metric units. For example:

The doctor ordered Atropine 400 **mcg** IM.

The bottle reads Atropine .4 **mg** per ml.

To calculate how much to give, you must first convert mcg to mg OR mg to mcg. In other words, the two units of measure must match.

To convert the units you need to know the following equations:

Weight

1mg = 1000 mcg
1gm = 1000 mg
1 kg = 1000 gm

Volume

1000ml = 1 L

The metric system is a decimal system. Note that for all of the equations used, only the numbers 1 and 1000 are used. To convert, you multiply or divide by 1000. This can be done by simply moving the decimal point.

To convert from a smaller unit to a larger unit, you divide by 1000 or move the decimal point 3 spaces to the LEFT. For Example.

$$1500 \text{ mg} = \underline{\quad?} \text{ gm}$$

OR

$$1500 \text{ mg} \div 1000 = 1.5 \text{ gm}$$

Move the decimal point:



To convert from a larger unit to a smaller unit you multiply by 1000 or move the decimal point 3 places to the right. For example:

$$2 \text{ gm} = \underline{\quad?} \text{ mg}$$

$$2 \text{ gm} \times 1000 = 2000 \text{ mg}$$

OR

move the decimal point:



Now try a few conversions:

1. 0.5 gm = _____ mg

2. 500 ml = _____ L

3. 600 mg = _____ gm

4. 5 mcg = _____ mg

5. 100 mg = _____ gm

6. 400 mcg = _____ mcg

7. 400 mcg = _____ mg

8. .3 gm = _____ mg

9. 0.25 gm = _____ mg

10. 500 mcg = _____ mg

11. 5 mcg = _____ mg

12. 0.5 L = _____ ml

13. 100 mcg = _____ mg

14. 1000mcg = _____ mg

15. 5.5 mg = _____ gm

16. 25mg = _____ gm

17. 0.05 gm = _____ mg

18. 0.2 mg = _____ mcg

Memory tip: when converting **to** a **L**arger unit, move decimal **L**eft (both start with L)
When converting **to** a smaller unit, move decimal right.

Answers:

1. 500mg	10. .5 mg
2. .5 L	11. .005 mg
3. .6 gm	12. 500 ml
4. .005 mg	13. .1 mg
5. .1 gm	14. 1 mg
6. 400,000 mcg	15. .0055 gm
7. .4 mg	16. .025 gm
8. 300 mg	17. 50 mg
9. 250 mg	18. 200 mcg

Unit II
Dosage Calculation
Study Sheet RN and LPN

A simple proportion formula can be used to calculate any dosage.

This formula will work every time, with all forms of medication, if set up correctly.
Always follow these steps:

- 1.) Set up the proportion: Put what you know on the left of the equal sign (=) and what you need to know on the right of the equal (=) sign.

For example: You know a medication comes in 250 mg tablets, and want to give 750 mg. To find out how many tablets to give, the set up is:

$$\begin{array}{ccc} \textit{YOU KNOW} & & \textit{YOU WANT} \\ \\ \frac{250 \text{ mg}}{1 \text{ tablet}} & = & \frac{750 \text{ mg}}{X \text{ tablets}} \end{array}$$

- 2.) Cross multiply: Multiply the numerator of each side by the denominator of the other side.

$$\begin{array}{ccc} \frac{250 \text{ mg}}{1 \text{ tablet}} & \begin{array}{c} \swarrow \quad \searrow \\ \quad \quad \quad \end{array} & \frac{750 \text{ mg}}{X \text{ tablets}} \end{array} \quad \text{which is} \quad 250 \cdot X = 750$$

- 3.) Isolate X: Divide both sides of the equation by the number on the "X" side.

$$\begin{array}{ccc} \frac{\cancel{250} X}{\cancel{250}} & = & \frac{750}{250} \\ \\ X & = & \frac{750}{250} \end{array}$$

- 4.) Solve for X: Divide the numbers on the right side of the equation.

$$\begin{array}{l} X = 750 \text{ divided by } 250 \\ X = 3 \text{ tablets} \quad \text{therefore, give 3 tablets} \end{array}$$

Note: A proportional formula is only used when the answer is not readily apparent. Obviously, this one could have been done "in your head".

Let's try another. A liquid medication comes in 30 mg per 5 ml. You want to give 300 mg.

1.) Set up proportion:

$$\begin{array}{r} \text{YOU KNOW} \\ \hline 30 \text{ mg} \\ \hline 5 \text{ ml} \end{array} = \begin{array}{r} \text{YOU WANT} \\ \hline 300 \text{ mg} \\ \hline X \text{ ml} \end{array}$$

2.) Cross multiply:

$$\begin{array}{r} 30 X \\ \hline 30 X \end{array} = \begin{array}{r} 300 \cdot 5 \\ \hline 1500 \end{array}$$

3.) Isolate X:

$$\begin{array}{r} \cancel{30} X \\ \hline \cancel{30} \end{array} = \begin{array}{r} 1500 \\ \hline 30 \end{array}$$
$$X = \frac{1500}{30}$$

4.) Solve for X:

$$X = 1500 \div 30$$
$$X = 50 \quad \text{therefore, give 50 ml}$$

Now try some on your own. You may have to do metric conversion first (see Unit I), to be sure the units of measure match.

1.) The order is for .25 mg of Digoxin. You have an ampule labeled 500 mcg per 2 ml. How many ml's will you give?

2.) The order is for Dilantin liquid 150 mg per NG tube. You have a bottle with 30 mg per 5 ml. How many ml's will you give?

- 3.) The order is for 50 mg of a drug. The dose available in stock is 25 mg. How many tablets will you give?

- 4.) The order is for 15 mEq of KCl per NG tube. The bottle contains 20 mEq per 15 ml. How many ml's will you give?

- 5.) The order is for 500 mg of a drug. The dose available is 200mg per 10 ml. How many ml's will you give?

- 6.) The order is for Prednisone 12.5 mg. You have 2.5 mg tablets of Prednisone. How many tablets will you give?

- 7.) The order is for Lactulose 25 gm. The bottle contains 10 gm. per 15 ml. How many ml's will you give?

- 8.) The order is for Digoxin 250 mcg. You have tablets of 0.25 mg of Digoxin. How many tablets will you give?

Answers:
1. 1ml 3. 2 tablets 5. 25ml 7. 37.5 ml
2. 25ml 4. 11.25 ml 6. 5 tablets 8. 1 tablet

Unit III
 Dosage Calculations for Injectables
 Study Sheet RN and LPN

The proportional formula found in Unit II works just the same for injectable drug calculations. Here's an example: 8 mg of morphine is ordered. You have a supply of 10 mg per ml morphine.

1.) Set up proportion:
$$\frac{10 \text{ mg}}{1 \text{ ml}} = \frac{8 \text{ mg}}{X \text{ ml}}$$

2.) Cross multiply:
$$10 X = 8$$

3.) Isolate X:
$$\frac{\cancel{10} X}{\cancel{10}} = \frac{8}{10} \quad X = \frac{8}{10}$$

4.) Solve for X:
$$X = 8 \div 10 \quad X = .8 \text{ ml}$$

If you have an injectable powdered drug to which you are going to add a diluent, the directions on the vial tell you how much diluent to add. The vial will also state the concentration of the drug after the diluent is mixed. It is this concentration that is used in the proportional formula for dosage calculation. The amount of diluent added does not figure into the dosage calculation.

For example: The order is for 1 gm of Cefadyl (Cephapirin). The directions on the vial read: "Add 2 ml sterile water for injection. Each 1.2 ml contains 500 mg. of cephalapirin. How many ml's do you give?"

First, convert 500 mg to .5 gm, so that the units match.

Then set up the proportion, and solve, using the concentration given on the vial.

$$\frac{.5 \text{ gm}}{1.2 \text{ ml}} = \frac{1 \text{ gm}}{X \text{ ml}}$$

$$.5 X = 1.2$$

$$X = 2.4 \text{ ml}$$

Here are some practice problems for injectable dosage calculation.

1.) The order is for 2 gm of the drug. The vial reads: Add 8.6 ml of diluent to contents of vial. Each ml will contain 500 mg. How many ml's will you give?

- 2.) You have an ampule of Digoxin labeled 500 mcg/2 ml. The order is for .25 mg of Digoxin. How many ml's will you give?
- 3.) The order is for Haloperidol 2 mg. It is supplied in a 5 mg/ml vial. How many ml's will you give?
- 4.) The order is for 0.5 gm Amoxicillin. The directions read: Add 3.5 ml of diluent. Resulting solution contains 250 mg Amoxicillin per ml. How many ml's will you give?
- 5.) The order is for naloxone (Narcan) 0.3 mg. The cartridge is labeled 0.4 mg/ml. How many ml's will you give?
- 6.) The order is for furosemide (Lasix) 20 mg. The vial is labeled 100mg/10 ml. How many ml's will you give?
- 7.) The order is for penicillin G 300,000 U. The vial directs you to add 4.2 ml of normal saline to make a concentration of 3 million units per 5 ml. How many ml will you give?

Answers:

1. 4 ml
2. 1 ml
3. .4 ml
4. 2 ml
5. .75 ml
6. 2 ml
7. .5 ml

Unit IV
Calculation of IV Drip Rates Study Sheet
LPN's with 3+ years of experience only

Please note: You will only be asked to do this part of the drug calculation test if you have 3 or more years of LPN experience.

When you calculate the IV drip rate, you are calculating the **number of drops per minute** at which you will administer IV fluids.

To do this calculation you need three pieces of information. They are:

- A. VOLUME (in ml or cc)
- B. DROP FACTOR OF THE IV TUBING (found on tubing package)
- C. TIME TO RUN (In minutes, for the volume being administered)

Using these three numbers, the formula for calculating the drip rate is:

$$\frac{\text{A. (VOLUME)} \cdot \text{B. (DROP FACTOR)}}{\text{C. (MINUTES for above volume to run)}} = \text{DROPS PER MINUTE}$$

Most tubing used at the MVAMC have drop factors of either:

15 drops/ml: "macrodrop" or "regular" tubing

60 drops/ml: "minidrip" tubing

Always check the drop factor on the tubing package.

EXAMPLES:

A. The IV order is for D₅W to run at 125 ml/hour.

Using regular macrodrip tubing, the problem is set up as follows:

$$\frac{125 \text{ (volume)} \cdot 15 \text{ (tubing drop factor)}}{60 \text{ (minutes)}} = \frac{1875}{60} = 31+ \text{ drops/minute}$$

OR if you reduce the numbers:

$$\frac{125 \cdot \cancel{15^1}}{\cancel{60}_4} = \frac{125}{4} = 31+ \text{ drops/minute}$$

B. The hourly rate of the IV is 90 ml. Using minidrip tubing:

$$\frac{90 \cdot 60}{60} = \frac{5400}{60} = 90 \text{ drops/min} \quad \text{OR} \quad \frac{90 \cdot \cancel{60^1}}{\cancel{60}_1} = 90$$

Note that in this example, you are taking an hourly volume of 90, multiplying by 60 and then dividing the product by 60, which is unnecessary. When using an **hourly** volume and **minidrip** tubing, the hourly volume and the drops per minute are the same.

C. You are to give an antibiotic in 50 ml D₅W with minidrip tubing to run over one hour.

$$\frac{50 \cdot 60}{60} = 50 \text{ drops per minute} \quad \text{OR} \quad \frac{50 \cdot \cancel{60^1}}{\cancel{60}_1} = 50$$

As noted in example "B", the drops per minute and the ml per hour are the same.

D. However, if you were to run the previous piggyback in 30 minutes, your calculation would change to:

$$\frac{50 \cdot 60}{30} = \frac{3000}{30} = 100 \text{ drops/min} \quad \text{OR} \quad \frac{50 \cdot \cancel{60^2}}{\cancel{30}_1} = 100$$

Remember to always adjust the minutes to the time ordered.

Now try a few on your own:

1. The order is for 1000 ml of D₅½NS with 20 mEq KCl to run over 8 hours.

- The hourly rate for this IV is _____ .
- Using macrodrip tubing (15 drops/ml), the drip rate is _____ .
- If you used minidrip tubing (60 drops/ml), the drip rate is _____ .

2. The order is for 1 liter of D₅NS to run over 10 hours.
 - a. The hourly rate is _____ .
 - b. Using regular tubing (15 drops/ml), the drip rate is _____ .
 - c. Using minibidrip tubing (60 drops/ml), the drip rate is _____ .

3. The order is for Vancomycin 1000 mg, diluted in 250 ml of NS. You want to run this piggyback over 2 hours.
 - a. Using regular tubing, the drip rate is _____ .
 - b. If you use minibidrip tubing, the drip rate is _____ .

4. 250 ml of packed red blood cells (PRBC's) is to run over 3 hours, and you are monitoring the progress. The blood tubing delivers 15 drops per ml. The drip rate is _____ .

5. An IVPB of Ceftizoxime 1 gm in 50 ml of NS is to run in 30 minutes. If you use minibidrip tubing, the drip rate is _____ .

6. If the IVPB in #5 were to run over 20 minutes, the drip rate is _____ .

Answers:

- | | | | |
|-----------------|----------------|----------------|--------------|
| 1. a. 125 ml/hr | 2. a. 100ml/hr | 3. a. 31 drops | 4. 21 drops |
| b. 31 drops | b. 25 drops | b. 125 drops | 5. 100 drops |
| c. 125 drops | c. 100drops | | 6. 150 drops |

Notes

Notes